

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
12 December 2002 (12.12.2002)


PCT

(10) International Publication Number
WO 02/099084 A2

- (51) International Patent Classification⁷: **C12N**
- (21) International Application Number: **PCT/US02/22272**
- (22) International Filing Date: **4 April 2002 (04.04.2002)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:
0108491.2 **4 April 2001 (04.04.2001)** **GB**
- (71) Applicant (*for all designated States except US*): **SANG-AMO BIOSCIENCES, INC** [US/US]; Point Richmond Tech Center, 501 Canal Boulevard, Suite A100, Richmond, CA 94804 (US).
- (72) Inventors; and
(75) Inventors/Applicants (*for US only*): **Moore, Michael** [GB/GB]; The Hideaway, Fagnall Lane, Winchmore Hill, Amersham, Bucks HP7 0PG (GB). **SEPP, Armin** [EE/GB]; 43 Bullen Close, Cambridge CB1 8YU (GB). **ISALAN, Mark** [GB/GB]; 24 Shottfield Avenue, East Sheen, London, SW14 8EA (GB). **CHOO, Yen** [GR/GB]; 10 Sidney Street, London SW3 6PP (GB).
- (74) Agents: **PASTERNAK, Dahna, S.** et al.; Robins & Pasternak llp, 545 Middlefield Road, Suite 180, Menlo Park, CA 94025 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK,

[Continued on next page]

(54) Title: COMPOSITE BINDING POLYPEPTIDES



(57) Abstract: Disclosed herein are polypeptides with novel DNA binding specificities, constructed from combinations of zinc fingers, and methods for their preparation and use.

WO 02/099084 A2



SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

COMPOSITE BINDING POLYPEPTIDES

TECHNICAL FIELD

5 The present disclosure is in the fields of molecular biology and protein design; in particular, the design of sequence-specific binding proteins for regulation of gene expression.

10 BACKGROUND

Protein-nucleic acid recognition is a commonplace phenomenon that is central to a large number of biomolecular control mechanisms that regulate the functioning of eukaryotic and prokaryotic cells. For instance, protein-DNA interactions form the basis of the
15 regulation of gene expression and are thus one of the subjects most widely studied by molecular biologists.

A wealth of biochemical and structural information explains the details of protein-DNA recognition in numerous instances, to the extent that general principles of recognition
20 have emerged. Many DNA-binding proteins contain independently folded domains for the recognition of DNA, and these domains in turn belong to a large number of structural families, such as the leucine zipper, the "helix-turn-helix" and zinc finger families.

Despite the great variety of structural domains, the specificity of the interactions observed
25 to date between protein and DNA most often derives from the complementarity of the surfaces of a protein α -helix and the major groove of DNA. See, e.g., Klug, (1993) Gene 135:83-92. In light of the recurring physical interaction of α -helix and major groove, the tantalising possibility arises that the contacts between particular amino acids and DNA bases could be described by a simple set of rules; in effect a stereochemical recognition
30 code which relates protein primary structure to binding-site sequence preference.

It is clear, however, that no code will be found which can describe DNA recognition by all DNA-binding proteins. The structures of numerous complexes show significant differences in the way that the recognition α -helices of DNA-binding proteins from different structural families interact with the major groove of DNA, thus precluding
5 similarities in patterns of recognition. The majority of known DNA-binding motifs are not particularly versatile, and any codes which might emerge would likely describe binding to a very few related DNA sequences.

Even within each family of DNA-binding proteins, moreover, it has hitherto appeared
10 that the deciphering of a code would be elusive. Due to the complexity of the protein-DNA interaction, there does not appear to be a simple "alphabetic" equivalence between the primary structures of protein and nucleic acid which specifies a direct amino acid to base relationship.

15 International patent application WO 96/06166 addresses this issue and provides a "syllabic" code that explains protein-DNA interactions for zinc finger nucleic acid binding proteins. A syllabic code is a code that relies on more than one feature of the binding protein to specify binding to a particular base, the features being combinable in the forms of "syllables", or complex instructions, to define each specific contact. Segal,
20 D. J., Dreier, B., Beerli, R. R. & Barbas, C. F. (1999) Proc. Natl. Acad. Sci. USA 96, 2758-2763 present a method of constructing zinc fingers polypeptides, based on 16 individual zinc finger domains which bind sequences of the form 5'-GXX-3', where X is any base. See also U.S. Patent No. 6,140,081. The latter method has the severe limitation that it does not provide instructions permitting the specific targeting of triplets
25 containing nucleotides other than G in the 5' position of each triplet, which greatly restricts the potential target sequences of such generated zinc finger peptides.

International patent application WO98/53057 addresses the above problems by recognizing that zinc fingers can specify overlapping 4 bp subsites, and therefore synergy
30 between adjacent zinc finger domains is an important consideration in selecting zinc finger nucleic acid-binding domains to specifically target any sequence.

With the recent completion of the human genome project and the rapidly advancing fields of transgenic animals and plants, thousands of uncharacterised (and characterised) genes have (and will) become valid targets for functional genomics and other such projects.

Concomitantly, 'designer' zinc finger peptides are emerging as one of the most universal

5 and desirable ways of regulating the expression of specific genes within cells. See, for

example, Choo, Y., Sanchez-Garcia, I. & Klug, A. (1994) *Nature* 372: 642-645; Beerli,

R. R., Dreier, B. & Barbas, C. F. III (2000) *Proc. Natl. Acad. Sci. USA* 97: 1495-1500;

Kim, J-S. & Pabo, C. O. (1998) *Proc. Natl. Acad. Sci. USA* 95: 2812-2817; Kang, J. S. &

Kim, J-S. (2000) *J. Biol. Chem.* 275: 8742-8748); Zhang *et al.* (2000) *J. Biol. Chem.*

10 275:33,850-33,860; Liu *et al.* (2001) *J. Biol. Chem.* 276:11,323-11,334; and Ren *et al.*

(2002) *Genes. Devel.* 16:27-32. See also WO 00/41566 and WO 01/19981. Hence, a

rapid method of creating multi-zinc finger peptides for the up- or down-regulation of any specific gene is highly desirable.

As stated above, synergy between adjacent zinc finger peptides is an important factor in

15 specific DNA recognition. Moreover, the findings reported in co-owned WO 01/53480,

which is hereby incorporated by reference, demonstrate that poly-zinc finger peptides

constructed from strings of 2-finger domains can provide greater DNA binding specificity.

20 Traditional strategies of zinc finger mutagenesis and selection, such as phage display,

particularly if employed for the selection of 2-zinc finger units to target any desired

binding site are limited by the size of the library that can be cloned into host/vector

systems, such as phage. Due to limitations in library size imposed by such constraints, it

is impossible to include an exhaustive combination of randomisations to cover all

25 potentially important sequence-space. Furthermore, for important applications of

engineered zinc finger peptides, such as for gene therapy or transgenic animal systems,

engineered zinc finger peptides run the significant risk of eliciting a harmful

immunological reaction in the host animal.

30 The human genome sequencing project has also revealed the presence of almost 700

endogenous zinc finger-containing proteins. Assuming that each of these proteins

contains at least 2 finger modules, there are probably at least 2,000 natural zinc finger modules in the human genome alone. Similar numbers are expected in other animal and plant genomes.

5 SUMMARY

The present invention recognises the potential importance of designer zinc finger peptides in therapeutic and transgenic applications in animals and plants. Furthermore the present invention acknowledges that the safety of such applications is of primary importance.

10

The present invention provides the isolation of natural zinc finger modules, from genomes such as human, mouse, chicken, arabidopsis and other species, and the construction of non-natural combinations of such zinc finger modules, to create multi-finger domains, and to provide and determine novel nucleic acid binding specificities.

15

Such a procedure will allow the identification of the novel zinc finger domains that bind any desired nucleic acid sequence, particularly sequences of between 6 and 10

nucleotides long. The first advantage of such technology is that millions of years of natural evolution, to create specific nucleotide-binding zinc finger modules, are captured to create novel nucleic acid-binding domains. Also, use of poly-zinc finger peptides

20

constructed from such units for targeted gene regulation avoids the potentially harmful effects of host immune responses. The present invention thus greatly enhances the possibilities for the use of zinc finger transcription factors for *in vivo* applications, such as gene therapy and transgenic animals.

25

In a first aspect, therefore, there is provided a composite binding polypeptide comprising a first natural binding domain derived from first natural binding polypeptide, and a second natural binding domain derived from a second natural binding polypeptide, wherein said first and second natural binding polypeptides may be the same or different; which polypeptide binds to a target, said target differing from the natural target of the

30

both the first and the second binding polypeptides.

Preferably, said first and second natural binding polypeptides are different polypeptides.

Binding polypeptides according to the invention comprise two or more natural binding domains, advantageously three or more natural binding domains; advantageously, six or more domains are included. These are preferably arranged in a 3x2 conformation,
5 separated by linker sequences.

The binding domains are preferably nucleic acid binding domains, and the composite polypeptide is preferably a nucleic acid binding polypeptide. Most preferably, the composite polypeptide is a zinc finger polypeptide, and the natural binding domains are
10 zinc finger domains.

Zinc finger binding domains can comprise any type of zinc finger or zinc-coordinated structure including, but not limited to, Cys2-His2 (SEQ ID NO:1) zinc finger binding domain or Cys3-His (SEQ ID NO:2) zinc finger binding domains.
15

In a further aspect, there is provided a library of natural binding domains. The natural binding domains are the domains that may be assembled into polypeptides according to the previous aspect of the invention. Preferably, the library is of natural zinc finger nucleic acid binding domains.
20

Said zinc finger domains may comprise a linker attached thereto. Any linker amino acid sequence known in the art can be used. Advantageously, the linker comprises the amino acid sequence TGEKP (SEQ ID NO:3).

25 In a further aspect, the invention provides a method for selecting a binding polypeptide capable of binding to a target site, comprising:

- (a) providing a library of natural binding domains;
- (b) assembling two or more of said domains to form a composite polypeptide;
- (c) screening said composite polypeptide against the target site in order to
30 determine its ability to bind the target site.

Preferably, the natural binding domains are zinc finger binding domains.

Furthermore, the invention provides methods for designing a composite binding polypeptide, comprising:

- (a) providing information defining a target site;
- 5 (b) selecting, from a database of natural binding domains, a sequence of binding domains, separated by linker sequences, which is predicted to bind to the target site;
- (c) displaying the sequence of binding domains and linkers and optionally assembling the binding polypeptide from a library of said domains.

10 In certain embodiments, the binding domains are zinc finger domains. In certain embodiments, a binding domain sequence that will bind a particular target site is predicted by the application of one or more rules that define target binding interactions for the binding domains. In additional embodiments, a nucleotide sequence encoding the binding domains is assembled and introduced into a cell such that the composite binding
15 polypeptide is expressed.

In one embodiment, zinc fingers can be considered to bind to a nucleic acid triplet, in which case domains can be selected according to one or more of the following rules:

- (a) if the 5' base in the triplet is G, then position +6 in the α -helix is Arg; or
20 position +6 is Ser or Thr and position ++2 is Asp;
- (b) if the 5' base in the triplet is A, then position +6 in the α -helix is Gln and ++2 is not Asp;
- (c) if the 5' base in the triplet is T, then position +6 in the α -helix is Ser or Thr and position ++2 is Asp;
- 25 (d) if the 5' base in the triplet is C, then position +6 in the α -helix may be any amino acid, provided that position ++2 in the α -helix is not Asp;
- (e) if the central base in the triplet is G, then position +3 in the α -helix is His;
- (f) if the central base in the triplet is A, then position +3 in the α -helix is Asn;
- (g) if the central base in the triplet is T, then position +3 in the α -helix is Ala, Ser
30 or Val; provided that if it is Ala, then one of the residues at -1 or +6 is a small residue;

(h) if the central base in the triplet is C, then position +3 in the α -helix is Ser, Asp, Glu, Leu, Thr or Val;

(i) if the 3' base in the triplet is G, then position -1 in the α -helix is Arg;

(j) if the 3' base in the triplet is A, then position -1 in the α -helix is Gln;

5 (k) if the 3' base in the triplet is T, then position -1 in the α -helix is Asn or Gln;

(l) if the 3' base in the triplet is C, then position -1 in the α -helix is Asp.

In a further embodiment, the zinc fingers can be considered to bind to a nucleic acid quadruplet and domains can be selected according to one or more of the following rules:

10 (a) if base 4 in the quadruplet is G, then position +6 in the α -helix is Arg or Lys;

(b) if base 4 in the quadruplet is A, then position +6 in the α -helix is Glu, Asn or Val;

(c) if base 4 in the quadruplet is T, then position +6 in the α -helix is Ser, Thr, Val or Lys;

15 (d) if base 4 in the quadruplet is C, then position +6 in the α -helix is Ser, Thr, Val, Ala, Glu or Asn;

(e) if base 3 in the quadruplet is G, then position +3 in the α -helix is His;

(f) if base 3 in the quadruplet is A, then position +3 in the α -helix is Asn;

(g) if base 3 in the quadruplet is T, then position +3 in the α -helix is Ala, Ser or Val; provided that if it is Ala, then one of the residues at -1 or +6 is a small residue;

(h) if base 3 in the quadruplet is C, then position +3 in the α -helix is Ser, Asp, Glu, Leu, Thr or Val;

(i) if base 2 in the quadruplet is G, then position -1 in the α -helix is Arg;

(j) if base 2 in the quadruplet is A, then position -1 in the α -helix is Gln;

25 (k) if base 2 in the quadruplet is T, then position -1 in the α -helix is His or Thr;

(l) if base 2 in the quadruplet is C, then position -1 in the α -helix is Asp or His;

(m) if base 1 in the quadruplet is G, then position +2 is Glu;

(n) if base 1 in the quadruplet is A, then position +2 Arg or Gln;

(o) if base 1 in the quadruplet is C, then position +2 is Asn, Gln, Arg, His or Lys;

30 (p) if base 1 in the quadruplet is T, then position +2 is Ser or Thr.

In a preferred embodiment, zinc fingers are considered to bind to a nucleic acid quadruplet and domains are selected according to one or more of the following rules:

(a) if base 4 in the quadruplet is G, then position +6 in the α -helix is Arg; or position +6 is Ser or Thr and position ++2 is Asp;

5 (b) if base 4 in the quadruplet is A, then position +6 in the α -helix is Gln and ++2 is not Asp;

(c) if base 4 in the quadruplet is T, then position +6 in the α -helix is Ser or Thr and position ++2 is Asp;

10 (d) if base 4 in the quadruplet is C, then position +6 in the α -helix may be any amino acid, provided that position ++2 in the α -helix is not Asp;

(e) if base 3 in the quadruplet is G, then position +3 in the α -helix is His;

(f) if base 3 in the quadruplet is A, then position +3 in the α -helix is Asn;

(g) if base 3 in the quadruplet is T, then position +3 in the α -helix is Ala, Ser or Val; provided that if it is Ala, then one of the residues at -1 or +6 is a small residue;

15 (h) if base 3 in the quadruplet is C, then position +3 in the α -helix is Ser, Asp, Glu, Leu, Thr or Val;

(i) if base 2 in the quadruplet is G, then position -1 in the α -helix is Arg;

(j) if base 2 in the quadruplet is A, then position -1 in the α -helix is Gln;

(k) if base 2 in the quadruplet is T, then position -1 in the α -helix is Asn or Gln;

20 (l) if base 2 in the quadruplet is C, then position -1 in the α -helix is Asp;

(m) if base 1 in the quadruplet is G, then position +2 is Asp;

(n) if base 1 in the quadruplet is A, then position +2 is not Asp;

(o) if base 1 in the quadruplet is C, then position +2 is not Asp;

(p) if base 1 in the quadruplet is T, then position +2 is Ser or Thr.

25

Two or more composite polypeptides comprising two or more domains which are selected for binding to two or more target sites can be combined to provide a composite polypeptide which binds to an aggregate binding site comprising the two or more target binding sites.

30

In a still further aspect, the invention provides a computer-implemented method for designing a zinc finger polypeptide that binds to a target nucleic acid sequence, comprising the steps of:

- 5 (a) providing a system comprising at least storage means for storing data relating to a library of zinc fingers; storage means for storing a rule table; means for inputting target nucleic acid sequence data; processing means for generating a result; and means for outputting the result;
- (b) inputting sequence data for a target nucleic acid molecule;
- (c) defining a first target zinc finger binding site in said nucleic acid molecule;
- 10 (d) interrogating the zinc finger library and rule table storage means, comparing zinc fingers to the target zinc finger binding site according to the rule table and selecting zinc finger data identifying a zinc finger capable of binding to said target site;
- (e) defining at least one further target zinc finger binding site and repeating step (d); and
- 15 (f) outputting the selected zinc finger data.

Such a method may further comprise sending instructions to an automated chemical synthesis system to assemble a zinc finger polypeptide as defined by the zinc finger data obtained in (f).

20

In additional embodiments, the sequence of one or more oligonucleotides encoding a composite binding polypeptide can be determined from the sequence of a composite binding polypeptide, and the one or more oligonucleotides can be synthesized by any number of well-known methods.

25

Preferably, a composite binding polypeptide is tested for binding to a target sequence, and data from said testing is used to select, from a plurality of possibilities, a composite binding polypeptide that binds with optimal affinity and specificity to the target site.

30

Advantageously, two or more zinc finger polypeptides are combined to form a zinc finger polypeptide capable of binding to an aggregate binding site comprising two or more target sites.

The rule table preferably comprises rules as set forth above.

BRIEF DESCRIPTION OF THE FIGURES

5 **Figure 1** shows a flowchart depicting part of the logic used in the selection of zinc fingers from a natural library in accordance with the invention. The logic set forth in Figure 1 may be supplemented, for example using Rules relating to zinc finger overlap. Functional testing of zinc fingers for binding to the desired binding site may be implemented in an automated fashion and integrated with the zinc finger design system.

10

Figure 2 is a schematic representation of the human zinc finger mini-library construction procedure. Synthetic zinc finger coding oligonucleotides are assembled into full-length ds expression constructs by overlap PCR.

15 **Figure 3** is a schematic representation of the fluorescent ELISA assay used to detect zinc finger peptides bound to double stranded DNA target sites. Streptavidin (7), biotinylated DNA target (5) linked to biotin (6), 3-finger peptide (4) fused to HA-tag (3), anti-HA antibody (2) fused to horseradish peroxidase (HRP, 1).

20 **Figure 4** depicts ELISA scores of 384 library 2 constructs screened against the 5'-GCG-TGG-GCG-3' (SEQ ID NO:4) target site. Six constructs showed significant binding, and are termed C8, G16, I19, I23, J19 and K19, according to their coordinates on the 384-well plate.

25 **Figure 5** depicts ELISA scores of selected library 2 members; B10, C8, G16, I23, J19, and K19, against different DNA target sites. The sequences of the target sites are (from back of graph to front): 5'-GCG-TGG-GCG-3' (SEQ ID NO:5) ; 5'-CCA-CTC-GGC-3' (SEQ ID NO:6); 5'-CCT-AGG-GGG-3'(SEQ ID NO:7); 5'-GGA-TAA-GCG-3' (SEQ ID NO:8); 5'-GGG-AGG-CCT-3' (SEQ ID NO:9); 5'-GCG-TAA-GGA-3' (SEQ ID NO:10); 5'-GCG-GGG-GGA-3' (SEQ ID NO:11); and no DNA control (front row).

30

Figure 6 depicts a schematic representation of the 3-zinc finger library constructed according to the procedure described in Example 2.

DETAILED DESCRIPTION

5

Definitions

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art (e.g., in cell culture, molecular genetics, nucleic acid chemistry, hybridisation techniques and biochemistry).
The practice of the present invention will employ, unless otherwise indicated, conventional techniques of chemistry, molecular biology, microbiology, recombinant DNA, immunology, chemical methods, pharmaceutical formulations and delivery and treatment of patients, which are within the capabilities of a person of ordinary skill in the art. Such techniques are explained in the literature. See, for example, J. Sambrook, E. F. Fritsch, and T. Maniatis, 1989, *Molecular Cloning: A Laboratory Manual*, Second Edition, Books 1-3, Cold Spring Harbor Laboratory Press; Ausubel, F. M. et al. (1995 and periodic supplements; *Current Protocols in Molecular Biology*, ch. 9, 13, and 16, John Wiley & Sons, New York, N.Y.); B. Roe, J. Crabtree, and A. Kahn, 1996, *DNA Isolation and Sequencing: Essential Techniques*, John Wiley & Sons; J. M. Polak and James O'D. McGee, 1990, *In Situ Hybridisation: Principles and Practice*; Oxford University Press; M. J. Gait (Editor), 1984, *Oligonucleotide Synthesis: A Practical Approach*, IRL Press; and, D. M. J. Lilley and J. E. Dahlberg, 1992, *Methods of Enzymology: DNA Structure Part A: Synthesis and Physical Analysis of DNA* Methods in Enzymology, Academic Press. Each of these general texts is herein incorporated by reference.

The term "library" is used according to its common usage in the art, to denote a collection of different polypeptides or, preferably, a collection of nucleic acids encoding different polypeptides. The libraries of natural zinc finger peptides referred to herein comprise or encode a repertoire of polypeptides of different sequences, each of which has a preferred binding sequence.

The terms “polypeptide”, “peptide” and “protein” are used interchangeably to refer to a polymer of amino acid residues, preferably including naturally occurring amino acid residues. Artificial amino acid residues are also within the scope of the invention, but the
5 exclusive use of naturally-occurring amino acids is preferred in order to maintain the natural nature of the binding domains. There are 20 common amino acids, each specified by a different arrangement of three adjacent DNA nucleotides by the genetic code. These are the building blocks of proteins. Joined together in a strictly ordered chain by peptide bonds, the sequence of amino acids determines each polypeptide molecule. The 20
10 common amino acids are: alanine, arginine, aspartic acid, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, valine, cysteine, methionine, lysine, and asparagine. Virtually all of these amino acids (except glycine) possess an asymmetric carbon atom, and thus are potentially chiral in nature.

15 As used herein, “nucleic acid” includes both RNA and DNA, and nucleic acids constructed from natural nucleic acid bases or synthetic bases, or mixtures thereof. Modified nucleic acids such as, for example, PNAs and morpholino nucleic acids, are also included in this definition.

20 A “gene”, as used herein, is the segment of nucleic acid (typically DNA) that is involved in producing a polypeptide chain or ribonucleic acid gene product. It includes regions preceding and following the coding region (leader and trailer) as well as intervening sequences (introns) between individual coding segments (exons). Preferably, “gene”
25 includes the necessary control sequences for gene expression, as well as the coding region encoding the gene product.

A “binding polypeptide” is a polypeptide capable of binding to a specific target. Although, as is well known, polypeptides are capable of non-specific binding to a wide
30 range of substrates, it is also known that certain polypeptides, such as antibodies and other members of the immunoglobulin superfamily, zinc fingers, leucine zipper polypeptides, peptide aptamers and the like can bind specifically to target sites or

molecules. Generally, specific binding is preferably achieved with a dissociation constant (K_d) of 100 μ M or lower; preferably 10 μ M or better; preferably 1 μ M or better; and ideally 0.5 μ M or better. Binding polypeptides can be nucleic acid binding polypeptides which bind to nucleic acid in a target sequence-specific manner, such as zinc finger polypeptides. Unless specifically noted, no difference is intended herein between terms such as "peptide", "polypeptide" and "protein".

A "natural binding polypeptide" is a binding polypeptide encoded by the genome of a living organism such as, for example, a plant or animal.

A "composite" polypeptide is a polypeptide that is assembled from a plurality of components. In a preferred embodiment, the invention provides composite binding polypeptides that are assembled from a plurality of individual natural binding domains as set forth in detail herein. Typically, such domains are zinc finger nucleic acid binding domains.

A "natural binding domain" (or module) is a domain of a naturally occurring polypeptide that is capable of specific binding to a target as defined above. The terms "domain" and "module", according to their ordinary signification in the art, refer to a discrete continuous part of the amino acid sequence of a polypeptide that can be equated with a particular function. Protein domains or modules are largely structurally independent and can retain their structure and function in different environments. In certain embodiments, a natural binding domain or module is a zinc finger that binds a triplet or quadruplet nucleotide sequence.

Preferably, each of the individual natural binding domains that make up a composite binding polypeptide contain no changes in sequence, as compared to the natural sequence. However, those skilled in the art will understand that certain changes including conservative amino acid substitutions, as well as additions or deletions, may be made without altering the function of a domain. Moreover, where the changes are consistent with sequences common to the species from which the domain is derived, such as for

example being present in consensus sequences, they are unlikely to give rise to immunological problems.

Conservative amino acid substitutions may be made, for example according to Table 1.

- 5 Amino acids in the same block in the second column and preferably in the same line in the third column may be substituted for one another:

Table 1

ALIPHATIC	Non-polar	G A P
		I L V
	Polar - uncharged	C S T M
		N Q
	Polar - charged	D E
		K R
AROMATIC		H F W Y

A domain is “derived” from a protein if it is effectively removed from a naturally-occurring protein for use in a composite binding polypeptide. Removal may be physical removal, by cleavage of the protein; more commonly, however, the sequence of the domain is determined and the domain is synthesised by protein synthesis techniques to be a copy of the naturally-occurring domain. Alternatively, a nucleic acid encoding the domain is synthesized and expressed in a cell. *In vitro* synthesised domains, or *in vitro* synthesized polynucleotides encoding naturally-occurring domains, are considered to be “derived” from the natural protein if they recapitulate the sequence of the naturally-occurring domain.

A “target” is a molecule or part thereof to which a binding polypeptide or a binding domain is capable of specific binding. The “natural target” of a binding polypeptide is the target to which that polypeptide binds in nature; *e.g.*, in a living cell. In the case of zinc finger polypeptides, for instance, the natural target is the nucleotide sequence to which the polypeptide binds in a living cell. Sequences other than the natural target, as defined herein, to which a zinc finger polypeptide may bind *in vitro* are not natural targets.

20

In the case of nucleic acid binding polypeptides, therefore, the term “target” may be substituted or supplemented with “binding site” or “binding sequence.” Where binding sites are assembled to form larger binding sites, which are bound by multi-domain

binding polypeptides, such binding sites are referred to as “aggregate binding sites”, indicating that they are formed by the juxtaposition of two or more individual binding sites. The aggregate binding sites can comprise contiguous individual binding sites, or individual binding sites interspersed by one or more intervening nucleotides or sequence
5 of nucleotides.

The present invention relates to naturally-occurring zinc fingers and their use as specific nucleic acid binding modules in combinations not present in nature. This invention provides methods of determining and/or predicting the nucleotide binding specificities of
10 natural zinc finger modules. Also provided are methods of constructing poly-zinc finger peptides containing at least one natural zinc finger module, from libraries of natural zinc finger peptides, and methods of screening such peptides to determine their preferred nucleotide binding specificity. Moreover, the invention provides for the use of combinations of such natural zinc finger modules in poly-zinc finger peptides not present
15 in nature, to bind any desired nucleotide sequence.

Poly-zinc finger peptides of this invention may contain 2, 3, 4, 5, 6 or more zinc finger modules. Natural zinc finger modules of this invention may preferably be linked by canonical, flexible or structured linkers, as set out below and in WO 01/53480, the disclosure of which is hereby incorporated by reference. More preferably, the linkers are
20 canonical linkers such as -TGEKP- (SEQ ID NO:3).

The poly-zinc finger peptides of this invention can be given useful biological functions by the addition of effector domains, creating chimeric zinc finger peptides. Preferably, such chimeric zinc finger peptides may be used to up- or down-regulate desired genes, *in vitro* or *in vivo*. Preferable effector domains include transcriptional repressor domains, transcriptional activator domains, transcriptional insulator domains, chromatin remodelling domains, enzymatic domains, and signalling / targeting sequences or domains. To cause a desired biological effect composite binding polypeptides can bind to one or more suitable nucleotide sequences *in vivo* or *in vitro*. Preferred DNA regions
25 from which to effect the up- or down-regulation of specific genes include promoters, enhancers or locus control regions (LCRs). Other suitable regions within genomes,
30

which may provide useful targets for composite binding polypeptides include telomeres and centromeres.

5 The expression of many genes is also achieved by controlling the fate of the associated RNA transcript. RNA molecules often contain sites for RNA-binding proteins, which determine RNA half-life. Hence, composite binding polypeptides can also control endogenous gene expression by specifically targeting RNA transcripts to either increase or decrease their half-life within a cell.

10 Composite binding polypeptides can also be fused to epitope tags, which can be detected by antibodies, and may therefore be used to signal the presence or location of a particular nucleotide sequence in a mixed pool of nucleic acids, or immobilised on the surface of a chip or other such surface.

15 Intracellular localization of composite binding polypeptides can be regulated, for example, by fusion to a localization domain, for example, a nuclear localization sequence or a localization domain as disclosed, for example, in PCT/US01/42377.

a. Nucleic Acid Binding Polypeptides

20

This invention preferably relates to nucleic acid binding polypeptides. Preferably, the binding polypeptides of the invention are DNA binding polypeptides. Particularly preferred examples of nucleic acid binding polypeptides are zinc finger peptides.

25 Zinc finger peptides typically contain strings of small nucleic acid binding domains, each stabilised by the co-ordination of zinc. These individual domains are also referred to as “fingers” and “modules”. A zinc finger recognises and binds to a nucleic acid triplet, or an overlapping quadruplet, in a DNA target sequence. However, zinc fingers are also known to bind RNA and proteins. Clemens, K. R. *et al.*, (1993) *Science* 260: 530-533;
30 Bogenhagen, D.F. (1993) *Mol. Cell. Biol.* 13: 5149-5158; Searles, M. A. *et al.*, *J. Mol. Biol.* 301: 47-60 (2000); Mackay, J. P. & Crossley, M. (1998) *Trends Biochem. Sci.* 23: 1-4.

Preferably, there are 2 or more zinc fingers, for example 2, 3, 4, 5, 6, or 7 zinc fingers, in each zinc finger polypeptide. Advantageously, there are 3 or more zinc fingers in each zinc finger polypeptide.

5

All of the DNA binding residue positions of zinc finger peptides, as referred to herein, are numbered from the first residue in the α -helix of the finger, ranging from +1 to +9. "-1" refers to the residue in the framework structure immediately preceding the α -helix in a zinc finger peptide. Residues referred to as "++" are residues present in an adjacent
10 (C-terminal) peptide. Where there is no C-terminal adjacent peptide, "++" interactions do not operate.

The α -helix of a zinc finger peptide aligns antiparallel to the target nucleic acid strand, such that the primary nucleic acid sequence is arranged 3' to 5' in order to correspond
15 with the N- terminal to C-terminal sequence of the zinc finger peptide. Since nucleic acid sequences are conventionally written 5' to 3', and amino acid sequences N-terminus to C-terminus, the result is that when a target nucleic acid sequence and a zinc finger peptide are aligned according to convention, the primary interaction of the zinc finger peptide is with the "minus" strand of the nucleic acid sequence, since it is this strand
20 which is aligned 3' to 5'. These conventions are followed in the nomenclature used herein. It should be noted, however, that in nature certain zinc finger modules, such as zinc finger 4 of the protein GLI, bind to the "plus" strand of the nucleic acid sequence. See Suzuki *et al.* (1994) *Nucl. Acids Rev.* 22: 3397-3405; and Pavletich & Pabo, (1993) *Science* 261: 1701-1707. The present invention encompasses incorporation of such zinc
25 finger peptides into DNA binding molecules.

Natural Zinc Finger Peptides.

In certain embodiments, this invention relates to natural zinc finger modules. As used
30 herein, the term 'natural' with reference to a zinc finger, means that the DNA sequence which encodes a particular zinc finger, whether normally expressed *in vivo* or not, is found in nature, *i.e.* is part of the genome of a cell. A natural human zinc finger is one

which is endogenous to the human genome, a natural mouse zinc finger is found in the mouse genome, and a natural viral zinc finger is found in a viral genome, *etc.* Natural zinc finger genes which have become integrated into the genome of a heterologous species by natural means, *e.g.*, integration of a viral genome into a host genome, are
 5 considered to be endogenous to the host species within the context of this disclosure. A zinc finger module constructed or produced *in vitro* or extracted from an *in vivo* source is considered to be natural if its amino acid sequence matches that of the amino acid sequence encoded by its natural gene. The DNA sequence of the natural gene is not the defining aspect. Thus, polynucleotides encoding natural zinc finger modules may have a
 10 different sequence from that of the naturally-occurring sequence encoding the module, *e.g.*, to adjust codon usage to optimise expression of the module in a particular expression system.

Preferably, sequences of zinc fingers used in the present invention are not mutated from
 15 their natural form. Advantageously, the natural zinc finger polypeptides are expressed in nature.

A natural zinc finger binding motif is a structure well known to those in the art and defined in, for example, Miller *et al.*, (1985) *EMBO J.* 4: 1609-1614; Berg (1988) *Proc.*
 20 *Natl. Acad. Sci. USA* 85: 99-102; Lee *et al.*, (1989) *Science* 245: 635-637; see also International patent applications WO 96/06166 and WO 96/32475, incorporated herein by reference.

In general, a natural zinc finger framework has the structure:

25 SEQ ID NO:12 X_{0-2} C X_{1-5} C X_{9-14} H X_{3-6} ^H/C

where X is any amino acid, and the numbers in subscript indicate the possible numbers of residues represented by X (Formula A).

In a preferred aspect of the present invention, natural zinc finger nucleic acid binding
 30 motifs may be represented as motifs having the following primary structure:

X_{0-2} C X_{1-5} C X_{2-7} X X X X X X X H X_{3-6} ^H/C (SEQ ID NO:14)
 (SEQ ID NO:13)

-1 1 2 3 4 5 6 7

where X is any amino acid, and the numbers in subscript indicate the possible numbers of residues represented by X (Formula A'). The numbers -1 through 7 refer to amino acid position with respect to the beginning of the alpha-helical region of the zinc finger.

- 5 The Cys and His residues, which together co-ordinate the zinc metal atom, are marked in bold text and are usually invariant. However, all naturally-occurring zinc finger modules, even if they diverge from the above formula, are encompassed within the scope of this invention.
- 10 Zinc finger modules of formula A' are often arranged in tandem within a natural zinc finger polypeptide, such that a zinc finger containing protein may have 2, 3, 4, 5, 6, 7, 8, 9 or more individual zinc finger motifs. In such a protein, individual zinc fingers are joined to each other by a polypeptide sequence known as a linker. Generally, such a natural linker lacks secondary structure, although the amino acids within the linker may
- 15 form local interactions when the protein is bound to its target site. By 'linker sequence' is meant an amino acid sequence that links together adjacent zinc finger modules. For example, in a natural zinc finger protein, the linker sequence is the amino acid sequence which lies between the last residue of the α -helix in a zinc finger and the first residue of the β - sheet in the next zinc finger. The linker sequence therefore joins together two zinc
- 20 fingers. For the purposes of the present invention, the last amino acid of the α -helix in a zinc finger is considered to be the final zinc coordinating histidine (or cysteine) residue, while the first amino acid of the following finger is generally a tyrosine / phenylalanine or another hydrophobic residue. Since some natural zinc fingers do not start with a hydrophobic residue (see Appendices), the start of a finger is sometimes harder to define
- 25 from amino acid sequence (or indeed zinc finger structure), and so some flexibility must be allowed in this definition. Accordingly, in a natural zinc finger protein, threonine is often considered to be the first residue in the linker, and proline is the last residue of the linker. Thus, for example, in the natural Zif268 peptide the linker sequence is -
- 30 TG(E/Q)(K/R)P- (SEQ ID NO:15). Although natural linkers can vary greatly in terms of amino acid sequence and length, on the basis of sequence homology, the canonical

natural linker sequence is considered to be -TGEKP- (SEQ ID NO:3). Hence, the preferred linker sequence to join zinc finger modules of the present invention is -TGEKP-.

5 Additionally, a 'leader' peptide may be added to the N-terminal zinc finger of a poly-zinc finger peptide to aid its expression, without changing the sequence of the natural zinc finger module. Preferably, the leader peptide is MAEERP (SEQ ID NO:16) or MAERP (SEQ ID NO:17).

10 In general, naturally occurring zinc finger modules may be selected from those proteins for which the DNA binding specificity is already known. For example, these may be the proteins for which a crystal structure has been resolved: namely Zif268 (Elrod-Erickson *et al.* (1996) *Structure* 4: 1171-1180), GLI (Pavletich & Pabo (1993) *Science* 261: 1701-1707), Tramtrack (Fairall *et al.* (1993) *Nature* 366: 483-487) and YY1 (Houbaviy *et al.* (1996) *Proc. Natl. Acad. Sci. USA* 93: 13577-13582). Furthermore, the sequence
15 specificity of many naturally-occurring zinc fingers and zinc finger proteins are known. In addition, this invention further provides for the determination of the binding specificity of natural zinc finger modules for use in the present invention. See "Prediction of Binding Specificity," *infra*.

20

Poly-Zinc Finger Peptides.

It is desirable that a 'designer' transcription factor for uses such as gene therapy and in transgenic organisms should have the ability to target virtually unique sites within any genome. For complex genomes such as in humans, an address of at least 16 bps is
25 required to specify a potentially unique DNA sequence. Shorter DNA sequences have a significant probability of appearing several times in a genome, raising the possibility of obtaining undesirable non-specific gene targeting with a designed transcription factor targeted to such a shorter sequence. As individual zinc fingers only bind 3 to 4 nucleotides, it is therefore necessary to construct multi-finger polypeptides to target these
30 longer sequences. A six-zinc finger peptide (with an 18 bp recognition sequence) could, in theory, be used for the specific recognition of a single target site and hence, the

specific regulation of a single gene within any genome. In addition, a significant increase in binding affinity might also be expected, compared to a protein with fewer fingers. In simple terms, if a three-finger peptide (with a 9 bp recognition sequence) binds DNA with nanomolar affinity, two tandemly linked three-finger peptides might be expected to bind an 18 bp sequence with an affinity of 10^{-15} - 10^{-18} M. However, most previous attempts at producing high-affinity 6-finger peptides (poly-zinc finger peptides) based on fusions of two 3-finger domains have been unsuccessful in generating much of an improvement in affinity over 3-finger peptides. Liu, Q., Segal, D. J., Ghiara, J. B. & Barbas, C. F. III (1997) *Proc. Natl. Acad. Sci. USA* 94: 5525-5530; Kim, J-S. & Pabo, C. O. (1998) *Proc. Natl. Acad. Sci. USA* 95: 2812-2817; Kamiuchi, T., Abe, E., Imanishi, M., Kaji, T., Nagaoka, M. & Sugiura, Y. (1998) *Biochemistry* 37: 13827-13834. To optimise both the affinity and specificity of 6-finger peptides, a fusion of three 2-finger domains has been shown to be advantageous. Moore, M., Klug, A. & Choo, Y. (2001) *Proc. Natl. Acad. Sci. USA* 98: 1437-1441; and WO 01/53480. Therefore, in one embodiment, 2-finger units are linked to make poly-zinc finger nucleotide-binding domains. A pool of 4096 such 2-finger units, that recognise all possible 6 bp sequences ($4^6=4096$), represents an archive sufficient to rapidly create universal nucleic acid recognition, by simple linkage, in an "off-the-shelf" manner. See Moore *et al.*, *supra* and WO 01/53480.

Poly-zinc finger peptides according to this invention may be constructed containing 2, 3, 4, 5, 6 or more zinc finger modules. Such poly-zinc finger peptides may contain inter-finger linkers other than the canonical (TGEKP) linker sequence, as described, for example, in WO 01/53479; Moore, M., Choo, Y. & Klug, A. (2001) *Proc. Natl. Acad. Sci. USA* 98: 1432-1436; and Moore, M., Klug, A. & Choo, Y. (2001) *Proc. Natl. Acad. Sci. USA* 98: 1437-1441. Briefly, linker sequences may be flexible or structured but, in general, will not form base-specific interactions with the target nucleotide sequence. A 'flexible' linker is defined as one which does not form a specific secondary structure in solution, whereas a 'structured' linker is defined as one that adopts a particular secondary structure in solution. Preferably, flexible linkers include the sequences GGERP (SEQ ID NO:18), GSERP (SEQ ID NO:19), GGGGSERP (SEQ ID NO:20), GGGGSGGSERP (SEQ ID NO:21), GGGGSGGSGGSERP (SEQ ID NO:22),

GGGGSGGSGGSGGSGGSERP (SEQ ID NO:23). Preferably, the structured linker comprises an amino acid sequence that is not capable of specifically binding nucleic acid. More preferably, the structured linker comprises the amino acid sequence of TFIIIA finger IV. Alternatively, or in addition, the structured linker is derived from a zinc finger
5 by mutation of one or more of its base contacting residues to reduce or abolish nucleic acid binding activity of the zinc finger. The zinc finger may be finger 2 of wild type Zif268 mutated at positions -1, 2, 3 and/or 6.

In one embodiment, this invention provides for the construction and screening of poly-
10 zinc finger peptides containing at least one natural zinc finger module.

In another embodiment, this invention provides for the construction and screening of poly-zinc finger peptides containing at least one natural zinc finger module, linked with the canonical linker sequence -TGEKP- (SEQ ID NO:3).

15

In one embodiment, methods for the construction and use of poly-zinc finger peptide comprising natural zinc finger modules are provided.

In another embodiment, methods for the construction and use of poly-zinc finger peptide
20 comprising natural zinc finger modules, linked with the canonical linker sequence -TGEKP- (SEQ ID NO:3), are provided.

In a further embodiment, methods for the construction and use of poly-zinc finger peptides comprising at least one natural zinc finger module, containing either flexible or
25 structured linkers (as described above and in WO 01/53480), are provided.

b. Advantages of Natural Zinc Finger Modules

Zinc finger modules are compact and stable structures of approximately 30 amino acids,
30 which contain the full information required to bind a nucleic acid triplet or overlapping quadruplet. As such, they have proven to be extremely versatile scaffolds for engineering novel DNA-binding domains. *See*, for example, Rebar, E. J. & Pabo, C. O. (1994)

- Science* 263, 671-673; Jamieson, A. C., Kim, S.-H. & Wells, J. A. (1994) *Biochemistry* 33, 5689-5695; Choo, Y. & Klug, A. (1994) *Proc. Natl. Acad. Sci. U.S.A.* 91, 11163-11167; Choo, Y., Sanchez-Garcia, I. & Klug, A. (1994) *Nature* 372, 642-645; Wu, H., Yang, W.-P. & Barbas III, C. F. (1995) *Proc. Natl. Acad. Sci. USA* 92, 344-348;
- 5 Greisman, H. A. & Pabo, C. O. (1997) *Science* 275, 657-661; Isalan, M., Klug, A. & Choo, Y. (1998) *Biochemistry* 37, 12026-12033; Choo, Y. (1998) *Nature Struct. Biol.* 5, 264-265; Segal, D. J., Dreier, B., Beerli, R. R. & Barbas, C. F. (1999) *Proc. Natl. Acad. Sci. USA* 96, 2758-2763; Isalan, M. & Choo, Y. (2000) *J Mol Biol* 295, 471-477; and Beerli, R. R., Dreier, B., Barbas, C.F. (2000) *Proc Natl Acad Sci U S A* 97, 1495-500.
- 10 The resulting engineered zinc finger domains have increased our knowledge of sequence-specific DNA recognition, as well as provided a wide range of potential tools for medicine and biotechnology.

As a result of these and other studies on zinc finger engineering, it has been recognised

15 that an individual zinc finger module does not necessarily recognise a simple nucleotide triplet, as was first thought; but instead, can bind to an overlapping quadruplet of double stranded DNA. *See*, for example, Isalan *et al.* (1997) *Proc Natl Acad Sci U S A* 94, 5617-5621; and WO98/53057). In this respect, zinc finger engineering strategies have been particularly important for deciphering the mechanism and specificity of these interactions.

20

With the recent completion of the human genome project and the rapidly advancing fields of transgenic animals and plants, thousands of uncharacterised (and characterised) genes have (and will) become valid targets for functional genomics and other such projects. Concomitantly, engineered zinc finger peptides (often as a component of “designer”

25 transcription factors) are emerging as one of the most universal and desirable ways of regulating the expression of specific genes within cells. *See*, for example, Choo, Y., Sanchez-Garcia, I. & Klug, A. (1994) *Nature* 372: 642-645; Beerli, R. R., Dreier, B. & Barbas, C. F. III (2000) *Proc. Natl. Acad. Sci. USA* 97: 1495-1500; Kim, J-S. & Pabo, C. O. (1998) *Proc. Natl. Acad. Sci. USA* 95: 2812-2817; Kang, J. S. & Kim, J-S. (2000) *J. Biol. Chem.* 275: 8742-8748; Zhang *et al.* (2000) *J. Biol. Chem.* 275:33,850-33,860; Liu *et al.* (2001) *J. Biol. Chem.* 276:11,323-11,334; Ren *et al.* (2002) *Genes. Devel.* 16:27-32; and WO 00/41566.

30

Notwithstanding the remarkable progress in zinc finger engineering, there remain several issues that limit the use of engineered zinc fingers for such applications. Points of particular concern include the potential immunogenicity of non-natural zinc fingers, and the 'fine-tuning' of particular aspects of the protein-DNA interactions to obtain optimal and specific zinc finger-nucleic acid contacts.

The present invention overcomes problems such as immunogenicity and optimal binding specificity, by exploiting the vast repertoire of naturally occurring zinc fingers to construct targeted zinc finger proteins having novel specificities.

Immunogenicity

The main function of the immune system is to detect, and render harmless, foreign particles which have invaded the body as a whole, or individual cells or organs. 'Foreign' in this context means non-host, i.e. a substance which has originated from a different species, or one which has originated as a result of a mutational event (such as might generate a malignant cell). On encountering such an antigenic particle, either in solution or on the surface of an infected cell, the body's defences rapidly destroy/remove it by complex pathways which involve the interaction of many members of the immune system. For a good overview of immunology see Roitt, *Essential Immunology*, Blackwell Science Ltd. and Roitt, I., Brostoff, J. & Male, D. *Immunology*, 4th Ed. Mosby. Hence, all biological therapeutic agents, such as peptides, nucleic acids, viruses, etc., risk eliciting an immune response in the recipient. Particularly for cases in which repeated doses of a therapeutic agent are required, this response can be strong and potentially dangerous to the host organism.

The immune system functions through either innate or adaptive responses. The innate response is usually the body's first internal line of defence. Phagocytic cells recognise and bind to foreign objects in extracellular environments. Once bound, the foreign object is internalised and destroyed. Foreign therapeutic agents such as peptides and nucleic acids, which are administered directly to the blood stream of the recipient, risk being

detected and possibly destroyed before they even reach their intended target. This response is one of primitive non-specific recognition of non-host agents, and does not adapt with time or exposure to the antigen.

5 Foreign therapeutic agents (or infectious agents such as bacteria and viruses), which evade the innate immune response and may have been successfully delivered to a particular cell have not necessarily avoided the host's immune system. Proteins that are expressed in cells are routinely degraded within lysosomes, and short peptide fragments, generally of between 6 and 9 amino acids, are transported to the cell surface and
10 presented to the host's immune system. This is the start of the host's second internal defence mechanism against invasion, the adaptive immune response. The proteins responsible for displaying such peptide fragments are known as major-histocompatibility complexes (MHC) proteins. Lymphocyte cells, known as T-lymphocytes, dock with the MHC proteins and scan the peptide fragments displayed. Contact of a T-lymphocyte with
15 a fragment specifically recognised as not belonging to the host organism initiates an immunological cascade which ultimately results in the host cell being destroyed or undergoing apoptosis. This mechanism is one of specific recognition, and once recognised as foreign, the antigen is 'remembered' so that any future invasions by the agent are dealt with more and more rapidly. B-cells are another type of lymphocyte that
20 recognise extracellular particles and then produce and release antibodies to help combat the agent.

To avoid potentially damaging the host organism and to ensure the successful delivery and action of a therapeutic peptide it is important to make it as much like a host protein as
25 is reasonably possible. In the case of synthesised therapeutic antibodies for human use, a great deal of work has gone in to the 'humanisation' of antibodies produced by other animal species (See EP 0239400). In this invention we present a solution for the equivalent problem associated with zinc finger therapeutic peptides.

30 To some extent, prior art zinc finger engineering strategies have attempted to minimise the risk of eliciting immune responses by using an engineering scaffold that is compatible with (*i.e.* that originates from) the recipient, and by limiting the sizes of the varied regions

within the final product. For example, typical engineered zinc fingers utilize a scaffold such as the three-finger DNA-binding domain of Zif268 (containing approximately 100 amino acid residues). Because the amino acid sequence of Zif268 is completely conserved in a variety of species, including mice and humans, the scaffold is not itself immunogenic in these species. However, in order to engineer new DNA-binding domains, stretches of approximately 7 amino acids must be varied within each zinc finger. These sequences of 7 amino acids represent modifications in positions -1, 1, 2, 3, 4, 5, and 6 of the α -helix of each finger. Although these engineered regions are considered to be relatively small, they are approximately the length of the peptide fragments displayed on the surface of cells by MHC molecules. Hence, they may provide antigenic peptide fragments in several registers of the amino acid sequence, which may result in dangerous and/or undesirable immune responses in the host.

Accordingly, it is not known whether this type of engineering strategy will be entirely sufficient to avoid all potential undesirable effects, or indeed whether it will create the most optimal framework for all zinc finger-nucleic acid interactions.

In addition to the zinc fingers themselves, it is also possible that inter-finger linker sequences could present potential immunological problems. Fortunately, natural zinc finger proteins display strong conservation and homology in their linker sequence. A very large number of natural fingers are joined by the canonical linker peptide -TGEKP- (SEQ ID NO:3), located between the final zinc chelating residue (usually histidine) of the first finger, and the first residue of the second finger (usually a large hydrophobic residue such as tyrosine or phenylalanine, which begins the β -sheet). Hence, the use of the canonical linker sequence -TGEKP- (SEQ ID NO:3), to join natural zinc finger modules in a non-natural order, will reduce the possibility of eliciting an undesirable immune reaction to a minimum. Furthermore, there are so many natural zinc fingers which are already joined by canonical linker sequences, that if deemed necessary, the database of natural zinc fingers used for the construction of poly-zinc finger peptides may be restricted to those already flanked by such linkers.

The periodicity of zinc fingers and their amenability to linkage using the TGEKP (SEQ ID NO:3) motif is illustrated in Table 2.

		α -HELIX		LINKER
5		-1123456		
	YA CPVESC	DRRFS	(SEQ ID NO:24)	RSDELTR H IR I H (SEQ ID NO:25) <u>TGEKP</u>
	FQ CRI	CMRNFS	(SEQ ID NO:26)	RSDHLST H IR T H (SEQ ID NO:27) <u>TGEKP</u>
	FA CDI	CGRKFA	(SEQ ID NO:28)	RSDEK R H T K I H (SEQ ID NO:29) <u>TGEKP</u>

Table 2. A functional three-finger DNA-binding domain based on the peptide sequence of Zif268. TGEKP linker motifs are underlined. The helical residues of each zinc finger are numbered relative to the first helical position, position +1. Conserved Cysteines and Histidines forming the classical Cys₂His₂ zinc finger core are shown in bold.

Fine-Tuning of Zinc Finger-Nucleic Acid Interactions.

It has previously been shown that zinc fingers cannot simply be regarded as independent nucleic acid-binding modules. Isalan, M., Klug, A. & Choo, Y. (1998) Biochemistry 37, 12026-12033; Isalan, M., Choo, Y. & Klug, A. (1997) Proc Natl Acad Sci 94, 5617-5621. The interactions between adjacent zinc fingers can be complex and involve overlap of binding sites, which means that optimal interfaces are not easily engineered through rational design. Combinatorial library selection systems, which if designed correctly necessarily result in interface compatibility, can help to engineer better optimisation of the zinc finger-nucleic acid interface. See, for example, WO98/53057. However, all library selection systems suffer from the problem of library size, whereby because of physical constraints, it is impossible to include an exhaustive combination of randomisations to cover all potentially important sequence-space. For example, to optimise the zinc finger-nucleic acid interface, subtle amino acid variations may be needed, even from positions outside the recognition α -helix. Furthermore, alternative approaches to zinc finger engineering, such as 'affinity maturation' through random mutation or gene shuffling, which may (to a limited extent) increase the coverage of sequence space, may also raise the probability of generating undesirable immunological problems. Hence, it is possible that the creation of truly optimal zinc finger domains for

recognition of specific nucleic acid sequences may be outside the scope of traditional engineering strategies.

In contrast, naturally occurring zinc finger modules have already been 'fine-tuned' by

5 thousands of years of natural selection and are, under normal circumstances, non-immunogenic in their host organism. The human genome project has revealed that zinc finger-containing proteins constitute the second most abundant family of proteins in humans, with well over 600 members. Since zinc finger proteins usually contain several individual zinc finger modules, the human genome provides a repertoire of thousands of
10 natural zinc finger modules for the creation of composite binding polypeptides.

Furthermore, because there are only 64 ($=4^3$) possible 3 bp sequences and 256 ($=4^4$) possible 4 bp sequences, it is likely that a natural zinc finger domain exists which is

capable of binding to every potential 3- or 4-nucleotide target sequence. Consequently, natural zinc fingers are a very useful resource for the production of composite binding

15 polypeptides comprising zinc fingers. At present, the natural binding site of many natural zinc finger modules is not known. Thus, to be useful for the construction of composite binding polypeptides, nucleotide sequence preferences for certain natural zinc fingers are determined according to rules tables disclosed in the following section ("Binding Specificity of Natural Zinc Finger Modules").

20

To create optimal poly-zinc finger peptides the potentially significant problem of

interface incompatibility must be addressed, since natural zinc finger modules will not necessarily be compatible with each other when juxtaposed. In this respect, a library construction and screening system is preferably employed which links natural zinc finger

25 modules in non-natural combinations, and screens them against possible target sequences

of greater than 3 or 4 bp in length (which represents the possible binding site of a single zinc finger module), to determine optimal 2- or 3-finger domains. In this way, the

cooperative nature of zinc finger binding is taken into account in the design and selection of composite binding polypeptides, and in the determination of the sequence specificity of

30 their binding. In one embodiment, a library of poly-zinc finger peptides containing at

least one natural zinc finger module is provided. Preferably, poly-zinc finger peptides of the library contain at least two natural zinc finger modules.

5 **c. Binding Specificity of Natural Zinc Finger Modules**

Disclosed herein are certain improvements to current limitations on the use of customised zinc finger nucleic acid binding domains, through the use of natural zinc finger modules. By using either natural 1-finger or 2-finger sub-domains, and/or novel combinatorially-
10 mixed, pre-selected 2-finger sub-domains, it is possible to construct poly-zinc finger peptides that bind any desired nucleotide target sequence, using non-natural combinations of natural zinc fingers.

This approach is particularly suited for human gene therapy applications, but the
15 invention is not just limited to zinc finger modules encoded by the human genome. For applications within transgenic animals such as mice, chicken, etc., the same system can be used, but incorporating natural zinc finger modules from those species instead (see Example 3). The genome of any organism (*e.g.*, animal, plant, bacterium, virus, *etc.*) can thus provide a genetic 'toolbox' of non-immunogenic, structurally optimised zinc fingers
20 for applications in that organism.

Before such zinc finger modules can be utilised, however, it is essential that their optimal binding site is determined, in isolation, or preferably as part of a 2- or 3-finger subdomain. Natural zinc finger modules are advantageously fused into subdomains
25 comprising two or three zinc finger modules in random arrangement, optionally comprising an anchor finger, then subjected to binding site analysis. An 'anchor' zinc finger is one for which the binding specificity is known, such as, for example, finger 1 or finger 3 of Zif268, each of which binds the sequence 5'-GCG-3'. An anchor finger is attached to the N- or C-terminus of the zinc finger module(s) or subdomain for which the
30 binding specificity is to be determined, and acts as an anchor to set the binding register for the binding site selection. For example, if the binding site preference of a pair of natural zinc fingers is to be determined, finger 1 of Zif268 may be fused to the N-

terminus of the pair of natural fingers, and a 5'-GCG-3' anchor sequence is placed at the 3' end of 6 or more randomised nucleotides. Selection of the optimal binding site may thus be conducted with an oligonucleotide containing the sequence 5'-XXX-XXX-GCG-3' (SEQ ID NO:30), where X is any specified nucleotide. The anchor sequence thereby
5 allows the binding site preference of the zinc finger libraries to be easily determined. Such procedures are described in the Examples.

Screening for Zinc Finger Binding Specificity

10 There are various approaches, known to those in the art, for screening nucleic acid binding peptides for their binding specificity. To determine the binding specificity of, for example, zinc finger peptides, procedures can be conducted using: (a) a library of zinc fingers and a specified target sequence – to select one or more zinc finger peptides with a particular binding preference; or (b) a single zinc finger peptide and a random population
15 of target sequences – to select one or more optimal binding sites for a particular peptide. For many applications, such as for the creation of transcription factors for regulating specific gene activity, it is often preferable to screen zinc finger libraries against specific target sequences. In this way, the search is geared towards a particular application. However, if the function or binding specificity of a natural protein is the object of the
20 investigation, a library of potential binding sites can be screened using a single peptide. Some such methods are outlined below.

A typical method for screening libraries of nucleic acid binding polypeptides against specific target sites is that of phage display. Phage display protocols generally involve
25 expressing the peptides under study as fusions with the gIII major coat protein of bacteriophage (J. McCafferty, R. H. Jackson, D. J. Chiswell, (1991) *Protein Engineering* 4, 955-961). Suitable protocols for the selection of zinc finger peptides have been described and are well known to those in the art. See, for example, Choo, Y. & Klug, A. (1994) *Proc. Natl. Acad. Sci. U.S.A.* 91, 11163-11167; Choo, Y., Sanchez-Garcia, I. &
30 Klug, A. (1994) *Nature* 372, 642-645; Choo, Y. (1998) *Nature Struct. Biol.* 5, 264-265; Choo, Y. & Klug, A. (1997) *Curr. Opin. Str. Biol.* 7, 117-125; 7 Isalan, M., Klug, A. & Choo, Y. (1998) *Biochemistry* 37, 12026-12033; Isalan, M. & Choo, Y. (2000) *J Mol*

Biol 295, 471-477; Isalan, M., Choo, Y. & Klug, A. (1997) Proc Natl Acad Sci 94, 5617-5621; WO 01/53480, WO 01/53479, WO 96/06166, WO 98/53057, WO 98/53058, WO 98/53059 and WO 98/53060 and references cited therein; see also Examples, *infra*. In general, sequences comprising target sites are bound, such as through biotin-streptavidin, to a solid support, such as a magnetic particle, or the surface of a tube or well. A solution of phage expressing members of a library of zinc finger peptides is then added to the immobilised target site. Non-bound phage are washed away and bound phage (containing the DNA encoding the bound zinc finger peptide), are collected. The collected phage sample is usually reused in further rounds of selection to enrich for the tightest binding zinc finger peptide.

Phage display protocols based on random mutagenesis of zinc finger modules are known to have a number of limitations. First, as discussed above, the library size that can be expressed on the surface of phage is limited by the efficiency of procedures such as cloning and transformation. Furthermore, the efficiency of incorporation of gIII-zinc finger fusions into phage and hence, zinc finger peptide expression, is determined by the number of zinc finger modules. Therefore, 2-finger peptides are expressed more efficiently than 3-finger peptides and so on. For this reason, phage display protocols are generally limited to the assay of polypeptides comprising 3 or fewer zinc finger modules.

An alternative to phage display is an *in vitro* selection system. In such a system, libraries of zinc fingers can be produced by PCR using degenerate primer oligonucleotides. Target binding sites are added to the end of the DNA encoding the zinc finger peptide. Zinc finger peptide expression may be performed directly from PCR products using an *in vitro* expression kit, such as the TNT T7 Quick Coupled Transcription/Translation System for PCR DNA (Promega, Madison, WI, USA), or another suitable expression system. The components of the expression reaction (including the zinc finger gene/binding site) are compartmentalised by suspension in an emulsion, in such a way that (on average) only one copy of the zinc finger gene / binding site is present in each compartment. See, for example, Tawfik, D.S. & Griffiths, A.D. (1998) *Nat. Biotechnol.* 16: 652-656. Zinc finger peptides which bind the specified target site (and the gene encoding them) can be collected using, for example, a suitable epitope tag (such as myc,

FLAG or HA tags), and the non-bound binding sites/zinc finger genes are removed. The genes encoding zinc finger peptides that bind the required target site can then be amplified by PCR and used in further rounds of selection if required.

- 5 A preferred method for selecting a zinc finger peptide which binds a specified target sequence is described in Example 4. Briefly, the DNA encoding a library of zinc finger peptides with an attached epitope tag is diluted into as many aliquots as it is possible to screen (e.g. 384 or 1534 aliquots). This creates pools of sub-libraries with reduced numbers of variants. The DNA is then amplified by PCR and used to produce protein,
10 from a suitable *in vitro* expression system, as described above. A specified binding site with an attached biotin molecule, and a horse radish peroxidase (HRP)-conjugated antibody to the peptide-attached epitope tag may then be added. Binding site / bound zinc finger / antibody complexes may be collected by binding to streptavidin and the samples are washed to remove unbound zinc finger and antibodies. The samples
15 containing the highest amount of bound zinc finger peptide can be detected by adding an HRP substrate solution. The original DNA stock from such positive samples may then be diluted into aliquots (as above), PCR-amplified and used for the next round of selection. In this way, pools of zinc finger encoding genes with the desired activity are isolated, subdivided into pools of reduced variation and re-isolated until the most active clone is
20 identified.

Principal advantages of the *in vitro* systems described above are: (a) there is virtually no limit to the library size which can be screened (up to 10^{12} different PCR products can easily be made); and (b) polypeptides comprising larger numbers of linked zinc finger
25 modules (e.g., 4, 5, 6, 7, or more) can be assayed. Another *in vitro* selection system which can be used is polysome/ribosome display. See, for example, Mattheakis, L.C., Bhatt, R.R. & Dower, W.J. (1994) *Proc. Natl. Acad. Sci. USA*. 91: 9022-9026; and WO 00/27878.

- 30 Protocols for the reverse selection procedure, *i.e.* the selection of a particular binding site from a mixed population using a single nucleic acid binding polypeptide, include SELEX (systematic evolution of ligands by exponential enrichment) and microarray techniques.

The SELEX procedure has been well described. See, for example, Drolet, D.W., Jenison, R.D., Smith, D.E., Pratt, D. & Hicke, B.J. (1999) *Comb. Chem. High Throughput Screen* 2: 271-278; Burden, D.A. & Osheroff, N. (1999) *J. Biol. Chem.* 274: 5227-5235; Shultzaberger, R.K. & Schneider, T.D. (1999) *Nucleic Acids Res.* 27: 882-887; Marozzi, A., Meneveri, R., Giacca, M., Gutierrez, M.I., Siccardi, A.G. & Ginelli, E. (1998) *J. Biotechnol.* 15: 117-128; and US Patents No. 5,270,163; 5,475,096; 5,595,877; 5,670,637; 5,696,249; 5,817,785 and 6,331,398. A single nucleic acid binding polypeptide is expressed, either *in vitro* or *in vivo*, and screened against a library of target sequences. Nucleic acid binding polypeptides are collected (along with any bound target sites) using an epitope tag (as above) or another suitable procedure. Bound target sites are amplified by PCR and may be used in further rounds of selection, to enrich for the optimal binding site, or sequenced.

Microarray technology provides a method of screening a particular polypeptide or nucleic acid against thousands to millions of target sequences on a single slid support such as, for example, a glass or nitrocellulose slide. For example, the members of a library encoding polypeptides comprising 2 linked zinc fingers will bind a 6 bp recognition sequence. Hence, there are 4096 ($=4^6$) unique binding sites for such a library. All 4096 of these sites can be arrayed onto a single glass slide, for example, allowing a specified 2-finger peptide to be screened simultaneously against every possible binding site. The amount of binding to each target sequence can be visualised and quantified using simple fluorescence measurements. For example, the zinc finger peptide may be expressed *in vitro*, or on the surface of phage. Isolated zinc finger peptides may contain an epitope tag for labelling purposes, whereas bound phage can be detected using a primary antibody against a phage coat protein, such as gVIII. A secondary antibody conjugated to, for example, R-phycoerythrin, horseradish peroxidase or alkaline phosphatase, can be used to provide a visible, quantifiable signal when a suitable substrate is applied. See, for example, Bulyk *et al.* (2001) *Proc. Natl. Acad. Sci. USA*:98:13, 7158-7163, which is incorporated, by reference, in its entirety.

Prediction of Binding Specificity

The screening approaches described above rely on the assay of large libraries of randomly-selected natural zinc finger modules, to obtain one or more zinc finger modules that optimally bind a particular target nucleic acid sequence. In order to simplify the process further and ensure a more rapid selection of optimal zinc finger modules for a particular target site, sub-libraries can be created. In this disclosure, the term 'sub-library' refers to a library of natural zinc finger modules that have been roughly categorised according to their predicted binding specificity. For example, the total population of natural zinc fingers can be sub-divided to create libraries comprising zinc finger modules whose predicted binding sites are guanine (G) rich, cytosine (C) rich, adenine (A) rich or thymine (T) rich. Alternatively, sub-libraries can be categorised as binding G in the 3' position, in the central position, or in the 5' position of a nucleotide triplet, etc. Alternatively, sub-libraries can be created which comprise zinc finger modules predicted to bind a particular triplet sequence such as, for example, GGG, GGA, GGC, GGT, GAG, GCG, GTG, etc. This approach combines knowledge of the modes of zinc finger-nucleic acid recognition, gained from studies on artificial zinc finger variants, with the benefits of combinatorial library selection. It also takes into account the fact that concerted interactions between adjacent zinc fingers, i.e. overlapping contacts, can affect the binding affinity and/or specificity of individual zinc fingers. *See*, for example, Isalan, M., Klug, A. & Choo, Y. (1998) Biochemistry 37, 12026-12033; Isalan, M., Choo, Y. & Klug, A. (1997) Proc Natl Acad Sci 94, 5617-5621. Thus, for example, a composite binding polypeptide comprising two fingers, each having a predicted binding specificity for a particular triplet, can be easily screened to determine if that pair of fingers are compatible with each other for binding to the 6-nucleotide target site comprising their individual target sequences. This strategy is described further in the Examples.

For the process of creating sub-libraries of natural zinc fingers according to predicted binding preference, the rules set forth in international patent applications WO 96/06166, WO 98/53057, WO 98/53058, WO 98/53059 and WO 98/53060, and described in more detail below, are used. These rules allow the assignment of an amino acid residue, in an

appropriate position of the recognition region of a zinc finger module (generally comprising amino acids -1 through +6, with respect to the start of the alpha-helical portion of the finger), which will bind a specified nucleotide in a triplet or quadruplet target subsite. However, these rules can also be used to predict the sequence of a target subsite that would be preferentially bound by a zinc finger of given amino acid sequence. In particular, the identity of the amino acid residing at a particular position in the recognition region of a natural zinc finger module can be used to predict the identity of a nucleotide at a particular location in a target subsite. These 'rules' should be considered as a guide to target site preference and not a guaranteed prediction, as binding site specificity may be determined by variations elsewhere in the zinc finger module (i.e. outside of the recognition region), may be influenced by context, or may be influenced by factors as yet unknown. It should also be noted that some rules may be more generally applicable than others.

In the application of these rules, it should be noted that the recognition region of a zinc finger aligns such that the N-terminal to C-terminal sequence of the finger is arranged along the nucleic acid strand to which it binds in a 3'-to-5' direction. As a result, when a zinc finger sequence and a nucleic acid sequence (to which the finger binds) are aligned, the primary interactions occur between the zinc finger and the 'minus' strand of the nucleic acid sequence (i.e. the strand which has a 3'-to-5' orientation). Furthermore, as stated above, the recognition region of a zinc finger comprises amino acids -1 through +6, with respect to the start of the alpha-helical portion of the finger. With respect to a particular zinc finger, an amino acid residue designated ++2 refers to the residue present in the adjacent (in the C-terminal direction) zinc finger, which (in certain instances) buttresses an amino acid-nucleotide interaction and/or participates in a cross-strand interaction with a nucleotide.

Thus, the following set of rules can be used to predict a 3 bp target subsite for a given natural zinc finger module: (a) if the 5' base in the triplet is G, then position +6 in the α -helix is Arg; or position +6 is Ser or Thr and position ++2 is Asp; (b) if the 5' base in the triplet is A, then position +6 in the α -helix is Gln and ++2 is not Asp; (c) if the 5' base in the triplet is T, then position +6 in the α -helix is Ser or Thr and position ++2 is Asp; (d) if

the 5' base in the triplet is C, then position +6 in the α -helix may be any amino acid, provided that position ++2 in the α -helix is not Asp; (e) if the central base in the triplet is G, then position +3 in the α -helix is His; (f) if the central base in the triplet is A, then position +3 in the α -helix is Asn; (g) if the central base in the triplet is T, then position +3
 5 in the α -helix is Ala, Ser or Val; provided that if it is Ala, then one of the residues at -1 or +6 is a small residue; (h) if the central base in the triplet is C, then position +3 in the α -helix is Ser, Asp, Glu, Leu, Thr or Val; (i) if the 3' base in the triplet is G, then position -1 in the α -helix is Arg; (j) if the 3' base in the triplet is A, then position -1 in the α -helix is Gln; (k) if the 3' base in the triplet is T, then position -1 in the α -helix is Asn or Gln; (l)
 10 if the 3' base in the triplet is C, then position -1 in the α -helix is Asp.

Furthermore, a natural zinc finger module may be capable of binding specifically to a four-nucleotide target subsite that overlaps with the target subsite of an adjacent zinc finger. In this case a different set of 'rules' can be used to determine predicted binding sites for each zinc finger module. Accordingly, in the description below, the overlapping
 15 4 bp binding site is described such that position 4 is the 5' base of a typical triplet binding site, position 3 is the central position of a typical triplet, position 2 is the 3' position of a typical triplet, and position 1 is the complement of the nucleotide which is contacted by the cross strand interaction from the +2 position of the zinc finger module. Position 1 can also be considered to be the 5' base of the triplet or quadruplet contacted by an adjacent
 20 (in the N-terminal direction) finger, if present.

Binding to each base of a quadruplet by an α -helical zinc finger nucleic acid binding motif in a natural protein can be predicted with reference to the following rules: (a) if base 4 in the quadruplet is G, then position +6 in the α -helix is Arg or Lys; (b) if base 4 in the quadruplet is A, then position +6 in the α -helix is Glu, Asn or Val; (c) if base 4 in the
 25 quadruplet is T, then position +6 in the α -helix is Ser, Thr, Val or Lys; (d) if base 4 in the quadruplet is C, then position +6 in the α -helix is Ser, Thr, Val, Ala, Glu or Asn; (e) if base 3 in the quadruplet is G, then position +3 in the α -helix is His; (f) if base 3 in the quadruplet is A, then position +3 in the α -helix is Asn; (g) if base 3 in the quadruplet is T, then position +3 in the α -helix is Ala, Ser or Val; provided that if it is Ala, then one of the

residues at -1 or +6 is a small residue; (h) if base 3 in the quadruplet is C, then position +3 in the α -helix is Ser, Asp, Glu, Leu, Thr or Val; (i) if base 2 in the quadruplet is G, then position -1 in the α -helix is Arg; (j) if base 2 in the quadruplet is A, then position -1 in the α -helix is Gln; (k) if base 2 in the quadruplet is T, then position -1 in the α -helix is His or Thr; (l) if base 2 in the quadruplet is C, then position -1 in the α -helix is Asp or His; (m) if base 1 in the quadruplet is G, then position +2 is Glu; (n) if base 1 in the quadruplet is A, then position +2 Arg or Gln; (o) if base 1 in the quadruplet is C, then position +2 is Asn, Gln, Arg, His or Lys; (p) if base 1 in the quadruplet is T, then position +2 is Ser or Thr.

- 10 The above rules may be further refined to those described below: (a) if base 4 in the quadruplet is G, then position +6 in the α -helix is Arg; or position +6 is Ser or Thr and position ++2 is Asp; (b) if base 4 in the quadruplet is A, then position +6 in the α -helix is Gln and ++2 is not Asp; (c) if base 4 in the quadruplet is T, then position +6 in the α -helix is Ser or Thr and position ++2 is Asp; (d) if base 4 in the quadruplet is C, then
- 15 position +6 in the α -helix may be any amino acid, provided that position ++2 in the α -helix is not Asp; (e) if base 3 in the quadruplet is G, then position +3 in the α -helix is His; (f) if base 3 in the quadruplet is A, then position +3 in the α -helix is Asn; (g) if base 3 in the quadruplet is T, then position +3 in the α -helix is Ala, Ser or Val; provided that if it is Ala, then one of the residues at -1 or +6 is a small residue; (h) if base 3 in the quadruplet
- 20 is C, then position +3 in the α -helix is Ser, Asp, Glu, Leu, Thr or Val; (i) if base 2 in the quadruplet is G, then position -1 in the α -helix is Arg; (j) if base 2 in the quadruplet is A, then position -1 in the α -helix is Gln; (k) if base 2 in the quadruplet is T, then position -1 in the α -helix is Asn or Gln; (l) if base 2 in the quadruplet is C, then position -1 in the α -helix is Asp; (m) if base 1 in the quadruplet is G, then position +2 is Asp; (n) if base 1 in the quadruplet is A, then position +2 is not Asp; (o) if base 1 in the quadruplet is C, then
- 25 position +2 is not Asp; (p) if base 1 in the quadruplet is T, then position +2 is Ser or Thr.

The rules therefore predict that the presence of an Asp (D) residue at position +2 will preclude binding to either A or C by an amino acid at position +6 in an adjacent N-

- 30 terminal finger. Isalan, M., Klug, A. & Choo, Y. (1998) Biochemistry 37, 12026-12033;

Isalan, M., Choo, Y. & Klug, A. (1997) Proc Natl Acad Sci 94, 5617-56212. Therefore, natural zinc fingers containing Asp, Glu, Asn or Gln at +6 are likely to be incompatible with any C-terminal finger containing an Asp residue at position +2. Although there are many such rules to describe the overlap between adjacent zinc fingers, a certain degree of degeneracy exists in these rules. Nonetheless, physical selection procedures (*e.g.*, library construction and screening) can be used to extract optimal pairs of fingers for any given target subsite interface.

Not all natural zinc fingers have a DNA-binding function. For example, it is known that many zinc fingers, such as those from TFIIIA, bind to RNA (Clemens, K. R. *et al.*, (1993) *Science* 260: 530-533; Bogenhagen, D.F. (1993) *Mol. Cell. Biol.* 13: 5149-5158; Searles, M. A. *et al.*, *J. Mol. Biol.* 301: 47-60 (2000)). The rules governing RNA binding by zinc fingers are less well understood than those of DNA binding, but some RNA binding zinc fingers can be identified on the basis of a characteristic sequence motif. Clemens, K. R. *et al.*, (1993) *Science* 260: 530-533; Bogenhagen, D.F. (1993) *Mol. Cell. Biol.* 13: 5149-5158; Searles, M. A. *et al.* (2000) *J. Mol. Biol.* 301: 47-60. Furthermore, some zinc fingers, such as those from the protein Ikaros, are able to form protein-protein interactions. Such zinc fingers often contain large hydrophobic patches. Mackay, J. P. & Crossley, M. (1998) *Trends Biochem. Sci.* 23: 1-4.

To this end, applied bioinformatic processing can help to determine which candidates in a particular genome are best suited to fulfilling a particular function, such as DNA-binding. In the case of zinc fingers, numerous documented databases exist denoting amino acid residues that are most likely to be found at particular positions within a DNA-binding zinc finger. See, for example, Isalan, M., Klug, A. & Choo, Y. (1998) Biochemistry 37, 12026-12033; Choo, Y. & Klug, A. (1997) Curr. Opin. Str. Biol. 7, 117-125; WO 98/53060; WO 98/53059; WO 98/53058. As an example, disclosed herein is a database of approximately 200 natural human zinc fingers which have been selected (on the basis of coded contacts) as having potentially useful DNA-binding activity (see Example 1). Also disclosed in Example 1 are the predicted DNA target sequences of these zinc fingers, assigned according to the rules set out above.

As the human genome contains almost 700 zinc finger-containing proteins, there are many other candidates that can be included in a more inclusive library of natural zinc fingers. A selection of these are disclosed in Example 2.

- 5 Similar work can be carried out in other organisms, such as farm (cows, pigs, sheep, chickens, etc.), laboratory (monkeys, rats, mice, etc.) and domestic (dogs, cats, etc.) animals. In this case, it is necessary to select natural zinc finger modules from the respective genomes of such organisms. Examples of zinc finger modules which have been selected from mouse, chicken and certain plant genomes, are disclosed in
10 Example 3.

d. Zinc Finger Chimeric Peptides

In a preferred embodiment, the composite binding polypeptides described herein comprise chimeric nucleic acid binding polypeptides.

- 15 A chimeric nucleic acid binding polypeptide, also referred to as a fusion polypeptide, comprises a binding domain (comprising a number of nucleic acid binding polypeptide modules or fingers) designed to bind specifically to a target nucleotide sequence, together with one or more further biological effector domains or functional domains. The terms "biological effector domain" and "functional domain" refer to any
20 polypeptide (of functional fragment thereof) that has a biological function. Included are enzymes, receptors, regulatory domains, transcriptional activation or repression domains, binding sequences, dimerisation, trimerisation or multimerisation sequences, sequences involved in protein transport, localisation sequences such as subcellular localisation sequences, nuclear localisation, protein targeting or signal sequences. Furthermore,
25 biological effector domains may comprise polypeptides involved in chromatin remodelling, chromatin condensation or decondensation, DNA replication, transcription, translation, protein synthesis, etc. Fragments of such polypeptides comprising the relevant activity (*i.e.*, functional fragments) are also included in this definition. Preferred biological effector domains include transcriptional modulation domains such as

transcriptional activators and transcriptional repressors, as well as their functional fragments.

The effector domain(s) can be covalently or non-covalently attached to the binding domain.

5 Chimeric nucleic acid binding polypeptides preferably comprise transcription factor activity, for example, a transcriptional modulation activity such as transcriptional activation or transcriptional repression activity. For example, a zinc finger chimeric polypeptide may comprise a binding domain designed to bind specifically to a particular nucleotide sequence, and one or more further biological effector domains, preferably a
10 transcriptional activation or repression domain, as described in further detail below. The zinc finger chimeric polypeptide may comprise one or more zinc fingers or zinc finger binding modules.

Preferably, in the case of a chimeric polypeptide comprising transcriptional modulation activity, a nuclear localisation domain is attached to the DNA binding domain
15 to direct the chimeric polypeptide to the nucleus.

Generally, a chimeric nucleic acid binding polypeptide, such as a chimeric zinc finger polypeptide, can also include an effector domain to regulate gene expression. The effector domain can be directly derived from a basal or regulated transcription factor such as, for example, transactivators, repressors, and proteins that bind to insulator or silencer
20 sequences. *See, for example, Choo & Klug (1995) Curr. Opin. Biotech. 6: 431-436; Choo, Y. & Klug, A. (1997) Curr. Opin. Str. Biol. 7, 117-125; Rebar & Pabo (1994) Science 263: 671-673; Jamieson et al. (1994) Biochem. 33: 5689-5695; Goodrich et al (1996) Cell 84: 825-830; Vostrov, A. A. & Quitschke, W. W. (1997) J. Biol. Chem. 272: 33353-33359 and WO 00/41566 and references disclosed therein. Other useful domains
25 are derived from receptors such as, for example, nuclear hormone receptors (Kumar, R. & Thompson, E. B. (1999) Steroids 64: 310-319), and their co-activators and co-repressors (Ugai, H. et al. (1999) J. Mol. Med. 77: 481-494).*

A chimeric nucleic acid binding polypeptide can also include other domains that may be advantageous within the context of the control of gene expression. Such domains include, but are not limited to, protein-modifying domains such as histone acetyltransferases, kinases, methylases and phosphatases, which can silence or activate genes by modifying DNA structure or the proteins that associate with nucleic acids. *See*, for example, Wolffe, *Science* 272: 371-372 (1996); Taunton *et al.*, *Science* 272: 408-411 (1996); Hassig *et al.*, *Proc. Natl. Acad. Sci. USA* 95: 3519-3524 (1998); Wang, *Trends Biochem. Sci.* 19: 373-376 (1994); and Schonthal & Semin, *Cancer Biol.* 6: 239-248 (1995). Additional useful effector domains include those that modify or rearrange nucleic acid molecules such as methyltransferases, endonucleases, ligases, recombinases etc. *See*, for example, Wood, *Ann. Rev. Biochem.* 65: 135-167 (1996); Sadowski, *FASEB J.* 7: 760-767 (1993); Cheng, *Curr. Opin. Struct. Biol.* 5: 4-10 (1995); Wu *et al.* (1995) *Proc. Natl. Acad. Sci. USA* 92:344-348; Nahon & Raveh, *Nucleic Acids Res* 1998 Mar 1;26(5):1233-9; Smith *et al.* *Nucleic Acids Res.* 1999 Jan 15;27(2):674-81; and Smith *et al.* (2000) *Nucleic Acids Res.* Sept 1; 28(17):3361-9. It will be appreciated that the biological effector domain portion of the chimeric polypeptide may itself also comprise such activities, without the need for further additional domains.

For the purpose of gene activation, zinc finger domains may be fused to the VP64 domain. *See*, for example, Seipel *et al.*, *EMBO J.* 11: 4961-4968 (1996). Other preferred transactivator domains include the herpes simplex virus (HSV) VP16 domain (Hagmann *et al.* (1997) *J. Virol.* 71: 5952-5962; Sadowski *et al.* (1988) *Nature* 335:563-564), transactivation domain 1 and/or domain 2 of the p65 subunit of nuclear factor- κ B (NF- κ B (Schmitz, M. L. *et al.* (1995) *J. Biol. Chem.* 270: 15576-15584). Other transcription factors are reviewed in, for example, Lekstrom-Himes J. & Xanthopoulos K. G. (C/EBP family) *J. Biol. Chem.* 273: 28545-28548 (1998); Bieker, J. J. *et al.*, (globin gene transcription factors) *Ann. N. Y. Acad. Sci.* 850: 64-69 (1998), and Parker, M. G. (estrogen receptors) *Biochem. Soc. Symp.* 63: 45-50 (1998).

Use of a transactivation domain from the estrogen receptor is disclosed in Metivier, R., Petit, FG., Valotaire, Y. & Pakdel, F. (2000) *Mol. Endocrinol.* 14: 1849-1871. Furthermore, activation domains from the globin transcription factors EKLF

(Pandya, K. Donze, D. & Townes T. (2001) *J. Biol. Chem.* 276: 8239-8243) may also be used, as well as a transactivation domain from FKLf (Asano, H. Li, XS. & Stamatoyannopoulos, G. (1999) *Mol. Cell. Biol.* 19: 3571-3579). C/EPB transactivation domains may also be employed in the methods described herein. The C/EBP epsilon
5 activation domain is disclosed in Verbeek, W., Gombart, AF, Chumakov, AM, Muller, C, Friedman, AD, & Koeffler, HP (1999) *Blood* 15: 3327-3337. Kowenz-Leutz, E. & Leutz, A. (1999) *Mol. Cell.* 4: 735-743 disclose the use of the C/EBP tau activation domain, while the C/EBP alpha transactivation domain is disclosed in Tao, H., & Umek, RM. (1999) *DNA Cell Biol.* 18: 75-84.

10 It is known that zinc finger proteins may be fused to transcriptional repression domains such as the Kruppel-associated box (KRAB) domain to form powerful repressors. These domains are known to repress expression of a reporter gene even when bound to sites a few kilobase pairs upstream from the promoter of the gene (Margolin *et al.*, 1994, *Proc. Natl. Acad. Sci. USA* 91: 4509-4513). Hence, in certain embodiments,
15 the KRAB repressor domain from the human KOX-1 protein is used to repress gene activity (Moosmann *et al.*, *Biol. Chem.* 378: 669-677 (1997); Thiesen *et al.*, *New Biologist* 2: 363-374 (1990)). In additional embodiments, larger fragments of the KOX-1 protein comprising the KRAB domain, up to and including full-length KOX protein, are used as transcriptional repression domains. See, for example, Abrink *et al.* (2001) *Proc.*
20 *Natl. Acad. Sci. USA* 98:1422-1426. Other preferred transcriptional repressor domains are known in the art and include, for example, the *engrailed* domain (Han *et al.*, *EMBO J.* 12: 2723-2733 (1993)), the *snag* domain (Grimes *et al.*, *Mol Cell. Biol.* 16: 6263-6272 (1996)) and the transcriptional repression domain of v-erbA (*e.g.*, Urnov *et al.* (2000) *EMBO J.* 19:4074-4090; Sap *et al.* (1989) *Nature* 340:242-244 and Ciana *et al.* (1999) *EMBO J.* 17:7382-7394).
25

Biological effector domains can be covalently or non-covalently linked to a binding domain. In one embodiment, a covalent linker comprises a flexible amino acid sequence; fusion polypeptides according to this embodiment comprise a nucleic acid binding domain fused, by an amino acid linker, to a biological effector domain.

30 Alternatively, a covalent linker may comprise a synthetic, non-amino acid based,

chemical linker, for example, polyethylene glycol. Synthetic linkers are commercially available, and methods of chemical conjugation are known in the art. Covalent linkers may comprise flexible or structured linkers, as described above.

Non-covalent linkages between a nucleic acid binding domain and an effector domain can be formed using, for example, leucine zipper/coiled coil domains, or other naturally occurring or synthetic dimerisation domains. *See e.g.*, Luscher, B. & Larsson, L. G. *Oncogene* 18:2955-2966 (1999) and Gouldson, P. R. *et al.*, *Neuropsychopharmacology* 23: S60-S77 (2000).

The expression of composite binding polypeptides (for example, zinc finger polypeptides) can be controlled by tissue specific promoter sequences such as, for example, the *lck* promoter (thymocytes, Gu, H. *et al.*, *Science* 265: 103-106 (1994)); the human CD2 promoter (T-cells and thymocytes, Zhumabekov, T. *et al.*, *J. Immunological Methods* 185: 133-140 (1995)); the alpha A-crystallin promoter (eye lens, Lakso, M. *et al.*, *Proc. Natl. Acad. Sci.* 89: 6232-6236 (1992)); the alpha-calcium-calmodulin-dependent kinase II promoter (hippocampus and neocortex, Tsien, J. *et al.*, *Cell* 87: 1327-1338 (1996)); the whey acidic protein promoter (mammary gland, Wagner, K.-U. *et al.*, *Nucleic Acids Res.* 25: 4323-4330 (1997)); the aP2 enhancer/promoter (adipose tissue, Barlow C. *et al.*, *Nucleic Acids Res.* 25: 2543-2545 (1997)); the aquaporin-2 promoter (renal collecting duct, Nelson R. *et al.*, *Am. J. Physiol.* 275: C216-C226 (1998)); and the mouse myogenin promoter (skeletal muscle, Grieshammer, U. *et al.*, *Dev. Biol.* 197: 234-247 (1998)). The expression of such polypeptides can also be controlled by inducible systems, in particular, controlled by small molecule induction such as the tetracycline-controlled systems (tet-on and tet-off), the RU-486 or tamoxifen hormone analogue systems, or the radiation-inducible early growth response gene-1 (EGR1) promoter. These promoter constructs and inducible systems have the benefit of being able to provide organ-specific and/or inducible expression of target genes for use in applications such as gene therapy and transgenic animals.

e. **Vectors**

The nucleic acid encoding the nucleic acid binding polypeptide such as a zinc finger polypeptide can be incorporated into intermediate vectors and transformed into prokaryotic or eukaryotic cells for expression or DNA amplification.

5 As used herein, vector (or plasmid) preferably refers to discrete elements that are used to introduce heterologous nucleic acid into cells for either expression or replication thereof. The term "heterologous to the cell" means that the sequence does not naturally exist in the genome of the host cell but has been introduced into the cell. The term "introduced into" means that a procedure is performed on a cell, tissue, organ or organism
10 such that the gene encoding the nucleic acid binding polypeptide (for example, a zinc finger polypeptide) previously absent from the cell or cells is then present in the cell or cells. Alternatively, or in addition, the gene may be initially present in the cell or cells and subsequently altered by introduction of heterologous DNA. A heterologous sequence may include a modified sequence introduced at any chromosomal site, or which is not
15 integrated into a chromosome, or which is introduced by homologous recombination such that it is present in the genome in the same position as the native allele. Selection and use of such vectors are well within the skill of the person of ordinary skill in the art. Many vectors are available, and selection of an appropriate vector will depend on the intended use of the vector, i.e. whether it is to be used for DNA amplification or for nucleic acid
20 expression, the size of the DNA to be inserted into the vector, and the host cell to be transformed with the vector, etc. Another consideration is whether the vector is to remain episomal or integrate into the host genome. Suitable vectors may be of bacterial, viral, insect or mammalian origin. Intermediate vectors for storage or manipulation of the nucleic acid encoding the nucleic acid binding polypeptide, or for expression and
25 purification of the polypeptide are typically of prokaryotic origin. Most expression vectors are shuttle vectors, i.e. they are capable of replication in at least one class of organisms but can be transfected into another class of organisms for expression. For example, a vector is cloned in *E. coli* and then the same vector is transfected into yeast or mammalian cells even though it is not capable of replicating independently of the host
30 cell chromosome. DNA may also be replicated by insertion into the host genome. The

nucleic acid binding polypeptides such as zinc finger polypeptides described here are preferably inserted into a vector suitable for expression in mammalian cells.

Prokaryote, yeast and higher eukaryote cells may be used for replicating DNA and producing the nucleic acid binding protein. Suitable prokaryotes include eubacteria, such as Gram-negative or Gram-positive organisms, such as *E. coli*, e.g. *E. coli* K-12 strains, DH5a and HB101, or Bacilli. Further hosts suitable for the vectors include eukaryotic microbes such as filamentous fungi or yeast, e.g. *Saccharomyces cerevisiae*. Higher eukaryotic cells include insect and vertebrate cells, particularly mammalian cells including human cells or nucleated cells from other multicellular organisms. In recent years propagation of vertebrate cells in culture (tissue culture) has become a routine procedure. Examples of useful mammalian host cell lines are epithelial or fibroblastic cell lines such as Chinese hamster ovary (CHO) cells, NIH 3T3 cells, HeLa cells or 293T cells. The host cells referred to in this disclosure comprise cells in *in vitro* culture as well as cells that are within a host animal.

Each vector contains various components depending on its function (amplification of DNA or expression of DNA) and the host cell for which it is compatible. The vector components generally include, but are not limited to, one or more of the following: an origin of replication, one or more selectable marker genes, a promoter, an enhancer element, a transcription termination sequence and a signal sequence.

Both expression and cloning vectors generally contain nucleic acid sequence that enable the vector to replicate in one or more selected host cells. Typically in cloning vectors, this sequence is one that enables the vector to replicate independently of the host chromosomal DNA, and includes origins of replication or autonomously replicating sequences. Such sequences are well known for a variety of bacteria, yeast and viruses. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria, the 2 μ plasmid origin is suitable for yeast, and various viral origins (e.g. SV 40, polyoma, adenovirus) are useful for cloning vectors in mammalian cells. Generally, the origin of replication component is not needed for mammalian expression vectors unless

these are used in mammalian cells competent for high level DNA replication, such as COS cells.

Advantageously, an expression and cloning vector contains a selection gene also referred to as selectable marker. This gene encodes a protein necessary for the survival or growth of transformed host cells grown in a selective culture medium. Host cells not transformed with the vector containing the selection gene will not survive in the culture medium. Typical selection genes encode proteins that confer resistance to antibiotics and other toxins, e.g. ampicillin, neomycin, methotrexate or tetracycline, complement auxotrophic deficiencies, or supply critical nutrients not available from complex media.

Since the replication of vectors is conveniently done in *E. coli*, an *E. coli* genetic marker and an *E. coli* origin of replication are advantageously included. These can be obtained from *E. coli* plasmids, such as pBR322, Bluescript© vector or a pUC plasmid, e.g. pUC18 or pUC19, which contain both *E. coli* replication origin and *E. coli* genetic marker conferring resistance to antibiotics, such as ampicillin and tetracycline. Vectors such as these are commercially available.

As to a selective gene marker appropriate for yeast, any marker gene can be used which facilitates the selection for transformants due to the phenotypic expression of the marker gene. Suitable markers for yeast are, for example, those conferring resistance to antibiotics G418, hygromycin or bleomycin, or provide for prototrophy in an auxotrophic yeast mutant, for example the URA3, LEU2, LYS2, TRP1, or HIS3 gene.

Suitable selectable markers for mammalian cells are those that enable the identification of cells competent to take up nucleic acid, such as dihydrofolate reductase (DHFR, methotrexate resistance), thymidine kinase, or genes conferring resistance to neomycin, G418 or hygromycin. The mammalian cell transformants are placed under selection pressure which only those transformants which have taken up and are expressing the marker are uniquely adapted to survive. In the case of a DHFR or glutamine synthase (GS) marker, selection pressure can be imposed by culturing the transformants under conditions in which the pressure is progressively increased, thereby

leading to amplification (at its chromosomal integration site) of both the selection gene and the linked DNA that encodes the nucleic acid binding protein. Amplification is the process by which genes in greater demand (such as one encoding a protein that is critical for growth), together with closely associated genes (such as one encoding a composite
5 binding polypeptide), are reiterated in tandem within the chromosomes of recombinant cells. Increased quantities of desired protein are usually synthesised from this amplified DNA.

Expression and cloning vectors usually contain control sequences that are recognised by the host organism and are operably linked to the nucleic acid encoding a
10 nucleic acid binding polypeptide. The term "control sequences" is intended to include, at a minimum, components whose presence can influence expression, and can also include additional components whose presence is advantageous, for example, leader sequences and fusion partner sequences. The term "operably linked" means that the components described are in a relationship permitting them to function in their intended manner.
15 Typical control sequences include promoters, enhancers and other expression regulation signals such as terminators. Such a promoter may be inducible or constitutive. A regulatory sequence operably linked to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under conditions compatible with the control sequences.

20 The term promoter is well known in the art and encompasses nucleic acid regions ranging in size and complexity from minimal promoters to promoters including upstream elements and enhancers. Suitable promoters for use in prokaryotic and eukaryotic cells are well known in the art, and described in for example, Current Protocols in Molecular Biology (Ausubel *et al.*, eds., 1994) and Molecular Cloning. A Laboratory Manual
25 (Sambrook *et al.*, 2nd ed. 1989).

Promoters suitable for use with prokaryotic hosts include, for example, the β -lactamase and lactose promoter systems, alkaline phosphatase, the tryptophan (Trp) promoter system and hybrid promoters such as the tac promoter. Their nucleotide sequences have been published, thereby enabling the skilled worker to ligate them to

DNA encoding a composite binding protein, using linkers or adapters to supply any required restriction sites. Promoters for use in bacterial systems will also generally contain an adjacent ribosome binding site (*e.g.*, a Shine-Dalgarno sequence) operably linked to the DNA encoding the composite binding polypeptide.

5 Preferred expression vectors are bacterial expression vectors, which comprise a promoter of a bacteriophage such as phage lambda, SP6, T3 or T7, for example, which is capable of functioning in bacteria. In one of the most widely used expression systems, the nucleic acid encoding the fusion protein can be transcribed from a vector by T7 RNA polymerase (Studier *et al*, *Methods in Enzymol.* 185: 60-89, 1990). In the *E. coli*
10 BL21(DE3) host strain, used in conjunction with pET vectors, the T7 RNA polymerase is produced from the λ -lysogen DE3 in the host bacterium, and its expression is under the control of the IPTG inducible lac UV5 promoter. This system has been employed successfully for over-production of many proteins. Alternatively, the polymerase gene may be introduced on a lambda phage by infection with an int^- phage such as the CE6
15 phage, which is commercially available (Novagen, Madison, WI, USA). Other vectors include vectors containing the lambda P_L promoter such as PLEX (Invitrogen, NL), vectors containing the *trc* promoters such as pTrcHisXpressTm (Invitrogen), or pTrc99 (Pharmacia Biotech, SE), or vectors containing the *tac* promoter such as pKK223-3 (Pharmacia Biotech), or PMAL (New England Biolabs, Beverly, MA, USA). A suitable
20 vector for expression of proteins in mammalian cells is the CMV enhancer-based vector such as pEVRF (Matthias, *et al.*, (1989) *Nucleic Acids Res.* 17, 6418).

Suitable promoting sequences for use with yeast hosts may be regulated or constitutive and are preferably derived from a highly expressed yeast gene, especially a *Saccharomyces cerevisiae* gene. Thus, the promoter of the TRP1 gene, the ADHI or
25 ADHII gene, the acid phosphatase (PH05) gene, a promoter of the yeast mating pheromone genes coding for the α - or α -factor or a promoter derived from a gene encoding a glycolytic enzyme such as the promoter of the enolase, glyceraldehyde-3-phosphate dehydrogenase (GAP), 3-phosphoglycerate kinase (PGK), hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-
30 phosphoglycerate mutase, pyruvate kinase, triose phosphate isomerase, phosphoglucose

isomerase or glucokinase genes, or a promoter from the TATA binding protein (TBP) gene can be used. Furthermore, it is possible to use hybrid promoters comprising upstream activation sequences (UAS) of one yeast gene and downstream promoter elements including a functional TATA box of another yeast gene, for example a hybrid promoter including the UAS(s) of the yeast PH05 gene and downstream promoter elements including a functional TATA box of the yeast GAP gene (PH05-GAP hybrid promoter). A suitable constitutive PH05 promoter is, for example, a shortened acid phosphatase PH05 promoter devoid of the upstream regulatory elements (UAS) such as the PH05 (-173) promoter element starting at nucleotide -173 and ending at nucleotide -9 of the PH05 gene.

The promoter is typically selected from promoters which are found in animal cells, although prokaryotic promoters and promoters functional in other eukaryotic cells can be used. Typically, the promoter is derived from viral or animal gene sequences, may be constitutive or inducible, and may be strong or weak.

Viral promoters can be derived from viruses such as polyoma virus, adenoviruses, adeno-associated viruses, poxviruses (*e.g.*, fowlpox virus), papilloma viruses (*e.g.*, BPV), avian sarcoma virus, cytomegalovirus (CMV), herpesviruses, retroviruses, lentiviruses and simian virus 40 (SV40). An example of a relatively weak viral promoter is thymidine kinase promoter from herpes simplex virus (HSV-TK).

Mammalian derived promoters can be heterologous to the animal in which composite binding polypeptide (such as zinc finger polypeptide) expression is to occur, or they can be host sequences. In some applications it is preferable to use a promoter that is active in all cell types, however it is often preferable to use promoter sequences that are active in specific cell types only.

The actin promoter and the strong ribosomal protein promoter are examples of promoter sequences that are active in all cell types. In contrast, by using promoters that are specific for certain cell or tissue types, the gene encoding the nucleic acid binding polypeptide can be expressed only in the required cell or tissue types. This may be of

extreme importance for applications such as gene therapy, and for the production of viable transgenic animals. Such promoters are known in the art and include the *lck* promoter (thymocytes, Gu, H. *et al.*, *Science* 265: 103-106 (1994)), the human CD2 promoter (T-cells and thymocytes, Zhumabekov, T. *et al.*, *J. Immunological Methods* 185: 133-140 (1995)); the alpha A-crystallin promoter (eye lens, Lakso, M. *et al.*, *Proc. Natl. Acad. Sci.* 89: 6232-6236 (1992)), the alpha-calcium-calmodulin-dependent kinase II promoter (hippocampus and neocortex, Tsien, J. *et al.*, *Cell* 87: 1327-1338 (1996)), the whey acidic protein promoter (mammary gland, Wagner, K.-U. *et al.*, *Nucleic Acids Res.* 25: 4323-4330 (1997)), the aP2 enhancer/promoter (adipose tissue, Barlow C. *et al.*, *Nucleic Acids Res.* 25: 2543-2545 (1997)), the aquaporin-2 promoter (renal collecting duct, Nelson R. *et al.*, *Am. J. Physiol.* 275: C216-C226 (1998)), the mouse myogenin promoter (skeletal muscle, Grieshammer, U. *et al.*, *Dev. Biol.* 197: 234-247 (1998)), retinoblastoma gene promoter (nervous system, Jiang, Z. *et al.*, *J. Biol. Chem.* 276: 593-600 (2001)).

The expression of nucleic acid binding polypeptides such as zinc finger polypeptides can also be controlled by small molecule induction or other inducible systems such as the tetracycline inducible systems (tet-on and tet-off), the RU-486 or tamoxifen hormone analogue systems, or the radiation-inducible early growth response gene-1 (EGR1) promoter, all of which are commercially available. By using such inducible promoter systems, transgenic lines can be established which carry a zinc finger chimeric polypeptide but express it only after addition of an inducer molecule. Thus the genes encoding the zinc finger polypeptides or other nucleic acid binding polypeptides can be expressed (or not expressed) in response to the small molecule, which can be easily administered. These systems may also allow the time and amount of polypeptide expression to be regulated.

Expression vectors typically contain expression cassettes that carry all the additional elements required for efficient expression of the nucleic acid in the host cell. Additional elements are enhancer sequences, polyadenylation and transcriptional termination signals, ribosome binding sites, and translational termination sequences.

Transcription of DNA by higher eukaryotes may be increased by inserting an enhancer sequence into the vector. Enhancers are relatively orientation and position independent. Many enhancer sequences are known from mammalian genes (e.g. elastase and globin). However, typically one will employ an enhancer from a eukaryotic cell virus. Examples include the SV40 enhancer on the late side of the replication origin (approx. bp 100-270) and the CMV early promoter enhancer. The enhancer may be spliced into the vector at a position 5' or 3' to the gene encoding the zinc finger polypeptide or nucleic acid binding polypeptide, but is preferably located at a site 5' from the promoter.

It has also been shown that the expression of a heterologous gene in an animal cell may be enhanced by retaining intron sequences (as opposed to using a cDNA clone). For example, intron 1 of the human CD2 gene has been shown to enhance the level of expression of CD2 in human cells (Festenstein, R. *et al.* 1996 *Science* 271: 1123).

Advantageously, a eukaryotic expression vector encoding a nucleic acid binding protein may comprise a locus control region (LCR). LCRs are capable of directing high-level integration site-independent expression of transgenes integrated into host cell chromatin. This is particularly important where the gene encoding the zinc finger polypeptide or the nucleic acid binding polypeptide is to be expressed over extended periods of time, for applications such as transgenic animals and gene therapy, as gene silencing of integrated heterologous DNA – especially of viral origin – is known to occur (Palmer, T. D. *et al.*, *Proc. Natl. Acad. Sci. USA* 88: 1330-1334 (1991); Harpers, K. *et al.*, *Nature* 293: 540-542 (1981); Jahner, D. *et al.*, *Nature* 298: 623-628 (1992); and Chen, W. Y. *et al.*, *Proc. Natl. Acad. Sci. USA* 94: 5798-5803 (1997)). Typical LCRs are exemplified by the human β -globin cluster, and the HS-40 regulatory region from the α -globin locus.

Eukaryotic vectors may also contain sequences necessary for the termination of transcription and for stabilising the mRNA transcript. Such sequences are commonly available from the 5' and 3' untranslated regions of eukaryotic or viral DNAs, and are known in the art. These regions contain nucleotide segments transcribed as

polyadenylated fragments in the untranslated portion of the mRNA encoding the relevant polypeptide. An appropriate terminator of transcription is fused downstream of the gene encoding the selected nucleic acid binding polypeptide such as a zinc finger protein. Any of a number of known transcriptional terminator, RNA polymerase pause sites and polyadenylation enhancing sequences can be used at the 3' end of the nucleic acid encoding for example a zinc finger polypeptide (see, for example, Richardson, J. P. *Crit. Rev. Biochem. Mol. Biol.* 28:1-30 (1993); Yonaha M. & Proudfoot, N. J. *EMBO J.* 19: 3770-3777 (2000); Ashfield, R. *et al.*, *EMBO J.* 10: 4197-4207 (1991); Hirose, Y. & Manley, J. L. *Nature* 395: 93-96 (1998)).

The nucleic acid binding polypeptides are generally targeted to the cell nucleus so that they are able to interact with host cell DNA and bind to the appropriate DNA target in the nucleus and regulate transcription. To effect this, a nuclear localisation sequence (NLS) is incorporated in frame with the expressible nucleic acid binding polypeptide (e.g., zinc finger polypeptide) gene construct. The NLS can be fused either 5' or 3' to the sequence encoding the binding protein, but preferably it is fused to the C-terminus of the chimeric polypeptide.

The NLS of the wild-type Simian Virus 40 Large T-Antigen (Kalderon *et al.* (1984) *Cell* 37: 801-813; and Markland *et al.* (1987) *Mol. Cell. Biol.* 7: 4255-4265) is an appropriate NLS and provides an effective nuclear localisation mechanism in animals. However, several alternative NLSs are known in the art and can be used instead of the SV40 NLS sequence. These include the NLSs of TGA-1A and TGA-1B.

Composite binding polypeptides can comprise tag sequences to facilitate studies and/or preparation of such molecules. Tag sequences may include FLAG-tags, myc-tags, 6his-tags, hemagglutinin tags or any other suitable tag known in the art.

Moreover, the nucleic acid binding protein gene according to the invention preferably includes a secretion sequence in order to facilitate secretion of the polypeptide from bacterial hosts, such that it will be produced as a soluble native peptide rather than

in an inclusion body. The peptide may be recovered from the bacterial periplasmic space, or the culture medium, as appropriate.

Construction of vectors employs conventional ligation techniques. Isolated
5 plasmids or DNA fragments are cleaved, tailored, and religated in the form desired to generate the plasmids required. If desired, analysis to confirm correct sequences in the constructed plasmids is performed in a known fashion. Suitable methods for constructing expression vectors, preparing in vitro transcripts, introducing DNA into host cells, and performing analyses for assessing nucleic acid binding protein expression and function
10 are known to those skilled in the art. Gene presence, amplification and / or expression may be measured in a sample directly, for example, by conventional Southern blotting, Northern blotting to quantify the transcription of mRNA, dot blotting (DNA or RNA analysis), or in situ hybridisation, using an appropriately labelled probe which may be based on a sequence provided herein. Those skilled in the art will readily envisage how
15 these methods may be modified, if desired.

f. Applications of Composite Binding Polypeptides

20 Nucleic acid binding proteins according to the invention can be employed in a wide variety of applications, including diagnostics and as research tools, and also in therapeutic applications and in transgenic organisms.

In Vitro Applications

25 Poly-zinc finger peptides of this invention may be employed as diagnostic tools for identifying the presence of nucleic acid molecules in a complex mixture. Nucleic acid binding molecules according to the invention can differentiate single base pair changes in target nucleic acid molecules.

Accordingly, the invention provides methods for determining the presence of a target nucleic acid molecule, wherein the target nucleic acid molecule comprises a target sequence, comprising the steps of:

- 5 a) preparing a nucleic acid binding protein, by a method set forth above, which is specific for the target nucleic acid sequence;
 - b) exposing a test system to the nucleic acid binding protein under conditions which promote binding of the protein to the target sequence, and removing any nucleic acid binding protein which remains unbound;
 - 10 c) testing for the presence of the nucleic acid binding protein in the test system;
- wherein, if the nucleic acid binding protein is detected, the target nucleic acid molecule is present and, if the nucleic acid binding protein is not detected, the target nucleic acid molecule is not present. In additional embodiments, quantitation of the amount of nucleic acid binding protein allows quantitation of the amount of the target nucleic acid molecule
- 15 present in the test system.

In a preferred embodiment, the nucleic acid binding molecules of the invention can be incorporated into an ELISA assay. For example, phage displaying composite binding polypeptides can be used to detect the presence of the target nucleic acid, and visualised

20 using enzyme-linked anti-phage antibodies.

Further improvements to the use of phage expressing a composite binding polypeptide for diagnosis can be made, for example, by co-expressing a marker protein fused to the minor coat protein (gVIII) of a filamentous bacteriophage. Since detection with an anti-phage

25 antibody would then be unnecessary, the time and cost of each diagnosis would be further reduced. Depending on the requirements, suitable markers for display might include fluorescent proteins (A. B. Cubitt, *et al.*, (1995) *Trends Biochem Sci.* **20**, 448-455; T. T. Yang, *et al.*, (1996) *Gene* **173**, 19-23), or an enzyme such as alkaline phosphatase (J. McCafferty, R. H. Jackson, D. J. Chiswell, (1991) *Protein Engineering* **4**, 955-961).

30 Labelling different types of diagnostic phage with distinct markers would allow multiplex screening of a single nucleic acid sample. Nevertheless, even in the absence of such refinements, the basic ELISA technique is reliable, fast, simple and particularly

inexpensive. Moreover it requires no specialised apparatus, nor does it employ hazardous reagents such as radioactive isotopes, making it amenable to routine use in the clinic. The major advantage of the protocol is that it obviates the requirement for gel electrophoresis, and so opens the way to automated nucleic acid diagnosis.

5

The invention provides nucleic acid binding proteins that have exquisite specificity. The invention lends itself, therefore, to the design of any molecule of which specific nucleic acid binding is required. For example, the proteins according to the invention may be employed in the manufacture of chimeric restriction enzymes, in which a nucleic acid
10 cleaving domain is fused to a nucleic acid binding domain comprising a zinc finger as described herein.

In Vivo Applications

15 The invention further provides composite binding polypeptides (and nucleic acids encoding them) that may be used in transgenic organisms (such as non-human animals), as therapeutic agents, and in gene therapy applications.

A transgenic animal is an animal, preferably a non-human animal, containing at least one foreign gene, called a transgene, in its genetic material. Preferably, the transgene is
20 contained in the animal's germ line such that it can be transmitted to the animal's offspring. Transgenic animals may carry the transgene in all their cells or may be genetically mosaic.

Constructs useful for creating transgenic animals according to the invention comprise genes encoding nucleic acid binding polypeptides, optionally under the control of nucleic
25 acid sequences directing their expression in cells of a particular lineage. Alternatively, nucleic acid binding polypeptide encoding constructs may be under the control of non-lineage-specific promoters, and/or inducibly regulated. Typically, DNA fragments on the order of 10 kilobases or less are used to construct a transgenic animal (Reeves, 1998, New. Anat., 253:19). A transgenic animal expressing one transgene can be crossed to a

second transgenic animal expressing second transgene such that their offspring will carry both transgenes.

Although the majority of previous studies have involved transgenic mice, other species of transgenic animal have also been produced, such as rabbits, sheep, pigs (Hammer et al., 1985, Nature 315:680-683; Kumar, et al., U.S. 05922854; Seebach, et al., U.S. Patent No. 6,030,833) and chickens (Salter et al., 1987, Virology 157:236-240). Transgenic animals are currently being developed to serve as bioreactors for the production of useful pharmaceutical compounds (Van Brunt, 1988, Bio/Technology 6:1149-1154; Wilmut, *et al.*, 1988, New Scientist (July 7 issue) pp. 56-59). Up-regulation of endogenous or exogenous genes expressing useful polypeptides, such as therapeutic polypeptides, by means of a heterologous nucleic acid binding polypeptide, may be used to produce such polypeptides in transgenic animals. Preferably, the polypeptides are secreted into an extractable fluid, such as blood or mammary fluid (milk), to enable easy isolation of the polypeptide.

Furthermore, the invention provides the use of polypeptide fusions comprising an integrase, such as a viral integrase, and a nucleic acid binding protein according to the invention to target nucleic acid sequences *in vivo* (Bushman, (1994) PNAS (USA) 91:9233-9237). In gene therapy applications, the method may be applied to the delivery of functional genes into defective genes, or the delivery of a heterologous nucleic acid in order to disrupt an endogenous gene. Alternatively, genes may be delivered to known, repetitive stretches of nucleic acid, such as centromeres, together with an activating sequence such as an LCR. This would represent a route to the safe and predictable incorporation of nucleic acid into the genome.

In conventional therapeutic applications, nucleic acid binding proteins according to this embodiment may be used to specifically eliminate cells having mutant vital proteins. For example, if a mutant ras gene is targeted, cells comprising this mutant gene will be destroyed because ras is essential to cellular survival. Alternatively, the action of transcription factors can be modulated, preferably reduced, by administering to the cell

agents which bind to the binding site specific for the transcription factor. For example, the activity of HIV tat may be reduced by binding proteins specific for HIV TAR.

Moreover, binding proteins according to the invention can be coupled to toxic molecules, such as nucleases, which are capable of causing irreversible nucleic acid damage and cell death. Such agents are capable of selectively destroying cells that comprise a mutation in their endogenous nucleic acid.

Nucleic acid binding proteins and derivatives thereof as set forth above may also be applied to the treatment of infections and the like in the form of organism-specific antibiotic or antiviral drugs. In such applications, the binding proteins can be coupled to a nuclease or other nuclear toxin and targeted specifically to the nucleic acids of microorganisms.

Transgenic animals comprising transgenes, optionally integrated within the genome, and expressing heterologous zinc finger and other nucleic acid binding polypeptides from transgenes, may be created by a variety of methods. Methods for producing transgenic animals are known in the art, and are described by Gordon, J. & Ruddle, F.H. *Science* 214: 1244-1246 (1981); Jaenisch, R. *Proc. Natl. Acad. Sci. USA* 73: 1260-1264 (1976); Gossler *et al.*, (1986) *Proc. Natl. Acad. Sci. USA* 83:9065-9069; Hogan *et al.*, *Manipulating the Mouse Embryo: A Laboratory Manual*, (1988); and US. Pat. Nos. 5,175,384; 5,434,340 and 5,591,669.

Pharmaceutical Preparations

25

The invention likewise relates to pharmaceutical preparations which contain the compounds according to the invention or pharmaceutically acceptable salts thereof as active ingredients, and to processes for their preparation.

30 The pharmaceutical preparations according to the invention which contain the compound according to the invention or pharmaceutically acceptable salts thereof are those for enteral, such as oral, furthermore rectal, and parenteral administration to (a) warm-

blooded animal(s), the pharmacological active ingredient being present on its own or together with a pharmaceutically acceptable carrier. The daily dose of the active ingredient depends on the age and the individual condition and also on the manner of administration.

5

The novel pharmaceutical preparations contain, for example, from about 10 % to about 80% (or any integral percentage therebetween), preferably from about 20 % to about 60 %, of the active ingredient. Pharmaceutical preparations according to the invention for enteral or parenteral administration are, for example, those in unit dose forms, such as
10 sugar-coated tablets, tablets, capsules or suppositories, and furthermore ampoules. These are prepared in a manner known per se, for example by means of conventional mixing, granulating, sugar-coating, dissolving or lyophilising processes. Thus, pharmaceutical preparations for oral use can be obtained by combining the active ingredient with solid carriers, if desired granulating a mixture obtained, and processing the mixture or granules,
15 if desired or necessary, after addition of suitable excipients to give tablets or sugar-coated tablet cores.

Suitable carriers are, in particular, fillers, such as sugars, for example lactose, sucrose, mannitol or sorbitol, cellulose preparations and/or calcium phosphates, for example
20 tricalcium phosphate or calcium hydrogen phosphate, furthermore binders, such as starch paste, using, for example, corn, wheat, rice or potato starch, gelatin, tragacanth, methylcellulose and/or polyvinylpyrrolidone, if desired, disintegrants, such as the abovementioned starches, furthermore carboxymethyl starch, crosslinked polyvinylpyrrolidone, agar, alginic acid or a salt thereof, such as sodium alginate;
25 auxiliaries are primarily glidants, flow-regulators and lubricants, for example silicic acid, talc, stearic acid or salts thereof, such as magnesium or calcium stearate, and/or polyethylene glycol. Sugar-coated tablet cores are provided with suitable coatings which, if desired, are resistant to gastric juice, using, inter alia, concentrated sugar solutions which, if desired, contain gum arabic, talc, polyvinylpyrrolidone, polyethylene glycol
30 and/or titanium dioxide, coating solutions in suitable organic solvents or solvent mixtures or, for the preparation of gastric juice-resistant coatings, solutions of suitable cellulose preparations, such as acetylcellulose phthalate or hydroxypropylmethylcellulose

phthalate. Colorants or pigments, for example to identify or to indicate different doses of active ingredient, may be added to the tablets or sugar-coated tablet coatings.

Other orally utilisable pharmaceutical preparations are hard gelatin capsules, and also soft closed capsules made of gelatin and a plasticiser, such as glycerol or sorbitol. The hard gelatin capsules may contain the active ingredient in the form of granules, for example in a mixture with fillers, such as lactose, binders, such as starches, and/or lubricants, such as talc or magnesium stearate, and, if desired, stabilisers. In soft capsules, the active ingredient is preferably dissolved or suspended in suitable liquids, such as fatty oils, paraffin oil or liquid polyethylene glycols, it also being possible to add stabilisers.

Suitable rectally utilisable pharmaceutical preparations are, for example, suppositories, which consist of a combination of the active ingredient with a suppository base. Suitable suppository bases are, for example, natural or synthetic triglycerides, paraffin hydrocarbons, polyethylene glycols or higher alkanols. Furthermore, gelatin rectal capsules which contain a combination of the active ingredient with a base substance may also be used. Suitable base substances are, for example, liquid triglycerides, polyethylene glycols or paraffin hydrocarbons.

Suitable preparations for parenteral administration are primarily aqueous solutions of an active ingredient in water-soluble form, for example a water-soluble salt, and furthermore suspensions of the active ingredient, such as appropriate oily injection suspensions, using suitable lipophilic solvents or vehicles, such as fatty oils, for example sesame oil, or synthetic fatty acid esters, for example ethyl oleate or triglycerides, or aqueous injection suspensions which contain viscosity-increasing substances, for example sodium carboxymethylcellulose, sorbitol and/or dextran, and, if necessary, also stabilisers.

The dose of the active ingredient depends on the warm-blooded animal species, the age and the individual condition and on the manner of administration. For example, an approximate daily dose of about 10 mg to about 250 mg is to be estimated in the case of oral administration for a patient weighing approximately 75 kg .

g. Transformation and Transfection

DNA can be stably incorporated into cells or can be transiently expressed using methods known in the art and described below. Stably transfected cells can be prepared by transfecting cells with an expression vector containing a selectable marker gene, and
5 growing the transfected cells under conditions selective for cells expressing the marker gene. To prepare transient transfectants, cells are transfected with a reporter gene to monitor transfection efficiency.

There are many well-known methods of introducing foreign nucleic acids into host cells, which include electroporation, calcium phosphate co-precipitation, particle
10 bombardment, microinjection, naked DNA, liposomes, lipofection, and viral infection etc (see, e.g. Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press, and Mountain, A. *Trends Biotechnol.* 18: 119-128 (2000) for a review). Any of the above methods can be used, as long as it is compatible with the host cell. Linear nucleic acid molecules have been found to be more
15 efficiently incorporated into mammalian genomes than circular plasmids. Additionally, nucleic acid molecules may be delivered to specific target tissues or to individual cells. Viral based gene transfer is often favoured for introducing nucleic acids into mammalian cells and specific target tissues, and several viral delivery approaches are in clinical trials for gene therapy applications. However, non-viral methods are attractive due to their
20 greater safety for the purpose of gene transfer to humans.

The preferred methods of particle bombardment use biolistics made from gold (or tungsten). Compared with other transfection procedures, particle bombardment requires a low amount of nucleic acid and a smaller number of cells, making the procedure generally more efficient (Heiser, W. C. *Anal. Biochem.* 217: 185-196 (1994); Klein, T. M. & Fitzpatrick-McElligott, S. *Curr. Opin. Biotechnol.* 4: 583-590 (1993)). The procedure
25 is particularly suited for organisms that are difficult to transfect, and for introducing DNA into organelles, such as mitochondria and chloroplasts. Although generally used for *ex vivo* applications, the procedure is also suitable for *in vivo* transfection of skin tissue. Suitable methods are known in the art and described, for instance, in US Patent Nos.

5,489,520 and 5,550,318. See also, Potrykus (1990) *Bio/Technol.* 8: 535-542; and Finnegan *et al.* (1994) *Bio/Technol.* 12: 883-888.

Microinjection is a common method of nucleic acid delivery to isolated cells (Palmiter, R. D. & Brinster, R. L. *Annu. Rev. Genet.* 20: 465-499 (1986); Wall, R. J. *et al.*, *J. Cell Biochem.* 49: 113-120 (1992); Chan, A. W. *et al.*, *Proc. Natl. Acad. Sci. USA* 95: 14028-14033 (1998)). DNA is generally injected into cells and the cells may then be re-introduced into animals. Procedures for such a technique are described in US Pat. Nos. 5,175,384 and 5,434,340, and improvements to the technique are described in WO 00/69257.

10 Efficient for gene transfer *in vivo* can be obtained following local injection of naked DNA. While expression of injected DNA in skin lasts for only a few days, injected DNA in mouse skeletal muscle has been shown to last for up to nine months (Wolff, J. A. *et al.*, *Hum. Mol. Genet.* 1: 363-369 (1992)). Naked DNA is particularly suited to gene therapy for preventive and therapeutic vaccines.

15 Cationic liposomes containing cholesterol are particularly suited for delivery of nucleic acids to humans as they are biodegradable and stable in the bloodstream. Liposomes can be injected intravenously, subcutaneously or inhaled as an aerosol. Stribling *et al.* (1992) *Proc. Natl. Acad. Sci. USA* 89:11,277-11,281. Liposomes can be targeted to certain cell types by incorporating ligands, receptors or antibodies
20 (immunolipids) into the lipid membrane (US. Pat. No. 4,957,773). On contacting target cells, entry of DNA from liposomes is via endocytosis and diffusion. Preparations of lipid formulations are commercially available and methods for their use are well documented (Bogdanenko, E. V. *et al.*, *Vopr. Med. Khim.* 46: 226-245 (2000); Natsume, A. *et al.*, *Gene Ther.* 6: 1626-1633 (1999)).

25 Uptake of DNA into animal cells can also be enhanced by using transfection agents. "Transfecting agent", as utilised herein, means a composition of matter added to the genetic material for enhancing the uptake of exogenous DNA segment (s) into a eukaryotic cell, preferably a mammalian cell, and more preferably a mammalian germ

cell. The enhancement is measured relative to the uptake in the absence of the transfecting agent. Examples of transfecting agents include adenovirus-transferrin-polylysine-DNA complexes. These complexes generally augment the uptake of DNA into the cell and reduce its breakdown during its passage through the cytoplasm to the nucleus of the cell. These complexes can be targeted to the male germ cells using specific ligands which are recognised by receptors on the cell surface of the germ cell, such as the c-kit ligand or modifications thereof. Other preferred transfecting agents include lipofectinTM, lipofectamineTM, DIMRIE C, Superfect, and Effectin (Qiagen), unifectin, maxifectin, DOTMA, DOGS (Transfectam; dioctadecylamidoglycylspermine), DOPE (1,2-dioleoyl-sn-glycero-3 phosphoethanolamine), DOTAP (1,2-dioleoyl-3-trimethylammonium propane), DDAB (dimethyl dioctadecylammonium bromide), DHDEAB (N, N-di-n-hexadecyl-N, N-dihydroxyethyl ammonium bromide), HDEAB (N-n-hexadecylN, N dihydroxyethylammonium bromide), polybrene, or poly (ethylenimine) (PEI). For example, Banerjee, R. *et al.*, Novel series of non-glycerol-based cationic transfection lipids for use in liposomal gene delivery, *J. Med. Chem.* 42 (21): 4292-99 (1999); Godbey, W. T. *et al.*, Improved packing of poly (ethylenimine)-DNA complexes increases transfection efficiency, *Gene Ther.* 6 (8): 1380-88 (1999); Kichler, A *et al.*, Influence of the DNA complexation medium on the transfection efficiency of lipospermine/DNA particles, *Gene Ther.* 5 (6): 855-60 (1998); Birchaa, J. C. *et al.*, Physico-chemical characterisation and transfection efficiency of lipid-based gene delivery complexes, *Int. J. Pharm.* 183 (2): 195-207 (1999). These non-viral agents have the advantage that they facilitate stable integration of xenogeneic DNA sequences into the vertebrate genome, without size restrictions commonly associated with virus-derived transfecting agents.

The most critical issues for applications such as gene therapy are the efficient delivery and appropriate expression of transgenes in host cells. For this purpose, viral systems are particularly well suited as viruses have evolved to efficiently cross the plasma membrane of eukaryotic cells and express their nucleic acids in host cells. Suitability of viral vectors is assessed primarily on their ability to carry foreign nucleic acids and deliver and express transgenes with high efficiency. Current applications utilise both RNA and DNA virus based systems, and 70% of gene therapy trials use viral vectors

derived from retroviruses, adenovirus, adeno-associated virus, herpesvirus and pox virus. See, for example, Flotte *et al.* (1995) *Gene Ther.* 2:357-362; Glorioso *et al.* (1995) *Ann. Rev. Microbiol.* 49:675-710; Smith (1995) *Ann. Rev. Microbiol.* 49:807-838; Prince (1998) *Pathology* 30:335-347; and Robbins *et al.* (1998) *Pharmacol. Ther.* 80:35-47.

5 Retroviruses represent the most prominent gene delivery system as they mediate high gene transfer and expression of therapeutic genes. Members of the DNA virus family such as adenovirus, adeno-associated virus or herpesvirus are popular due to their efficiency of gene delivery. Adenoviral vectors are particularly suited when transient transfection of nucleic acid is preferred. Retroviruses express particular envelope
10 proteins that bind to specific cell surface receptors on host cells, in order for the virus to enter the cell. Hence, the type of viral vector used should be determined by the tissue type to be targeted. See *e.g.*, Dornburg (1995) *Gene Ther.* 2:301-310; Gunzburg, *et al.* (1996) *J. Mol. Med.* 74:171-182; Vile *et al.* (1996) *Mol. Biotechnol.* 5:139-158; Miller (1997) "Development and Applications of Retroviral Vectors" Cold Spring Harbor Laboratory
15 Press, Cold Spring Harbor, New York; Karavanas *et al.* (1998) *Crit. Rev. Oncol. Hematol.* 28:7-30; Hu *et al.* (2000) *Pharmacol. Rev.* 52: 493-511; and Walther *et al.* (2000) *Drugs* 60: 249-271 for reviews.

Safety is a critical issue for viral based gene delivery because most viruses are either pathogens or have pathogenic potential. Generally, when a replication-competent
20 virus infects an animal cell it can express viral genes and release many new infectious viral particles in the host organism. Hence, it is very important that during transgene delivery the host animal does not receive a pathogenic virus with full replication potential. For this reason, viral-host cell systems have been developed for gene therapy treatments to prevent the creation of replication-competent viruses. In this method, viral
25 components are divided between a vector and a helper construct to limit the ability of the virus to replicate (Miller 1997). The viral vector contains the gene(s) of interest and cis-acting elements that allow gene expression and replication, but contain deletions of some or all of the viral proteins. Helper cells (or occasionally, helper virus) are engineered to express the viral proteins needed to propagate the viral vectors. These new viral particles
30 are able to infect target cells, reverse transcribe the vector RNA and integrate its DNA copy into the genome of the host, which can then be expressed. However, the vector can

not express the viral proteins required to create new infectious particles. Helper cell lines are known in the art (see Hu, W-S & Pathak, V. K. *Pharmacol. Rev.* 52: 493-511 (2000), for a review).

In general, retroviral vectors are able to package reasonably long stretches of foreign DNA (up to 10 kb). Oncoviruses are a type of retrovirus, which only infect rapidly dividing cells. For this reason they are especially attractive for cancer therapy. Murine leukaemia virus (MLV)-based vectors are the most commonly used of this class. Spleen necrosis virus (SNV), Rous sarcoma virus and avian leukosis virus are other types. Lentiviral vectors are retroviral vectors that can be propagated to produce high viral titres and are able to infect non-dividing cells. They are more complex than oncoviruses and require regulation of their replication cycle. Lentiviral vectors which may be used include human immunodeficiency virus (HIV-1 and -2) and simian immunodeficiency virus (SIV) based systems. HIV infects cells of the immune system, most importantly CD4⁺ T-lymphocytes, and so may be useful for targeted gene therapy of this cell type. Another type of retrovirus is the spumavirus. Spumaviruses are attractive because of their apparent lack of toxicity. Linial (1999) *J. Virol.* 73:1747-1755.

Adenoviral vectors have high transduction efficiency and are able to transfect a number of different cell types, including non-dividing cells. They have a high capacity for foreign DNA and can carry up to 30 kb of non-viral DNA (for a review see, Kochanek, S. *Hum. Gene Ther.* 10: 2451-2459 (1999)). Recombinant adenoviral (rAd) vectors are becoming one of the most powerful gene delivery systems available and have been used to deliver DNA to post-mitotic neurons of the central nervous system (CNS) (Geddes, B. J. *et al.*, *Front. Neuroendocrinol.* 20: 296-316 (1999), and are used to treat diseases such as colon cancer (Alvarez *et al.*, *Hum. Gene Ther.* 5: 597-613 (1997)). Adeno-associated virus (AAV) vectors and recombinant AAV (rAAV) vectors are proving themselves to be safe and efficacious for the long-term expression of proteins to correct genetic disease. Snyder, R. O. J. (*Gene. Med.* 1: 166-175 (1999)) provides a review of gene delivery approaches using such vectors. Construction of such vectors is described in, for example, Samulski *et al.*, *J. Virol.* 63: 3822-3828 (1989), and US. Pat. No. 5,173,414.

Many gene therapy trials have been conducted and are underway (over 3,500 people have been treated with gene therapy systems), and several reviews can be studied for details of the protocols and results (Hwu & Rosenberg, *Ann N Y Acad Sci.* 1994 May 31;716:188-97; Blaese, *Hosp Pract (Off Ed).* 1995 Nov 15;30(11):33-40; Blaese, *Hosp Pract (Off Ed).* 1995 Dec 15;30(12):37-45; Breau & Clayman, *Curr Opin Oncol.* 1996 May; 8(3):227-31; Dunbar *Annu Rev Med.* 1996;47:11-206; Lotze *Cancer J Sci Am.* 1996 Mar;2(2):63). The first gene therapy trial was carried out by Blaese *et al.*, (1995), to correct a genetic disorder known as adenosine deaminase (ADA) deficiency, which leads to severe immunodeficiency. Several cancer gene therapy strategies are being developed, which involve eliminating cancer cells by suicide therapy (Oldfield *et al.*, *Hum Gene Ther.* 1993 Feb;4(1):39-69), modification of cancer cells to promote immune responses (Lotze *et al.*, *Hum Gene Ther.* 1994 Jan;5(1):41-55), and reversion by delivery of a tumor suppressor gene (Roth *et al.*, *Hum Gene Ther.* 1996 May 1;7(7):861-74). Another successful gene therapy trial has been conducted to combat graft-versus-host disease, which can result following transplant procedures such as bone marrow transplants (Bonini *et al.*, *Science.* 1997 Jun 13;276(5319):1719-24). This procedure was carried out using an HSV-based vector. Several gene therapy treatments are under investigation for the treatment of HIV-1 infection. Most treatments involve modification of lymphocytes, *ex vivo*, to suppress the expression of viral genes, by means of ribozymes, antisense RNA, mutant trans-dominant regulatory proteins and modification to elicit a host immune response (Nabel *et al.*, *Cardiovasc Res.* 1994 Apr;28(4):445-55; Galpin *et al.*, *Hum Gene Ther.* 1994 Aug;5(8):997-1017; Morgan RA, Walker R. *Hum Gene Ther* 1996 Jun 20;7(10):1281-306 Gene therapy for AIDS using retroviral mediated gene transfer to deliver HIV-1 antisense TAR and transdominant Rev protein genes to syngeneic lymphocytes in HIV-1 infected identical twins; Wong-Staal *et al.*, *Hum Gene Ther.* 1998 Nov 1;9(16):2407-25). Vectors currently in use for gene therapy treatments and animal tests include those derived from Moloney murine leukemia virus, such as MFG and derivative thereof, and the MSCV retroviral expression system (Clontech, Palo Alto, California). Many other vectors are also commercially available.

Viral vectors are especially important in applications when a specific tissue type is to be targeted, such as for gene therapy applications. There are two available methods for

targeting genes to specific cell or tissue type. One strategy is designed to control expression of the required gene using a tissue specific promoter (discussed above), and another strategy is to control viral entry into cells. Viruses tend to enter specific cell types according to the envelope proteins that they express. However, by engineering the envelope proteins to express specific proteins as fusions, such as erythropoietin, insulin-like growth factor I and single chain variable fragment antibodies, viral vectors can be targeted to specific cell-types (Kasahara *et al.*, Science. 1994 Nov 25;266(5189):1373-6; Somia *et al.*, Proc Natl Acad Sci U S A. 1995 Aug 1;92(16):7570-4; Jiang *et al.*, J Virol. 1998 Dec;72(12):10148-56; Chadwick *et al.*, J Mol Biol. 1999 Jan 15;285(2):485-94).

In one example of tissue specific targeting in transgenic mice, a novel transgene delivery system has been developed in which the target tissue type expresses an avian viral receptor (TVA), under the control of a tissue specific promoter. Transgenic mice expressing the TVA receptor are then infected with avian leukosis virus, carrying the transgene(s) of interest (Fisher, G. H. *et al.*, *Oncogene* 18: 5253-5260 (1999)).

h. Construction of Zinc Finger libraries

Zinc finger libraries may be constructed from naturally-occurring human zinc finger modules. Thus, the invention provides libraries of zinc finger modules. Module libraries according to the invention may be assembled combinatorially into zinc finger polypeptides. The combinatorial assembly may be carried out biologically, using random assembly and selection technologies, or in a directed manner under computer control, assembling desired modules to produce zinc fingers having defined or random specificity. In accordance with the invention, libraries may be constructed entirely from natural zinc finger polypeptide modules from which zinc finger polypeptides having any desired specificity may be isolated. The invention, in its most preferred aspect, does not require the engineering of the specificity of any zinc finger module in order to produce a zinc finger polypeptide having specificity for any desired nucleic acid sequence.

Selection of appropriate zinc finger modules for assembly into libraries of composite binding polypeptides having a predetermined binding specificity can be

accomplished by applying the rules for zinc finger binding specificity set forth herein. In the case of zinc finger assembly under computer control, a rule table may be used to select zinc fingers for binding to the target site. Figure 1 shows a flowchart depicting part of the logic used in the selection of zinc fingers from a natural library in accordance with the invention. The logic set forth in Figure 1 may be supplemented, for example using Rules relating to zinc finger overlap. Functional testing of zinc fingers for binding to the desired binding site may be implemented in an automated fashion and integrated with the zinc finger design system.

The invention thus provides libraries of zinc finger modules. In one embodiment, the modules are human zinc finger modules. Preferably, the modules are DNA-binding zinc finger modules.

In a preferred aspect the invention provides a library of DNA-binding human zinc finger modules as set out in Example 1 below. Moreover, the invention provides a library of human zinc finger modules as set forth in Example 2 below. Sub-libraries can be prepared from either of the libraries of the invention.

The invention furthermore encompasses libraries in which zinc finger modules as set forth in Examples 1 or 2 herein are combined with other zinc finger modules to provide further libraries that may be used to generate zinc finger polypeptides.

In a still further aspect, the invention relates to libraries derived from animals other than humans, for use in said organisms in order to derive some or all of the same advantages as may be obtained with human zinc fingers for use in humans. Example 3 sets forth databases of zinc fingers from mouse, chicken and plants. Sequences of zinc fingers can be identified in other organisms by the same means, *i.e.* by analysis of sequence information and identification of zinc fingers in accordance with the guidance given herein.

EXAMPLES**Example 1. List of selected human DNA-binding zinc fingers.**

These fingers have been selected from the human genome on the basis of a prediction that they have a DNA-binding potential. This prediction is based on coded contacts (WO 96/06166, WO 98/53057, WO 98/53058; WO 98/53059 and WO 98/53060); accordingly, for each peptide unit, a 3-nucleotide DNA target subsite is shown, as the preferred sequence to which the zinc finger binds. Hence, by constructing 2- or 3-finger libraries from these 200 or so units, in the manner described in the Examples *infra*, there exists the potential to screen a large variety of novel DNA target sites. Note that the predicted DNA target subsites listed below are merely intended to be a guide to the DNA-binding potential. It is anticipated that, in practice, an even wider range of DNA sequences can be targeted using a library engineered from this database, through the exertion of a positive selection pressure in the library screening system.

15

The fingers listed below are in a format that can be linked with classical wild-type canonical "TGEKP" (SEQ ID NO:3) linkers (i.e. ...TGEKP – zinc finger peptide sequence – TGEKP – zinc finger peptide sequence – TGEKP - etc...). For each peptide sequence, an oligonucleotide is designed to encode the peptide sequence; the oligonucleotide can then be linked into a library selection system, as described in the Examples *infra*.

20

Database of predicted human DNA-binding zinc fingers

25 227 finger units

Zinc finger	DNA site	SEQ ID NO	Peptide sequence
ZIF268 F1	GCG	31	YACPVESCDRRFSRSDDELTRHIRIH
ZIF268 F2	TGG	32	FQCRICMRNFSRSDHLSTHIRTH
ZIF268 F3	GCG	33	FACDICGRKFARSDEKRHTKIH
Kr-like13	NGT	34	HKCHYAGCEKVYGKSSHLKAHLRTH
MAZ F1	AGG	35	YQCPVCQQRFKRKDRMSYHVRSH

MAZ F2	TGG	36	YNCSHCGKSFSRPDHLNSHVRQVH
MAZ F3	NGT	37	FKCEKCEAAAFATKDRLRAHTVRH
TIEG2 (SP1) F3	GGG	38	FVCPVCDRRFMRS DHLT KHARRH
SP1 F1	GGG	39	HKCHYAGCEKVY GKSSHLKAHLRTH
SP1 F2	GCG	40	FACSWQDCNKKFARSDELARHYRTH
SP1 F3	GGG	41	FSCPTICEKR FMRS DHLT KHARRH
WT1 F1	TGT	42	FMCAYPGCNKRYFKLSHLQMHSRKH
WT1 F2	GAG	43	YQCDFKDCERRFSRSDQLKRHQRRH
WT1 F3	TGG	44	FQCKTCQRKFSRSDHLKTHTRTH
WT1 F4	GCG	45	FSCRWPSCQKKFARSDELVRHHNMH
TY Y1	TAT	46	FQCTFEGCGKRFSLDNFNLRTHVRIH
TY Y1	NAA	47	YVCPFDGCNKKFAQSTNLKSHILTH
TF3A	GGG	48	FVCDYEGCGKAFIRDYHLSRHILTH
TF3A	GGC	49	FKCTQEGCGKHFASPSKLKRHAKAH
MAZ	GGC	50	HACEMCGKAFRDVYHLNRHKLSH
GLI1	GCA	51	YMCEHEGCSKAFSNASDRAKHQNRTH
ZIC3	GCA	52	FKCEFE GCDRRFANS SDRKKHMHVH
SP4	NGG	53	HICHIEGCGKVY GKTSHLRAHLRWH
SP2	NTG	54	HVCHIPDCGKTFRKTSLLRAHVRLH
BTE1	NGG	55	HKCPYSGCGKVY GKSSHLKAHYRVH
GLI2	TAG	56	HKCTFEGCSKAYSRLLENLKTHLRSH
Q14872	TAT	57	YQCTFEGCPRTYSTAGNLRTHQKTH
Q14872	TGC	58	FRCDHDGCGKAFAASHHLKTHVVRTH
ZIC3	TAG	59	FPCPFPGCGKIFARSEN LK I HKRTH
Z143	CTT	60	FKCPFE GCGRSFTTSNIRKVHVVRTH
Z143	CGT	61	FRCEYD GCGKLYTTAHLKVHERSH
O00153	AAT	62	FMCHESGCGKQFTTAGNLKNHRRRIH
Z143	AAC	63	YYCTEPGCGGRAFASATNYKNHVRIH
Q14872	TCT	64	FVCNQEGCGKAFLTSHSLRIHVVRH
O00153	TGT	65	FICPAEGCGKS FYVLQRLKVHMRTH
Q14872	GCT	66	FNCESEGC SKYFTT LSDLRKHIRTH
Z143	GCT	67	YRCSEDNCTKSFKTSGDLQKHIRTH
BTE1	GCG	68	FPCTWPDCLKKFSRSDDELTRHYRTH
O15391	TAA	69	FVCPFDVCNRKFAQSTNLKTHILTH
Z143	GNC	70	YVCTVPGCDKRFT EYSSLYKHHVVH
O43591	GGT	71	HVCEHCNAAFRTNYHLQRHVFIH
BCL6	TAG	72	YRCNICGAQFNRPANLKTHTRIH
O75626	TAC	73	HECQVCHKRFSSTS NLKTHLRLH
O75626	YAA	74	YECNVCAKTFGQLSNLKVHLRVH
BCL6	NGA	75	YKCETCGARFVQVAHLRAHVLIH

O75626	GGA	76	FKCQTCNKGFTQLAHLQKHLYLVH
ZN45	N (N/T) A	77	YRCDVCGKRFRQRSYLQAHQRVH
BCL6	YTY	78	YPCEICGTRFRHLQTLKSHLRH
GFI1	GCA	79	YPCQYCGKRFHQKSDMKKHTFIH
Z263	GAN	80	YQCNICGKCFSCNSNLHRHQRT
ZN75	TAY	81	YRCSWGKSFHNTNLHTHQRIH
Z186	TTT (YYY)	82	YKCIECGKTFTVNQLLTLHHRT
Z136	TTT (YYY)	83	FKCKQCGKAFSCSPTLRIHERTH
Z136	TGA	84	YKCKVCGKAFDYPFRFRTHERSH
Z136	TTT (YYY)	85	YKCKVCGKPFHSLSSFQVHERIH
Z177	TTA	86	YECKECGKAFRNSSCLRVRHVRTH
Z136	TNN	87	FECKRCGKAFRSSSSFRLHERTH
O60765	A/T-YT	88	YRCNECGKGFTSISRLNRHRIIH
ZN42	TYT	89	YHCGECGLGFTQVSRLTEHQRIH
ZN42	CGG	90	FVCGDCGQGFVRSARLEEHRRVH
O14913	TCG	91	YKCEKCGKGFFRSSLQHHQKIH
O14913	C-G/T-G	92	YKCEECGKGFSRSSKLQEHQTIH
ZN45	YYC	93	YKCEECGKGFCRASNLLDHQRGH
ZN45	AAA	94	YKCEECGKGFSQASNLLAHQRGH
ZN45	NAG	95	YQCEECGKGFCRASNFLAHRGVH
Z239	YYG	96	YKCEQCGKGFTRSSSLLIHQAVH
O94892	YNY	97	YRCSECGKGFI VNSGLMLHQRT
ZN45	AAY	98	YQCAECGKGFSVGSQQLQAHQRC
ZN45	NGY	99	YKCEECGKGFSVGSHLQAHQISH
ZN45	YCG	100	YQCDACGKGFSRSSDFNIHFRVH
ZN45	CCG	101	YKCGTCGKGFSRSSDLNVHCRIH
ZN45	TGA	102	YKCNACGKSFSYSSHLNIHCRIH
Z239	TCA	103	YQCYECGKGFSQSSDLRIHLRVH
Z239	YAA	104	YKCGECGKGFSQSSNLHIHRCIH
Z239	YGA	105	YKCDKCGKGFSQSSKLHIHQRVH
Z239	CGA	106	YHCGKCGKGFSQSSKLLIHQRVH
O60765	AYA	107	FKCSECGRAFSQSASLIQHERIH
O60792	GYG	108	YECKECGKA FIRSSSLAKHERIH
ZN07	ATA	109	YPCKECGKA FSQSSTLAQHQRMH
O43296	AYY	110	YKCEECGKA FSRSSSLTQHQRMH
Z134	ATG	111	YKCEECGKA FS RKDTLVQHQRH
Z134	ATG	112	YECSECGKA FS RKATLVQHQRH
ZN84	AYC	113	YECSECGKA FSEKLSLTNHQRH
Z191	AYG	114	YGCVECGKA FSRS SILVQHQRVH
ZN24	ACG	115	YGCVECGKA FSRS SILVQHQRVH

O43338	GTA	116	YVCGQCGKSFSQRATLIKHHRVH
O43339	GTA	117	YECSQCGKSFSQKATLVKHQRVH
O43338	AYA	118	YDCGQCGKSFIQKSSLIQHQQVH
O43339	ANA	119	YECGQCGKSFSQKSGLIQHQQVH
O43338	CAA	120	YECGECGKSFSQSSNLIHCRH
Q13398	AAA	121	YECGECGKSFSQRSNLMQHRRVH
Z135	CYA	122	YECGECGKAFSQSTLLTEHRRH
Q13398	ACA	123	YECSECGKSFSQSSSLIQHRRVH
O14709	AAA	124	YKCNECGKAFSQSAYLLNHQRIH
O14709	CAA	125	YKCNECGKVFSQNAYLIDHQRH
O14709	CAA	126	YKCTECGKAFTQSAYLFDHQRH
O14709	CAA	127	YKCDECGKTFAQTTYLIDHQRH
O60792	AAA	128	YNCNECRKTFSQSTYLIQHQRH
O15535	ANA	129	YHCKECGKVFSQSAGLIQHQRH
Q15776 (a)	TNA	130	YHCKECGKAFSQNTGLILHQRH
Q15776 (b)	TNA	131	YQCNQCGKAFSQSAGLILHQRH
Q15776	CNA	132	YKCNECGRAFSQKSGLIEHQRH
ZN84	AAC	133	YGCNECGRAFSEKSNLINHQRH
Z191	ANA	134	YKCLECGKAFSQNSGLINHQRH
ZN24	ANA	135	YKCLECGKAFSQNSGLINHQRH
O60765	AYA	136	YRCEECGISFGQSSALIQHRRH
ZN07	YYA	137	YRCEECGKAFGQSSSLIHHQRH
O43340	ACA	138	YECDECGKSYSQSSALLQHRRVH
Z135	CYY	139	YKCQECGKAFSHSSALIEHHRH
O43340	AYA	140	YDCSECGKSFRQVSVLIQHQRVH
O43340	AYA	141	YVCSECGKSFGQKSVLIQHQRVH
Q13398	AYT	142	YQCSQCGKSFGCKSVLIQHQRVH
O15535	GNA	143	HKCDECGKSFTQSSGLIRHQRH
Q15776	GNA	144	HKCDECGKSFAQSSGLVRHWRH
O75802	ANG	145	HKCEECGKAFSRSSGLIQHQRH
Z189	ANG	146	HKCEECGKAFSRSSGLIQHQRH
O75802	ANG	147	HKCDECGKAFSRNSGLIQHQRH
Q13398	YYG	148	HECNECGKSFSRSSSLIHHRRH
Z195	YAA	149	YKCDECGKNFTQSSNLIVHKRH
O43309	CYA	150	YKCDKCGKAFTQRSVLTEHQRH
Z195	CGA	151	YKCDECGKAYTQSSHLSEHRRH
ZN45	YYA	152	YKCERCGKAFSQFSSLQVHQRVH
O60893	YYN	153	YECEDCGKTFIGSSALVIHQRVH
ZN07	TAT	154	YECLQCGKAFSMSTQLTIHQRVH
O60893	CYA	155	YECDDCGKTFSQSCSLLEHHKH

Q15776	NGG	156	YECDECGKTFRRSSHLIGHQRSH
ZN84	YGG	157	YECGECGKAFSRKSHLISHWRTH
Z177	YGA	158	YECDHCGKSFSQSSHLNVHKRTH
O43296	AYG	159	YECMECGKA FNRSYLTQHQRTH
O43296	GNG	160	YECVECGKAFTRM SGLTRHKRIH
O43340	AGG	161	YECRECGKS FTRKNHLIQHKT VH
Z134	AAG	162	YECSECGKTF SRKDNLTQH KRIH
O43338	CGA	163	YECSECGKSFSQTSHLNDHRRIH
O75467	AGA	164	YECAQCGKA FSQTSHLTQHQRTH
Z135	AGA	165	YECSECGKA FRQSIHLTQH LRIH
Z135	AGA	166	YEC HDCGKSFRQSTH LTQHRRIH
Z205	AGG	167	YACTDCGKR FGRSSHLIQHQI IH
O43296	AGG	168	YECTECGKT FIKSTHLLQH HMIH
O75290	AAG	169	YECKECGKYFSRSANLIQH QSIH
O75290	AGG	170	YECKECGKGFNRGAHLIQH QKIH
O75290	AGG	171	YECKECGKGFNRGAHLIQH QKIH
O60792	CGA	172	YTCNECGKA FSQRGHFMEHQ KIH
O75123	CGA	173	YTCDQCGKGFGQSSHLMEHQRIH
O43337	GYA	174	YECNACGKA FSQSSTLIRHYLIH
O75802	GYG	175	YECNYCGKTFSVSSTLIRHQRIH
Z165	GGY	176	YECSECGKTF RVSSHLIRHFRIH
Z124	CYY	177	YVCNNCGKGFRCS SSLRDHERTH
Z135	AYY	178	YGCNECGKTFSHSSSL SQHERTH
O15361	GAY	179	YDCNHCGKS FNHKTNLNKHERIH
O75123	AAA	180	YVCNECGKRFSQTSNFTQHQRTH
Q13398	AAV	181	YVCGECGKSFSHSSNLKNHQRVH
ZN35	YYA	182	YTCNECGKA FRQRSSLTVHQ RTH
Z157	YYC	183	YECTECGKT FSEKATLTIHQ RTH
O43338	GGY	184	YECDECGKA FGSKSTLVRHQ RTH
ZN84	TYC	185	YECSECGKA FGEKSSLATHQ RTH
ZN07	GAA	186	YGCRECGKA FSQQSQLVRHQ RTH
ZN84	YAA	187	YNCSQCGKA FSQKSQLTSHQ RTH
Z186	YGY	188	YACDHCEKA FSHKSKLTVHQ RTH
O43338	GGC	189	YVCGECGKA FMFKSKLVRHQ RTH
OZF	YYA	190	YECNVCGKA FSQSSSLTVHV RSH
O95779	YYY	191	YKCECGKA FNHC SLLTIHERTH
Z135	GGY	192	YACRDCGKA FTHSSSLTKHQ RTH
ZN80	GYA	193	YECKECGKGFYYSYSLTRHTRSH
Z177	GYC	194	YECSDCGKA FIDQSSLKKHTRSH
Z177	GGY	195	YDCKECGKA FTVPSSLQKHVRTH

O43337	ACT	196	YDCMACGKAFCRSSELIQHQRH
Q14585	AGY	197	YECKECEKAFRSGSKLIQHQRMH
Q14585	AAY	198	YECIDCGKAFGSGSNLTQHRRH
Q14585	GYG	199	YECKACGMAFSSGSALTRHQRH
Q14585	AYY	200	YECKECGKAFFYSGSSLTQHQRH
Q14585	AAY	201	YECKECGKAFGSGANLAYHQRH
Q14585	GAY	202	FECCKECGKAFGSGSNLTHHQRH
Q14585	ACY	203	YVCKECGKAFFNSGSDLTQHQRH
O60792	ACY	204	YQCHECGKTF SYGSSLIQHRKH
O60893	GNA	205	HYCHECGKSFAQSSGLTKHRRH
Z165	GCC	206	YECNECGKSFAESSDLTRHRRH
O60893	GAY	207	YECCECGKVFSHSSNLIKHQRT
Q15776	NGY	208	YECNECGKAFFSHSSHLIGHQRH
Z135	GYG	209	YQCGECGKAFFSHSSSLTKHQRH
Z165	GGY	210	HQCNECGKAFFRHSSKLARHQRH
Z135	TYG	211	YECHECLKGFRNSSALTKHQRH
O43361	YGC	212	YECNECGKFFLDSYKLVIHQH
O43361	YGC	213	YECSECGKFFRDSYKLIHQH
Z140	YYG	214	YGCHECGKTFGRRFSLVLHQRT
O60792	AAA	215	YECNECGKAFFSQHSNLTQHOKH
Z135	ANA	216	YKCTQCGRTFNQIAPLIHQH
Z135	ANA	217	YECNQCGRAFSQLAPLIHQH
Z135	ANA	218	YECHECGKAFTQITPLIHQH
O43309	AGA	219	YKCNECGKAFFGRWSALNQHQLH
ZN83	AGA	220	YKCNECGKVFNHMSHLAQHRRH
ZN83	AGY	221	YRCNVCGKVFFHHISHLAQHQRH
ZN83	AGA	222	YKCNECGKVFNQISHLAQHQRH
O14709	CAY	223	FECSECGRAFSSNRNLIHKKRH
ZN74	GYA	224	YKCSECGRAFSQNHCLIKHQKH
Q13398	ANA	225	YECSECGKSFSQNFSLIYHQH
O75123	GYA	226	FECCKECGKGFSQSSLLIRHQRH
Z132 (a)	GGA	227	FECSECGRDFSQSSHLRLHQKH
Z132	GYA	228	YECNECGKFFSQNSILIKHQKH
Z132 (b)	GGA	229	YECDECGKAFFSNRSHLIRHEKH
Z132	GGN	230	YECSECGRAFSSNSHLVRHQH
Z132	AAA	231	YECSECGRAFNNNSNLAQHOKH
Z134	ATY	232	YKCSDCGKVFRHKSTLVQHESI
O75290	AAT	233	YECKECGKAFFRLYLQLSQHOKH
Z157	AYC	234	YECGECGKNFRAKKSLNQHQRH
Z157	TTT	235	YECGECGKFFRMKMTLNNHQRT

ZN07	AAT	236	YECAECGKVFRLCSQLNQHQRIH
Z157	AYT	237	YECSECGKIFSMKKSLCQHRRTTH
O43361	GGY	238	YECNKCGKFFMYNSKLIRHQKVH
O43361	GTY	239	YKCSKCGKFFRYRCTLSRHQKVH
Z157	CGY	240	YECNECGNAFYVKARLIEHQRMH
Z157	CGY	241	YECSECGNAFYVKVRLIEHQRIH
O75123	AGG	242	FECNECGKAFIRSSKLIQHQRIH
ZN07	AGT	243	FKCTECGKAFRLSSKLIQHQRIH
O75123	GYT	244	YECNECGKAFFLSSYLIRHQKIH
O75802	AAT	245	HKCGECGKAFRLSTYLIQHQKIH
Z174	GCG RNA	246	YKCDDCGKSFTWNSELKRHKRVH
Z202	GCG RNA	247	YRCDDCGKHFRWTSIDLVRHQRTTH
O43345	GTG RNA	248	YKCEECGKAYKWPSTLSYHKKIH
O43345	CA? RNA	249	YKCEECGKAFFNWSSNLMEHKKIH
O75346	TAA	250	YRCEECGKAFFNQSANLTTHKRIH
ZN43	TAA	251	YKCEECGKAFTQSSNLTTTHKKIH
ZN85	GGA	252	YKCEECGKAFFNQSSKLTCHKKIH
ZN85	GAA	253	YTCEECGKAFFNQSSNLTKHKRIH
Q02313	GAA	254	YKCEECGKAFFNQLSNLTRHKVIH
Q02313	CAA	255	YKCEECGKAFFKQFSNLTDHKKIH
Z141	GTG	256	YKCEECGKAFFNRSTTLTKHKRIH
ZN91	TTG	257	YKCEECGKAFFSRSSTLTKHKTIH

Example 2: List of all human C₂H₂ zinc fingers

This list represents an even more comprehensive database of human zinc fingers, including those with non-DNA-binding activities such as those mediating protein-protein interactions and those involved in RNA binding. By including fingers from this database into a natural finger selection system as disclosed herein, many new zinc finger proteins having unique target specificities can be obtained. All of these peptides would necessarily possess properties required for potential therapeutic agents, such as non-immunogenicity.

The fingers listed below are in a format that can be linked with classical canonical "TGEKP" linkers (i.e. ...TGEKP – zinc finger peptide sequence – TGEKP – zinc finger peptide sequence – TGEKP - etc...). For each peptide sequence, an oligonucleotide is designed to encode the peptide sequence; the oligonucleotide can then be linked into a library selection system, as described in the Examples *infra*.

Human zinc finger database

968 finger units

Name	SEQ ID NO	Peptide sequence
Q92981_HUMAN	258	HQCAHCEKTFNRKDHLKNHFQTH
O76019_HUMAN	259	HQCAHCEKTFNRKDHLKNHLQTH
ZFY_HUMAN	260	HRCEYCKKGFRRPSEKNQHIMRH
ZFX_HUMAN	261	HRCEYCKKGFRRPSEKNQHIMRH
ZFX_BOVIN	262	HRCEYCKKGFRRPSEKNQHIMRH
Q15558_HUMAN	263	HRCEYCKKGFRRPSEKNQHIMRH
ZFX_HUMAN	264	HKCDMCDKGFHRPSELKKHVA AH
ZFY_HUMAN	265	HKCEMCEKGFHRPSELKKHVAVH
Q15558_HUMAN	266	HKCEMCEKGFHRPSELKKHVAVH
Z161_HUMAN	267	YTCSVCGKGFSRDPDHL SCHVKHVH
MAZ_HUMAN	268	YNCSHCGKSFSRDPDHLNSHVRQVH
O43829_HUMAN	269	YSCEVCGKSFI RAPDLKKHERVH
O00403_HUMAN	270	YSCEVCGKSFI RAPDLKKHERVH
Z151_HUMAN	271	HKCPHCDKKFNQVGNLKAHLKIH
Q92618_HUMAN	272	YKCPYCDHRASQKGNLKIHIRSH
ZFX_HUMAN	273	FRCKRCRKGFRQQSELKKHMKTH
Q14526_HUMAN	274	YPCTICGKKFTQ RGTMTMRHMRSH
HKR3_HUMAN	275	FECTECGYKFTRQAHLRRHMEIH
Q14526_HUMAN	276	YACDACGMRFT RQYRLTEHMRIH
O75626_HUMAN	277	YECNVCAKTFGQLSNLKVHLRVH
CTCF_HUMAN	278	HKCPDCDMAFVTSGELVRHRRYKH
O75701_HUMAN	279	YSCPDCSLRFAYTSL LAIHRRIH

O75701_HUMAN	280	YACSDCKSRFTYPYLLAIHQQRKH
O43167_HUMAN	281	YACKDCGKVFKYNHFLAIHQQRSH
O75850_HUMAN	282	CACPDCGRSFTQRAHMLLHQQRSH
O75850_HUMAN	283	YACPDCGRGFSGHQLARHPRVH
ZN42_HUMAN	284	FVCGDCGQGFVRSARLEEHRVH
O75467_HUMAN	285	FRCVDCGKAFAKGAVLLSHRRIH
O15015_HUMAN	286	YKCSECGRAYRHRGSLVNHRHSH
O75701_HUMAN	287	YPCPDCGRRFRQRGSLAIHRRAH
Q92951_HUMAN	288	YECAICQRSFRNQSNLAVHRRVH
BCL6_HUMAN	289	YKCDRCQASFRYKGNLASHKTVH
ZN42_HUMAN	290	YACQDCGRRFHQSTKLIQHQRVH
O75701_HUMAN	291	YPCPDCGRRFTYSSLLLSHRRIH
O75701_HUMAN	292	HVCTDCGRRFTYPSLLVSHRRMH
O75701_HUMAN	293	HSCPDCGRNFSYPSSLASHQRVH
ZN42_HUMAN	294	YACVECGERFGRRSVLLQHRRVH
O43298_HUMAN	295	YCGVCGKFKMKHHLVGHMKIH
O15209_HUMAN	296	YDCPVCNKKFKMKHHLTEHMKTH
O43829_HUMAN	297	YACHMCDKAFKHKSHLKDHERRH
O00403_HUMAN	298	YACHMCDKAFKHKSHLKDHERRH
O60315_HUMAN	299	HQCQICKKAFKHKHHLIEHSRLH
Q12924_HUMAN	300	HECGICKKAFKHKHHLIEHMRLH
NIL2_HUMAN	301	HECGICKKAFKHKHHLIEHMRLH
Q12924_HUMAN	302	FKCTECGKAFFKYKHHLKEHLRIH
O60315_HUMAN	303	FKCTECGKAFFKYKHHLKEHLRIH
NIL2_HUMAN	304	FKCTECGKAFFKYKHHLKEHLRIH
O95780_HUMAN	305	YKCEECGKAFFKRCSHLNEHKRVQ
O95779_HUMAN	306	YKCEECGKAFFKRCSHLNEHKRVQ
O43296_HUMAN	307	FKCSECGKVFNKKHLLAGHEKIH
O14709_HUMAN	308	YKCKECGKGFIYRHSGLIIHLRRH
O14709_HUMAN	309	HKCKECGKGFIQRSSLLMHLRNH
ZN80_HUMAN	310	CKCVECGKVFNRRSHLLCYRQIH
O43337_HUMAN	311	YKCIECGKAFFKRRSHLLQHQRVH
O60765_HUMAN	312	YICKECGKAFTLSTSLYKHLRTH
Z136_HUMAN	313	FECKRCGKAFFRSSSSFRLHERTH
Z136_HUMAN	314	FVCKQCGKAFFRSASTFQIHERTH
Z136_HUMAN	315	YVCKHCGKAFFVSSTSIRIHERTH
Z136_HUMAN	316	FKCKQCGKAFFSCSPTLRIHERTH
Z124_HUMAN	317	YVCNCGKGFRCSSSLRDHERTH
Z177_HUMAN	318	YECKECGKAFFRNSSCLRVRVTH
Z124_HUMAN	319	YECKHCGKAFFRYSNCLHYHERTH
O95780_HUMAN	320	YKCKECGKAFFNHCSLLTIHERTH
O95779_HUMAN	321	YKCKECGKAFFNHCSLLTIHERTH
Z124_HUMAN	322	YPCKQCGKAFFRYASSLQKHEKTH
Z136_HUMAN	323	YECKQCGKAFFSYLNSFRTHEMI
Z136_HUMAN	324	YECKQCGKAFFSYLPSLRLHERIH
O15060_HUMAN	325	YSCKVCGKRFATSEFNHYHRRIH
Z136_HUMAN	326	YKCKVCGKPFHSLSPFRIHERTH
Z136_HUMAN	327	YKCKVCGKPFHSLSSFQVHERIH

Z136_HUMAN	328	YKCKVCGKAFDYPSRFRTHERSH
ZN35_HUMAN	329	YVCNECGKAFTCSSYLLIHQRIH
O15322_HUMAN	330	YNCKECGKSFRWSSYLLIHQRIH
Q92951_HUMAN	331	YRCDQCGKAFSQKGS LIVHIRVH
Q92951_HUMAN	332	YQCKECGKSFSQRGSLAVHERLH
Q92951_HUMAN	333	YECQECGKSFRQKGS LTLHERIH
OZF_HUMAN	334	YECNECGKAFSQRTSLIVHVRIH
OZF_HUMAN	335	YECNVCGKAFSQSSSLTVHVRS
ZN07_HUMAN	336	YVCNDCGKAFSQSSSLIYHQRIH
Z151_HUMAN	337	CQCV MCGKAFTQASSLIAHV RQH
Z177_HUMAN	338	YDCKECGKAFTVPSSLQKHV RTH
OZF_HUMAN	339	FECKDCGKAFIQKSNLIRHQ RTH
Z177_HUMAN	340	YECSDCGKAFIDQSSLKKHTRSH
Z177_HUMAN	341	YECSDCGKAFFIQSSLKKHMRSH
O60792_HUMAN	342	YECKECGKA FIRSSSLAKHERIH
Z161_HUMAN	343	YACTYCSKA FRDSYHLRRHESCH
Z161_HUMAN	344	HACEMCGKA FRDVYHLNRHKL SH
MAZ_HUMAN	345	HACEMCGKA FRDVYHLNRHKL SH
O60792_HUMAN	346	FKCDECDKTFT RSTH LTQH QKIH
O60792_HUMAN	347	YKCNECDKA FSRSTH L TEHQ NTH
Z263_HUMAN	348	YKCNECGKSFRQGMHLTRHQ RTH
Z263_HUMAN	349	HKCLECGKCF SQNTHL TRHQ RTH
Z135_HUMAN	350	YEC SQCGKA FRQSTH LTQH QRIH
Z135_HUMAN	351	YEC HDCGKSFRQSTH LTQH RRIH
Z135_HUMAN	352	YECSECGKA FRQSIH LTQH LRIH
O75467_HUMAN	353	YEC AQCGKA FSQTSH LTQH QRIH
ZN07_HUMAN	354	YEC LQCGKA F SMSTQL TIHQ RVH
O95270_HUMAN	355	YPCQFCGKR FHQKSDMKKHTY IH
GFI1_HUMAN	356	YPCQYCGKR FHQKSDMKKHTFIH
O75850_HUMAN	357	FPCTECEKR FRKKTH LIRHQ RIH
Q15552_HUMAN	358	FRCDECGMR SIQKYHMERHKRTH
O43591_HUMAN	359	FRCDECGMR FIQKYHMERHKRTH
Q15552_HUMAN	360	FQCSQCDMR FIQKYLLQRHEK IH
O43591_HUMAN	361	FQCSQCDMR FIQKYLLQRHEK IH
O75850_HUMAN	362	FPCSECDKR FSKKAHLTRHLRTH
O75850_HUMAN	363	YPCAECGKR FSQKIHLGSHQKTH
O94892_HUMAN	364	FMCSECGKGFTMKRYLIVHQ QIH
O43336_HUMAN	365	YQCSECGKSFIYKQSLLDHHR IH
O43167_HUMAN	366	FKCNECGKGFAQKHS LQVHTRMH
O43167_HUMAN	367	YTC DQCGKYFSQNRQLKSHYRVH
PLZF_HUMAN	368	YECNGCDKKFSLKHQLETHYRVH
HKR3_HUMAN	369	YACPTCHKKFLSKYYLKVHNRKH
O43336_HUMAN	370	YVCNVCGKSFRHKQTFVGHQQRIH
O43336_HUMAN	371	YVCNICGKSFLHKQTLVGHQQRIH
Z134_HUMAN	372	YDCSDCGKSFGHKYT LIKHQRIH
Z200_HUMAN	373	YDCNHCGKSFNHKTNLNKHERIH
O15361_HUMAN	374	YDCNHCGKSFNHKTNLNKHERIH
ZN84_HUMAN	375	YDCNHCGKA FSRKSQ LVRHQ RTH

ZN84_HUMAN	376	FECRECGKAFSRKSQVLVTHHRTTH
ZN07_HUMAN	377	YGCRECGKAFSQQSQVLVRHQRTTH
ZN84_HUMAN	378	YRCIECGKAFSQKSQVLINHQRTTH
ZN84_HUMAN	379	YGCSECRKAFSQKSQVLVNHQRIH
ZN84_HUMAN	380	HGCIQCGKAFSQKSHLISHQMTTH
ZN84_HUMAN	381	YNCSQCGKAFSQKSQVLTSHQRTTH
ZN84_HUMAN	382	YVCSECGKAFSQKSHLISHQRTTH
Z157_HUMAN	383	FECNECGKSFGRKSQVLILHTRTH
ZN84_HUMAN	384	FECSECGKAFSRKSHLIPHQRTTH
ZN84_HUMAN	385	YECGECGKAFSRKSHLISHWRTH
Z136_HUMAN	386	YHCKECGKAYSCRASFQRHMLTH
Z136_HUMAN	387	YECKECGEAFSCIIPSMRRHMIKH
Z136_HUMAN	388	YECQECGKAFTCITSVRRHMIKH
ZN80_HUMAN	389	YECQECGKAFFPEKVDFVRHMRIH
O43338_HUMAN	390	YVCGECGKAFFMFKSKLVVRHQRTTH
O43338_HUMAN	391	YECDECGKAFFGSKSTLVVRHQRTTH
Z133_HUMAN	392	YACGECGRGFSQKSNLVAHQRTTH
Z133_HUMAN	393	YMCSECGRGFSQKSNLIIHQRTTH
Z133_HUMAN	394	YACKDCGRGFSQKSNLIRHQRTTH
Z133_HUMAN	395	YACSDCGLGFSDRSNLISHQRTTH
Z133_HUMAN	396	YACRECGRGFNKSTLIIHERTH
Z133_HUMAN	397	YVCRECGRGFSHQAGLIRHKRKH
Z133_HUMAN	398	CVCRECGQGFLQKSHLTLHQMTTH
Z133_HUMAN	399	YVCRECGKGFSSQKSAVVVRHQRTTH
O94892_HUMAN	400	YICSECGKGFPRKSNLIVHQRTTH
O94892_HUMAN	401	YICNECGKGFPGKRNLIVHQRTTH
O94892_HUMAN	402	YTCSECGKGFPLKSRLIVHQRTTH
O94892_HUMAN	403	YICSECGKGFTTKHYVIIHQRTTH
O94892_HUMAN	404	YICSECGKGFTGKSMLIIHQRTTH
O94892_HUMAN	405	YLCSECGKGFTVKSMLIIHQRTTH
O94892_HUMAN	406	YGCNECGKGFTMKSRNLIVHQRTTH
O94892_HUMAN	407	YICNECGKGFTMKSRMIEHQRTTH
O94892_HUMAN	408	FICSECGKVFTMKSRNLIEHQRTTH
O94892_HUMAN	409	YICNECGKGFAFKSNLVVHQRTTH
Z186_HUMAN	410	YECNECGKTFHQKSFLTIVHQRTTH
Z186_HUMAN	411	YECNELGKTFHCKSFLTIVHQKTH
Z186_HUMAN	412	YGCNECGKTVRCKSFLTIVHQRTTH
ZN35_HUMAN	413	YTCNECGKAFFRQRSSLTVHQRTTH
Z186_HUMAN	414	YQCSECGKTFSSQKSYLTIHHRTTH
Z157_HUMAN	415	YECSECGKTFRVKISLTQHHRTH
Z186_HUMAN	416	YKCIIECGKTFVTNQLLTLHHRTTH
Z157_HUMAN	417	YECTECGKTFSEKATLTIHQRTTH
ZN84_HUMAN	418	YACSDCRKAFFFEKSELIRHQRTTH
ZN84_HUMAN	419	YECSLCRKAFFFEKSELIRHLRTH
Z140_HUMAN	420	YECNECRKALRCHSFLIKHQRIH
ZN84_HUMAN	421	YECNECRKAFFREKSSLINHQRIH
ZN84_HUMAN	422	YECSECRKAFFRERSSSLINHQRTTH
ZN84_HUMAN	423	YECSECGKAFFGEKSSLATHQRTTH

ZN84_HUMAN	424	YECSECGKAFSEKLSLTNHQRIH
O43339_HUMAN	425	YEC SKCGKA FRGKYS LVQHQRVH
Z157_HUMAN	426	YECSECGKIFSMKKSLCQHRRTH
Z157_HUMAN	427	YECGECGKFFRMKMTLNNHQORTH
Z157_HUMAN	428	YECGECGKNFRAKKSLNQHQRIH
O43361_HUMAN	429	YKCSECGKA FSLKHNVVQHLKIH
Z134_HUMAN	430	YECSECGKA FSRKATLVQHQRH
Z134_HUMAN	431	YKCSECGKA FSRKDTLVQHQRH
Z134_HUMAN	432	YECSECGKTFSRKDNLTQHKRIH
O14709_HUMAN	433	YKCKECGKV FIRSKSLLLHQRVH
O14709_HUMAN	434	YECDECGKCFILKKSLIGHQRH
O14709_HUMAN	435	YECNECGKV FILKKSLILHQRFH
O14709_HUMAN	436	YKCNKCQKA FILKKSLILHQRIH
Z140_HUMAN	437	YACAECDKA FSRFSLILHQORTH
Z140_HUMAN	438	YGCHECGKT FGRRFSLVLHQORTH
O95878_HUMAN	439	YACAQCGKT FNNTSNLRTHQRH
O14709_HUMAN	440	YKCDMCCKH FNKISHLINHRRH
ZN83_HUMAN	441	FKCDICGKI FNKKS NLASHQRH
ZN07_HUMAN	442	HQCEDCEKI FRWRSHLI IHQRH
Z137_HUMAN	443	HKCDDCGKVLTSRSHLIRHQRIH
Z140_HUMAN	444	HECKDCNKTF SYLSFLIEHQORTH
Z189_HUMAN	445	HKCSDCGKA FSWKSHLIEHQORTH
O75802_HUMAN	446	HKCSDCGKA FSWKSHLIEHQORTH
O14709_HUMAN	447	YKCNDCGKVFSYRSNLIAHQRIH
O43309_HUMAN	448	YGCDDCGKA FSQHSHLIEHQRIH
O75123_HUMAN	449	YTCDCGKGFGQSSHLMEHQRIH
O43336_HUMAN	450	YNCTACEKA FIYKNKLVEHQRIH
O43309_HUMAN	451	YKCDVCEKA FIQRTSLTEHQRIH
O60792_HUMAN	452	YKCDQCGKG FIEGPSLTQHQRH
O43309_HUMAN	453	YKCDKCGKA FTQRSVLTEHQRIH
ZN91_HUMAN	454	YKCEECGKA FKLSTLTTHKRIH
ZN91_HUMAN	455	YKCKECGKA FKFSTLTTHKIIH
ZN91_HUMAN	456	YKCKECDKTFKRLSTLTCHKIIH
ZN91_HUMAN	457	YKCKECDKTFKRLSTLTCHKIIH
ZN85_HUMAN	458	YKCEKCGKA FNHFSLTTHKIIH
ZN85_HUMAN	459	YKCEECGKA FNRFSTLTTHKIIH
ZN43_HUMAN	460	YKCEECGKA FNQFSTLTCHKIIH
ZN43_HUMAN	461	YTCEECGKV FNWSSRLTTHKRIH
ZN43_HUMAN	462	YKCEECGKA FNKSSILTTHKIIH
O75437_HUMAN	463	YKWEKFGKA FNRSSHLTTDKITH
O43345_HUMAN	464	YKCEEGGKA FNWSSLTYYKSAH
ZN91_HUMAN	465	YKCEECGKA FNQSSNLTTHKIIH
ZN91_HUMAN	467	YKCEECGKA FNRSSKLTTHKIIH
Q02313_HUMAN	468	YKCEECGKA FNQSSTLTTHNIIH
ZN91_HUMAN	469	YKCEECGKA FNHSSSLSTHKIIH
ZN43_HUMAN	470	YKCEECGKA FKLSTLTSTHKIIH
ZN91_HUMAN	471	YKCEECGKA FSQSSTLTTHKIIH
Q02313_HUMAN	472	YKCEECGKA FNQSSTLTTHKRIH

O95780_HUMAN	473	YKCEECGKAFNSSSILTEHKVIH
O95779_HUMAN	474	YKCEECGKAFNSSSILTEHKVIH
ZN91_HUMAN	475	YKCKECGKAFKHSSALAKHKIIH
ZN85_HUMAN	476	YKCKECGKAFKHSSTLTCHKIIH
ZN85_HUMAN	477	YKCEECDKAFKWSSVLTKHKIIH
ZN43_HUMAN	478	YKCEECGKAFKWSSTLTCHKIIH
ZN85_HUMAN	479	YKCEECGKGFKWPSTLTIIHKIIH
ZN91_HUMAN	480	YKCGECGKAFKESSALTKHKIIH
ZN91_HUMAN	481	YKCEECGKAFRKSSTLTEHKIIH
ZN91_HUMAN	482	YKCEECGKAFRQSSSTLTCHKIIH
Q02313_HUMAN	483	YKCGECGKAFNQSSALNTHKIIH
ZN91_HUMAN	484	CKCKECEKTFHWSSTLTNHKEIH
O75437_HUMAN	485	YKCKECGKTFNWSSTLTNHRKIY
ZN91_HUMAN	486	YKCKECGKAFSNSSTLANHKITH
ZN91_HUMAN	487	YKCKECGKAFSNSSTLANHKITH
O43345_HUMAN	488	YKCKECGKTFIKVSTLTTHKAIH
O43345_HUMAN	489	YKCEECGKTFSKVSTLTTHKAIH
O43345_HUMAN	490	YKCEECGKTFSKVSTLTTHKAIH
O43345_HUMAN	491	YKCEECGKAFSKVSTLTTHKAIH
O43345_HUMAN	492	YKCKECGKAFSKVSTLITHKAIH
O95270_HUMAN	493	YACRMCGKAFKRSSTLSTHLLIH
GFI1_HUMAN	494	YDCKICGKSFKRSSTLSTHLLIH
O75346_HUMAN	495	YKCIICGKAFKRSSTLTTHKKIH
ZN43_HUMAN	496	YKCKECGKAFNQYSNLTTNKKIH
ZN85_HUMAN	497	YKCKECGKAFNRSSTLTTHRKH
ZN91_HUMAN	498	YKCSEECDKAFIWSSTLTEHKRIH
ZN91_HUMAN	499	YKCEECGKAFISSSTLNGHKRIH
ZN43_HUMAN	500	YKCEECGKAFNYSSHLNTHKRIH
O95780_HUMAN	501	YKCEECGKAFNWSSILTEHKRIH
O95779_HUMAN	502	YKCEECGKAFNWSSILTEHKRIH
O43345_HUMAN	503	YKCEECGKAFNWSSNLMEHKRIH
O43345_HUMAN	504	YKCEECGKAFNWSSNLMEHKRIH
O43345_HUMAN	505	YKCEECGKAFNWSSNLMEHKKIH
O43345_HUMAN	506	YKCEECGKAFNWSSNLMEHKKIH
ZN91_HUMAN	507	FKCKECGKAFIWSSTLTRHKRIH
ZN91_HUMAN	508	FKCKECGKGFIWSSTLTRHKRIH
ZN91_HUMAN	509	YKCEECGKAFLWSSTLRRHKRIH
ZN91_HUMAN	510	YKCEECGKAFLWSSTLTRHKRIH
Q02313_HUMAN	511	YKCEAYGRAFNWSSTLNKHKRIH
ZN91_HUMAN	512	YKFEECGKAFRQSLTLNKHKIIH
Z141_HUMAN	513	YKCEECGKAFRRSTDRSQHKKIH
O75346_HUMAN	514	YKCEECGKAFNWSSDLNKHKKIH
ZN91_HUMAN	515	YKCEECGKAFNWSSSLTKHKRIH
ZN91_HUMAN	516	YKCEECGKAFNWSSSLTKHKRFH
ZN85_HUMAN	517	YKCEECGKAFNWSSSTLTCHKRIH
ZN43_HUMAN	518	YKCEECGKAFNWPSTLTCHKRIH
ZN43_HUMAN	519	YKCEECGKAFNWPSTLTCHKRIH
O75437_HUMAN	520	YKCEECGKAFFWSSTLTCHKRIH

O95780_HUMAN	521	YKCEECGKAFNWCSSTLTKHKRIH
O95779_HUMAN	522	YKCEECGKAFNWCSSTLTKHKRIH
ZN43_HUMAN	523	YKCEECGKAFSRSSNLTKHKKIH
ZN43_HUMAN	524	YKCTECGEAFSRSSNLTKHKKIH
ZN91_HUMAN	525	YKCEECGKAFSRSSTLTKHKTIH
O75437_HUMAN	526	YKCEECGKAFNRSSTFTKHKVIH
Z141_HUMAN	527	YKCEECGKAFNRFTTLTKHKRIH
Z141_HUMAN	528	YKCEECGKAFNRSTTLTKHKRIH
ZN43_HUMAN	529	CKCEKCGKAFNCPSIITKHKRIN
O43345_HUMAN	530	YKCEACGKAYNTFSILTKHKVIH
O43345_HUMAN	531	YKCEECGKAFSTFSILTKHKVIH
O43345_HUMAN	532	YKCEECGKSFSSTFSILTKHKVIH
O43345_HUMAN	533	YKCEECGKSFSSTFSVLTKHKVIH
O43345_HUMAN	534	YKCEECGKGFMFSILAKHKVIH
O43345_HUMAN	535	YKCEECGKGFSMFSILTKHEVIH
O43345_HUMAN	536	YKCEECGKGFSMFSILTKHEVIH
O43345_HUMAN	537	YKCKEKGKAFSKFSILTKHKVIH
O43345_HUMAN	538	YKCKEKGKAFSKFSILTKHKVIH
O43345_HUMAN	539	YKCKEKGKAFSKFSILTKHKVIH
O43345_HUMAN	540	YRCKEKGKAFSKFSILTKHKVIH
Z195_HUMAN	541	FKCEECDSIFKWFSDLTKHKRIH
O95780_HUMAN	542	YKCEKCDKVFKRFSYLTCHKRIH
O95779_HUMAN	543	YKCEKCDKVFKRFSYLTCHKRIH
O95780_HUMAN	544	CICEECGKTFKWFSYLTCHKRIH
O95779_HUMAN	545	CICEECGKTFKWFSYLTCHKRIH
ZN43_HUMAN	546	YKCEECGKAFNHFSILTKHKRIH
ZN91_HUMAN	547	YKCEKCCKAFNQSSILTNHKKIH
Q02313_HUMAN	548	YKCEKCVRAFNOASKLTEHKLIH
ZN85_HUMAN	549	YKSKECEKAFNQSSKLTEHKKIH
ZN43_HUMAN	550	YKCKECAKAFNQSSNLTEHKKIH
ZN85_HUMAN	551	YKCEECGKAFNQSSKLTKHKKIH
ZN85_HUMAN	552	YKCEECGKAFNQSSNLIKHKKIH
O43345_HUMAN	553	YKCEECGKAFNRSAILIKHKRIH
O43345_HUMAN	554	YKCEECGKAFNQSAILIKHKRIH
O43345_HUMAN	555	YKCEECGKAFNQSAILTKHKI IH
ZN43_HUMAN	556	YKCEVCGKAFNQFSNLTTCHKRIH
ZN43_HUMAN	557	YTCEECGKAFNQFSNLTTCHKRIH
O75346_HUMAN	558	YRCEECGKAFNQSANLTTCHKRIH
ZN85_HUMAN	559	YTCEECGKAFNQSSNLTKHKRIH
Z141_HUMAN	560	YKCKDCDKAFKRFSLNKHKKIH
Z141_HUMAN	561	YKCKECDKAFKQFSLLSQHKKIH
Q02313_HUMAN	562	YKCEECGKAFKQFSLNLDHKKIH
ZN43_HUMAN	563	YKCEECGKAFTQSSNLTTCHKI IH
ZN43_HUMAN	564	YKCEECGKAFTQSSNLTTCHKI IH
ZN85_HUMAN	565	YKCEECGKAFKQSSNLTTHKI IH
Q02313_HUMAN	566	YKCEECGKAFNQLSNLTRHKVIH
ZN85_HUMAN	567	YECEKCGKAFNQSSNLTRHKKSH
O95780_HUMAN	568	YNCEECGKAFNRCSHLTRHKKIH

O95779_HUMAN	569	YNCEECGKAFNRCSHLTRHKKIH
O95780_HUMAN	570	YTCEDCGRAFNRRHSHLTKHKTIH
O95779_HUMAN	571	YTCEDCGRAFNRRHSHLTKHKTIH
Q02313_HUMAN	572	YECEECGKAFNRSSKLTEHKYIH
ZN91_HUMAN	573	YKCEECGKAFNRSSNLTIHKFIH
ZN91_HUMAN	574	YKCEECGKAFNRSSNLTIHKFIH
ZN43_HUMAN	575	YKCEKCGKAFNRPSNLTIEHKKIH
Z141_HUMAN	576	YTCEECRKIFTSSSNFAKHKRIH
Z141_HUMAN	577	FTCEECGSIFTTSSHFAKHKIIH
Z141_HUMAN	578	YTCEECGKAFKWSLIFNEHKRIH
Z141_HUMAN	579	YTCEECGKAFRQSSKLNEHKKVH
O43345_HUMAN	580	YKCEECGKAYKWSSTLSYHKKIH
O43345_HUMAN	581	YKCEECGKAYKWSSTLSYHKKIH
O43345_HUMAN	582	YKCEECGKAYKWPSTLSYHKKIH
O43345_HUMAN	583	YKCEECGKAYKWPSTLSYHKKIH
O43345_HUMAN	584	YKCEECGKAYKWPSTLRYHKKIH
O43345_HUMAN	585	YKCEECGKGFSWSSTLSYHKKIH
O43345_HUMAN	586	YKCEECGKAFSWLSVFSKHKKIH
O43345_HUMAN	587	YKCEECGKAFSWLSVFSKHKKTH
O95780_HUMAN	588	YKCEECGKAFWCSPFVRHKKIH
O95779_HUMAN	589	YKCEECGKAFWCSPFVRHKKIH
Z195_HUMAN	590	YTCEECGNIFKQLSDLTKHKKTH
Z195_HUMAN	591	YKCEECGRAFMWFSDITKHKQTH
O43345_HUMAN	592	YKCEECGKAFSWPSRLTEHKATH
O43345_HUMAN	593	YKCEECDKAFSWPSSLTEHKATH
ZN43_HUMAN	594	YKCEECGKAFKWSSKLTEHKITH
ZN43_HUMAN	595	YKCEECGKAFKWSSKLTEHKLTH
ZN91_HUMAN	596	YKCEECGKAFSHSSALAKHKRIH
ZN91_HUMAN	597	YKCEECGKAFSHSSALAKHKRIH
ZN91_HUMAN	598	YKCEECGKAFSHSSTLAKHKRIH
ZN91_HUMAN	599	YKCEECGKAFSQPSHLTTHKRMH
ZN91_HUMAN	600	YKCEECGKAFSQSSTLTRHKRLH
ZN91_HUMAN	601	YKCEECGKAFSQSSTLTRHTRMH
Z124_HUMAN	602	YECMECGKALGFSRSLNRHKRIH
Z141_HUMAN	603	YKCDECGKAFGRSRVLNEHKKIH
ZN74_HUMAN	604	YKCDECGKAFTWSTNLLEHRRIH
Z195_HUMAN	605	YKCDECGKAYTQSSHLSEHRRIH
Z195_HUMAN	606	YKCDECGKNFTQSSNLIVHKRIH
Z195_HUMAN	607	YKCDECGKNFTQSSNLIVHKRIH
ZN80_HUMAN	608	YKCKECSVFNKNLSLLVRHQQIH
Z165_HUMAN	609	FGCKECCGRAFNLSNHLIRHQRIH
Q02313_HUMAN	610	YKCKECCGKAFNQTSHLIRHKRIH
O60792_HUMAN	611	YKCNECGRAFNQNIHLTQHKRIH
ZN74_HUMAN	612	YRCGECGKAFNQRTHLTRHHRIH
Q15776_HUMAN	613	YKCKECCGKAFTNGNTGLIQHLRIH
O43309_HUMAN	614	YKCDECGNAFRGITSLIQHQRH
O43309_HUMAN	615	YKCEECGKAFRGRTVLIRHKIIH
O75123_HUMAN	616	YVCNECGKRFSQTSNFTQHQRIH

O60792_HUMAN	617	YKCNECGKAFNGPSTFIRHHMIH
O43296_HUMAN	618	FVCSECGKAFTHCSTFILHKRAH
O43337_HUMAN	619	YECSQCRKAFTHRSTFIRHNRTH
O43296_HUMAN	620	YKCNECGKAFTHRSNFVLHNRRH
OZF_HUMAN	621	YGCNECGKAFSOFSTLALHLRIH
ZN83_HUMAN	622	YKCNERGKAFHQGLHLPIHQIIH
ZN07_HUMAN	623	YKCNECGKAFSQNSTLFQHQIIH
ZN83_HUMAN	624	YKCNECGKVFSRNSYLAQHLLIIH
ZN83_HUMAN	625	YECNKCCKVFSRNSYLVQHLLIIH
ZN83_HUMAN	626	YKCNECGKVFGLNSSLAHHRKIH
ZN83_HUMAN	627	YKCNECGKVFHQISHLAQHRTIH
ZN83_HUMAN	628	YKCNECGKVFNMSHLAQHRRIH
ZN83_HUMAN	629	YKCNECGKVFNQISHLAQHQRH
ZN83_HUMAN	630	YRCNVCGKVFHHSHLAQHQRH
ZN83_HUMAN	631	YKCDECGKVFSQNSYLAHWRIH
Z189_HUMAN	632	YKCDECGKTFVSVAHLVQHQRH
O75802_HUMAN	633	YKCDECGKTFVSVAHLVQHQRH
ZN83_HUMAN	634	YKCDECDKAFSQNSHLVQHHRH
O60792_HUMAN	635	YKCDECGKAFSQRTHLVQHQRH
O43361_HUMAN	636	YECGESSKVFKYNSSLIKHQIIH
ZN83_HUMAN	637	FKCNECGKAFSMRSSLTNHHAIH
O60792_HUMAN	638	YKCNECGKAFSYCSSLTQHRRH
Z137_HUMAN	639	YKYHDCGKVFSQASSYAKHRRH
O14709_HUMAN	640	YKCEDCGKAFSYNSSLLVHRRH
Z124_HUMAN	641	YVCMCECGKAFSCLSSLQGHIAH
O60792_HUMAN	642	YQCHECGKTFSYGSSLIQHRKIH
O60792_HUMAN	643	YDCAECGKSFSYWSSLAQHLKIH
ZN83_HUMAN	644	YKCNECGKVFSHKSSLVNHWRH
ZN83_HUMAN	645	YKCNECGKVFSHKSSLVNHWRH
Z132_HUMAN	646	YKCSECGKFFSRKSSLICHWRVH
O43339_HUMAN	647	YKCNECGKFFSQTSHLNDHRRH
O43338_HUMAN	648	YECSECGKSFSQTSHLNDHRRH
ZN45_HUMAN	649	YKCNACGKSFSYSSHLNIHCRIH
ZN45_HUMAN	650	YKCGTCGKGFSRSSDLNVHCRIH
Z263_HUMAN	651	YKCPLCGKNFSNNSNLIRHQRIH
Z202_HUMAN	652	YTCPTCGKSFSRQYHLIRHQTH
O75850_HUMAN	653	FSCPQCGKSFSRKTHLVRHQLIH
Z205_HUMAN	654	YACPLCGKSFSRRSNLHRHEKIH
O15535_HUMAN	655	HQCIIECGKSFNRRHCNLIRHQKIH
ZN24_HUMAN	656	YECVQCGKSYSQSSNLFRHQRRH
Z191_HUMAN	657	YECVQCGKSYSQSSNLFRHQRRH
Q99592_HUMAN	658	YTCTQCGKSFSQYSHNLSRHAVVH
Q13397_HUMAN	659	YTCTQCGKSFSQYSHNLSRHAVVH
Z189_HUMAN	660	YLCRQCGKSFSQLCNLIRHQGVH
O75802_HUMAN	661	YLCRQCGKSFSQLCNLIRHQGVH
Z189_HUMAN	662	YQCKECCGKSFSQLCNLTRHQRIH
O75802_HUMAN	663	YQCKECCGKSFSQLCNLTRHQRIH
Z263_HUMAN	664	YKCTLGGENFSHRSNLIRHQRIH

Z263_HUMAN	665	YKCPECGEIFAHSSNLLRHQRIH
O95878_HUMAN	666	YKCSECGKSFSRSSNRIRHERIH
Z263_HUMAN	667	YTCHECGDSFSHSSNRIRHLRTH
O43336_HUMAN	668	YVCIICGKSFIRSSDYMRHQRIH
O43336_HUMAN	669	YVCMCECGKSFIHSYDRIRHQRVH
BCL6_HUMAN	670	YRCNICGAQFNRPANLKTHTRIH
Z133_HUMAN	671	YKCGECGLSFSKMTNLLSHQRIH
ZN75_HUMAN	672	YRCSWCGKSFSHNTNLHTHQRIH
O60893_HUMAN	673	YKCNECERSFTRNRSLIEHQKIH
ZN74_HUMAN	674	YKCSECGRAFSQNHCLIKHQKIH
O14709_HUMAN	675	YACSECGKGFTYNRNLIHQRIH
Z177_HUMAN	676	YKCFQCEKAFSTSTNLIMHKRIH
O60792_HUMAN	677	YKCNECEKAFSRSENLIHQRIH
O94892_HUMAN	678	YGCTLCAKVFSRKSRLNEHQRIH
Z189_HUMAN	679	YHCTKCKKSFSRNSLLVEHQRIH
O75802_HUMAN	680	YHCTKCKKSFSRNSLLVEHQRIH
O43309_HUMAN	681	YQCTQCCKKSFSRRSILTQHOGVH
O15535_HUMAN	682	YQCSQCSKSYSRRSFLIEHQRS
Z205_HUMAN	683	YTCPACRKSFSHHSTLIQHORI
Z189_HUMAN	684	YTCIECGKSFSRSSFLIEHQRI
O75802_HUMAN	685	YTCIECGKSFSRSSFLIEHQRI
Z189_HUMAN	686	FQCNECGKSFSRSSFVIEHQRI
O75802_HUMAN	687	FQCNECGKSFSRSSFVIEHQRI
Z189_HUMAN	688	YLCTVCGKSFSRSSFLIEHQRI
O75802_HUMAN	689	YLCTVCGKSFSRSSFLIEHQRI
O14709_HUMAN	690	YECHVCRKVLTSRNLMMVHQRI
O14709_HUMAN	691	YEC DKCRKSFTSKRNLVGHQRI
ZN35_HUMAN	692	YECNECGKTFTRSSNLIVHQRI
O75123_HUMAN	693	YECNECGKSFIRSSSLIRHYQI
O43296_HUMAN	694	YECVECGKSF CWSTNLIRHAI
O43296_HUMAN	695	YECSECGKVFLSAALIH HYVI
O43337_HUMAN	696	YECTQCGKAFHRSTYLIQH SVI
O43296_HUMAN	697	YECTECGKTFIKSTHLLQH HMI
O75290_HUMAN	698	YECKECGKYFSRSANLIQH QSI
Z205_HUMAN	699	YACTDCGKRFRGRSSHLIQHQI
Z165_HUMAN	700	YECSECGKTFRVSSHLIRHFRI
Q15776_HUMAN	701	YECDECGKTFRRSSHLIGHQRSH
Q15776_HUMAN	702	YECNECGKAFSHSSHLIGHQRI
Z189_HUMAN	703	YECNYCGKTF SVSSTLIRHQRI
O75802_HUMAN	704	YECNYCGKTF SVSSTLIRHQRI
O43337_HUMAN	705	YECNACGKAFSQSSTLIRHYLI
ZN07_HUMAN	706	YECSECGKAFSRSSYLIEHQRI
Z132_HUMAN	707	YECSECGKAFAHSSTLIEHWRV
O43340_HUMAN	708	YECSECGKAFCNIIYLIIHQRFH
Z135_HUMAN	709	YECGECGKAFSQSSTLLTEHRR
O43338_HUMAN	710	YECGECGKSFSQSNNLIEHCRI
O43338_HUMAN	711	YECGKCGKSFTQHSGLILHRKSH
Z140_HUMAN	712	YECDECGKVFTWHASLIQHTKSH

Q13398_HUMAN	713	YACPECGKSFSQIYSLNSHRKVH
Q13398_HUMAN	714	YECCKCGKSFKQSSSFSSHRKVH
O43340_HUMAN	715	YECSECGKSFSHSTNLFRRHVRH
O43340_HUMAN	716	YECSECGKSFSHSTNLYRHRSAH
O43340_HUMAN	717	YECSECGKSFSQSSGLLRHRRVH
O43340_HUMAN	718	YKCSECGKSFSQSSGFLRHRKAH
O43340_HUMAN	719	YECSECGKVFSQSSGLFRHRRAH
O43340_HUMAN	720	YECDECGKSYSQSSALLQHRRVH
Q13398_HUMAN	721	YECSECGKSFSQSSSLIQHRRVH
Q13398_HUMAN	722	YECGECGKSFSQRSNLMQHRRVH
Z132_HUMAN	723	YECSECRKSFSRSSSLIQHWRIH
Z132_HUMAN	724	YECSQCGKSFSRSSSALIQHWRVH
Q13398_HUMAN	725	HECNECGKSFSRSSSLIHHRRRLH
O43339_HUMAN	726	YKCGECGNSFSQSAILNQHRRIH
O43339_HUMAN	727	YKCGDCGKSFSQSSILIQHRRIH
O60765_HUMAN	728	YRCEECGISFGQSSALIQHRRIH
O43338_HUMAN	729	YECGQCGKSFSCLKGLIQHQLIH
O43339_HUMAN	730	YECGQCGKSFSQKSGLIQHQQVH
O43338_HUMAN	731	YDCGQCGKSFIQKSSLIQHQQVH
Q13398_HUMAN	732	YQCSQCGKSFGCKSVLIQHQRVH
O43340_HUMAN	733	YVCSECGKSFGQKSVLIQHQRVH
O43340_HUMAN	734	YDCSECGKSFRQVSVLIQHQRVH
Q13398_HUMAN	735	YECSECKSFSCKSNLIKHLRVH
O43339_HUMAN	736	YECGQCGKSFSQKATLIKHQVRH
O43338_HUMAN	737	YVCGQCGKSFSQRATLIKHHRVH
O43339_HUMAN	738	YECSQCGKSFSQKATLVKHQVRH
Q13398_HUMAN	739	YECSECGKSFSQNFSLIYHQVRH
O43340_HUMAN	740	YECVCGKSFIKTHLIRHQTVH
O43340_HUMAN	741	YECSECEKSFSCKTDLIRHQTVH
O43340_HUMAN	742	YECRECGKSFTRKNHLIQHKTVH
Z189_HUMAN	743	HKCEECGKGFVRKAHFIIHQVRH
O75802_HUMAN	744	HKCEECGKGFVRKAHFIIHQVRH
O43340_HUMAN	745	HECSECGKSFSRKTHLTQHQRVH
O43309_HUMAN	746	YQCKECGKSFSQSGLIQHQRH
Q15776_HUMAN	747	YQCNQCGKAFSQSAGLILHQRIH
O15535_HUMAN	748	YHCKECGKVFSQSAGLIQHQRH
O60792_HUMAN	749	YNCNECRKTFSQSTYLIQHQRH
Q15776_HUMAN	750	YHCKECGKAFSQNTGLILHQRIH
ZN84_HUMAN	751	YGCNECGRAFSEKSNLINHQRIH
Q15776_HUMAN	752	YKCNECGRAFSQKSGLIEHQRIH
Z189_HUMAN	753	HKCDECGKAFSRNSGLIQHQRIH
O75802_HUMAN	754	HKCDECGKAFSRNSGLIQHQRIH
Z189_HUMAN	755	HKCEECGKAFSRSSGLIQHQRIH
O75802_HUMAN	756	HKCEECGKAFSRSSGLIQHQRIH
ZN24_HUMAN	757	YKCLECGKAFSQNSGLINHQRIH
Z191_HUMAN	758	YKCLECGKAFSQNSGLINHQRIH
OZF_HUMAN	759	YQCSECGKAFSQKSHHIRHQKIH
Q15776_HUMAN	760	YQCNCECGKAFTQRSSLIHQRIH

ZN35_HUMAN	761	YDCSECGKAFSQLSSLIVHQRH
ZN07_HUMAN	762	YRCEECGKAFGQSSSLIHHQRH
O60765_HUMAN	763	FKCNTCGKTFRQSSSRIAHQRH
OZF_HUMAN	764	FKCSECGTAFGQKKYLIKHQNIH
OZF_HUMAN	765	FECNECGKAFSQKQYVIKHQNT
Q92951_HUMAN	766	FECTHCGKSFRAGNLTTHQRH
OZF_HUMAN	767	FECNECGKSFSQKENLLTHQKI
ZN74_HUMAN	768	FKCNECGKAFFSHAYLIVHRRH
ZN74_HUMAN	769	FKCADCGKGFSCSHAYLLVHRRH
O60765_HUMAN	770	FKCSECGRAFSQSASLIQHERH
ZN35_HUMAN	771	FECHECGKAFIQSANLVVHQRH
ZN35_HUMAN	772	FTCSVCGKGFSSQSANLVVHQRH
ZN35_HUMAN	773	FACNDCGKAFTQSANLIVHQRSH
O14709_HUMAN	774	YKCNECGKDFSQNKNLVVHQRMH
O14709_HUMAN	775	YKCDECGKTFAQTTYLIDHQRH
O14709_HUMAN	776	YKCNECGKVFSQNAYLIDHQRH
O14709_HUMAN	777	YKCTECGKAFTQSAYLFDHQRH
O14709_HUMAN	778	YKCNECGKAFSQSAYLLNHQRH
Z157_HUMAN	779	YQCNECGKSFRVHSSLGIHQRH
O60765_HUMAN	780	YNCNECGKALSSHSTLIHERH
EVI1_HUMAN	781	YKCDQCPKAFNWKSNLIRHQMSH
Q15776_HUMAN	782	YQCNVCGKAFSYRSALLSHQDIH
O43309_HUMAN	783	YECNECGKAFVYNSSLVSHQEIH
Z200_HUMAN	784	YGCKKCGRRFGRLSNCTRHEKTH
O15361_HUMAN	785	YGCKKCGRRFGRLSNCTRHEKTH
ZN07_HUMAN	786	YKCNDGCGKAFNRSSRLTQHQKI
ZN74_HUMAN	787	YQCGSCGKAFTCHSSLTVHEKIH
ZN35_HUMAN	788	YVCSKCGKAFTQSSNLTVHQKI
Z140_HUMAN	789	YECIECGKAFFRFSHLTRHQSIH
O60893_HUMAN	790	YQCNMCGKAFFRNSHLLRHQRH
Q13396_HUMAN	791	YSCTECEKSFVQKQHLLQHQKI
O43361_HUMAN	792	YECTQCAKAFVRKSHLVQHEKIH
O43361_HUMAN	793	YECTECEKAFVRKSHLVQHQKI
O75123_HUMAN	794	YECKECGKAFLQKAHLTEHQKI
O75290_HUMAN	795	YECKECGKGFNRGHLLIQHQKI
O75290_HUMAN	796	YECKECGKGFNRGHLLIQHQKI
O75290_HUMAN	797	FECKECGKAFLHMQLIRHQKLH
O75290_HUMAN	798	FECKECGKAFLHMLIRHQKLH
O75290_HUMAN	799	FECKECGKAFLHIQFTRHQKFH
O75290_HUMAN	800	YECKECGKAFLRLYLQLSQHQKTH
Z140_HUMAN	801	YECTECGKAFSRASNLTRHQRH
O43296_HUMAN	802	YECVECGKAFTRMGLTRHKRIH
O43296_HUMAN	803	YECMECGKAFNRKSYLTQHQRH
O14913_HUMAN	804	HECVECGKRFSSSSRLQEHQKI
EVI1_HUMAN	805	HACPECGKTFATSSGLKQHKHIH
O15535_HUMAN	806	YECNECGKAFSRSSGLFNHRGIH
Z132_HUMAN	807	YECNDCGKAFSNSSTLIQHQKVH
Z132_HUMAN	808	YECIQCGKAFSERSTLVRHQKVH

Z132_HUMAN	809	YECDECGKAFSNRSHLIRHEKVVH
Z124_HUMAN	810	YECQKCGKAFSRASTLWKHKKTH
ZN35_HUMAN	811	FKCNECEKAFSYSSQLARHQKVH
O60792_HUMAN	812	FECSECGKAFSYLSNLTQHQQKTH
O75467_HUMAN	813	FRCSECGKAFSHGSLNSQHRKIH
O75467_HUMAN	814	FACPQCGRAFSHSSNLTQHQLLH
OZF_HUMAN	815	FACKVCGKVFESHKSNLTEHEHFH
Z132_HUMAN	816	YECSQCGKLFSHLCNLAQHKKIH
O60765_HUMAN	817	YECNTCGKLFNHRSSLTNHYKIH
O60792_HUMAN	818	YECAECGKAFRHCSSLAQHQQKTH
O43336_HUMAN	819	CECSECGKCFRHRTSLIQHQKVH
O43336_HUMAN	820	CECNECGKVFESHQKRLLEHQKVH
O95878_HUMAN	821	YECTECGRTFSDISNFGAHQORTH
O60792_HUMAN	822	YECNECGKAFSQHSNLTQHQQKTH
O43309_HUMAN	823	YHCNDCGKAFSQKAGLFHHIKIH
O43336_HUMAN	824	YECSDCGKAFISKQTLKHHKIH
O60893_HUMAN	825	YECDDCGKTFSQSCSLLEHHKIH
O43338_HUMAN	826	FECDECGKSFSQRTTLNKHHKVH
O75123_HUMAN	827	YVCSYCGKGFIQRSNFLQHQKIH
O60792_HUMAN	828	YTCNECGKAFSQRGHFMEHQKIH
ZN42_HUMAN	829	YTCDVCGKVFSQRSNLLRHQKIH
O14709_HUMAN	830	YGCNDCSKVFRQRKNLTVHQKIH
O43361_HUMAN	831	YVCSECGKAFLTQAHLVGHQKIQ
O43361_HUMAN	832	YTCSECGKAFLTQAHLVGHQKIH
O43361_HUMAN	833	YECTQCGKAFLTQAHLVGHQKTH
Z157_HUMAN	834	YECGECAKTFSARSYLIAHQKTH
O75123_HUMAN	835	YECNECGKAFFLSSYLIRHQKIH
Q13398_HUMAN	836	YECNECGKFFTYYSFIIHQRVH
O43361_HUMAN	837	YKCSKCGKFFRYRCTLSRHQKVH
O43361_HUMAN	838	YECNKCCKGKFFMYNSKLIRHQKVH
Z132_HUMAN	839	YECNECGKFFSQNSILIKHQKVH
Q13396_HUMAN	840	YECGYCGKSFSHPSDLVRHQRIH
O75467_HUMAN	841	YACPVCCKAFRHSSSLVRHQRIH
Z165_HUMAN	842	HQCNECGKAFRHSSSKLARHQRIH
Z205_HUMAN	843	YHCLDCGKSFSHSSSLTAHQORTH
Z135_HUMAN	844	YACRDCGKAFTHSSSLTKHQORTH
Z135_HUMAN	845	YECNDCGKAFSHSSSLTKHQRIH
Z135_HUMAN	846	YQCGECGKAFSHSSSLTKHQRIH
ZN74_HUMAN	847	FDCSQCWKAFSCHSSLIMHQRIH
ZN74_HUMAN	848	YTCGECGKAFSCHSSLNVHQRIH
ZN35_HUMAN	849	YECKECGKAFSCHFSLIVHQRIH
O43309_HUMAN	850	YKCNECGKAFGRWALNQHQRLLH
ZN24_HUMAN	851	YGCVECGKAFSRSSILVQHQRVH
Z191_HUMAN	852	YGCVECGKAFSRSSILVQHQRVH
O43296_HUMAN	853	YKCSECGKAFSRSSSLTQHQRMH
ZN75_HUMAN	854	FKCQECGKSFRVSSDLIKHHRIH
O75290_HUMAN	855	FVCKECGMAFRYHYQLIEHCQIH
O75467_HUMAN	856	FVCTQCGRAFRERPALFHHQRIH

ZN74_HUMAN	857	FKCEKCGEMFNWSSHLTEHQRLH
ZN85_HUMAN	858	FKCTKCGKSFGMISCLTEHSRIH
ZN43_HUMAN	859	FKCKEKGKSFCLPHLAQHKKI IH
Z195_HUMAN	860	FKCQCEGKSFQMLSFLTEHQKIH
ZN07_HUMAN	861	FKCDECGKAFRWISRLSQHQLIH
Z189_HUMAN	862	HKCGECGKAFLSTYLIQHQKIH
O75802_HUMAN	863	HKCGECGKAFLSTYLIQHQKIH
ZN07_HUMAN	864	FKCTECGKAFLSSKLIQHQRH
O75290_HUMAN	865	FECCKEKGKAFLLTKLVRHQKIH
O75290_HUMAN	866	FECCKEKGKVFSLPTQLNRHKNIH
O75290_HUMAN	867	FECRECGKAFSLNQLNRHKNIH
O75290_HUMAN	868	FECCKEKGKAFSNRAHLIQHYI IH
O43296_HUMAN	869	FECCKEKGKAFSNRKDLIRHFSIH
O62425_CAEL	870	FVCKVCGKAFRQASTLCRHKI IH
O75123_HUMAN	871	FECCKDCGKAFLIQSSKLLHQLI IH
O75290_HUMAN	872	FECCKEKGKFFRRGSNLNQHRSIH
O75290_HUMAN	873	FECCKEKGKSFNRSSNLVQHQSIH
O75290_HUMAN	874	FECCKEKGKSFNRSSNLVQHQSIH
O75290_HUMAN	875	FECQDCGKAFNRGSSLVQHQSIH
O94892_HUMAN	876	FVCSECRKAFSSKRNLIVHQRTH
O14709_HUMAN	877	FECSECGRAFSSNRNLIEHKRIH
Z135_HUMAN	878	YECNQCGRASARATLLIEHQRIH
Z157_HUMAN	879	FECQCEGKAFCRKAHLTEHQRTH
Z157_HUMAN	880	FECNECGKAYCRKSNLVEHLRIH
O75123_HUMAN	881	FECNECGKAFLRSSKLIQHQRH
ZN42_HUMAN	882	FRCAECGQSFRQRSNLLQHQRH
ZN42_HUMAN	883	FACPECGQSFRQHANLTQHRRH
ZN42_HUMAN	884	FACAECGQSFRQRSNLTQHRRH
ZN42_HUMAN	885	--CAECGKAFRQRPTLTQHRLRH
ZN42_HUMAN	886	YACPECGKAFRQRPTLTQHRLRH
O14913_HUMAN	887	YKCEEKGNSFYYPAMLKQHQRH
Z174_HUMAN	888	YTCGECGNCFRQSTLKLHQRIH
PLZF_HUMAN	889	YECFECGSCFRDESTLKSHKRIH
BCL6_HUMAN	890	YPCEICGTRFRHLQTLKSHLRIH
O43296_HUMAN	891	FECLECGKAFNHRSYLKRHQRIH
O43337_HUMAN	892	YKCLECGKAFFKRRSYLMQHHPH
O43296_HUMAN	893	YECLECGKVFKHRSYLMWHQQTH
O75123_HUMAN	894	YECCKEKGKAFLRHRSDLIEHQRIH
O43336_HUMAN	895	YECCKEKGKAFLHKKRLLEHQRIH
Z157_HUMAN	896	YECSECGNAFYVKVRLIEHQRIH
Z157_HUMAN	897	YECNECGNAFYVKARLIEHQRMH
OZF_HUMAN	898	FVCCKEKGKTFSGKSNLTEHEKIH
Z134_HUMAN	899	YKCSDCGKVFRHKSTLVQHEH
O60893_HUMAN	900	YECEDCGKTFIGSSALVHQRVH
O43339_HUMAN	901	YECSECGKLFRQNSSLVDHQKIH
O43338_HUMAN	902	FECSECGKFFRQSYTLVEHQKIH
O43338_HUMAN	903	YECGECGKLFRQSFSLVVHQRIH
O43361_HUMAN	904	YECSECGKLFMDSF TLGRHQRVH

O43361_HUMAN	905	YECSECGKFFRDSYKLI IHQRVH
O43361_HUMAN	906	YECNECGKFFLDSYKLVIHQRIH
O43336_HUMAN	907	YECSECGKGFYLEVKLLQHQRH
ZN07_HUMAN	908	YECAECGKVFRLC SQLNQHQRIH
Z132_HUMAN	909	HVCKECGKAFSHSSKLRKHQKFH
TYY1_HUMAN	910	HVCAECGKAFVESSKLRHQLVH
O15391_HUMAN	911	HVCAECGKAFLESSKLRRHQLVH
O94892_HUMAN	912	HVCSECGKAFVKKSQ L TDHERVH
ZFX_HUMAN	913	HICVECGKGFRHPSELKKHMRH
ZFY_HUMAN	914	HICVECGKGFRYPSELRKHMRH
Q15558_HUMAN	915	HICVECGKGFRHPSELRKHMRH
Z135_HUMAN	916	YECHECLKGFRNSSALTKHQRIH
ZN74_HUMAN	917	YTCGECGKA FRQSSSLTLHRRWH
Z174_HUMAN	918	YQCGQCGKSFRQSSNLHQHRLH
Z195_HUMAN	919	YQCEECEGKVFR TCSSLSNHKRTH
HKR3_HUMAN	920	FQCHLCGKTFR TQASLDKHNRTH
O43337_HUMAN	921	YDCMACGKA FRCSSELIQHQRH
O60765_HUMAN	922	YLCNECGNTFKSSSSSLRYHQRIH
O60765_HUMAN	923	YKCNECGKTFR CNSSLSNHQRH
Z140_HUMAN	924	YKCNECGKA FSSSGSELIRHQITH
Q14585_HUMAN	925	YECKECGKA F SFGSGLIRHQI IH
Q14585_HUMAN	926	YICNECGKA F SFGSALTRHQRIH
Q14585_HUMAN	927	YECKECGKS FSSSGSALNRHQRIH
Q14585_HUMAN	928	YECKACGMA FSSSGSALTRHQRIH
Q14585_HUMAN	929	YECKECGKS F S FESALIRHHRH
Q14585_HUMAN	930	YECKECGKT FSSSGSDLTQHHRH
Q14585_HUMAN	931	YVCKECGKA FN SSGSDLTQHQRH
Q14585_HUMAN	932	YECKECGKA F YSGSSLTQHQRH
Q14585_HUMAN	933	FECKECGKA F GSGSNLTHHQRIH
Q14585_HUMAN	934	YECKECGKA F GSGGANLAYHQRIH
Q14585_HUMAN	935	YECIDCGKA F GSGSNLTQHRRH
Q14585_HUMAN	936	YECKECGKA F GSGSKLIQHQLIH
Q14585_HUMAN	937	YECKECEKA FRSGSKLIQHQRMH
ZN80_HUMAN	938	YECKECGKT FYYNSSLTRHMKIH
ZN80_HUMAN	939	YECKECGKG FYYSYSLTRHTRSH
Z165_HUMAN	940	YECNECGKS FAESSDLTRHRRH
Z202_HUMAN	941	YKCTICGKS F SQKSVLTTHQRH
O43167_HUMAN	942	YTCEICGKS FTAKSSLQTHIRH
Q92618_HUMAN	943	HTCCICGKS FPFQSSLSQHMRKH
Q15776_HUMAN	944	HKCDECGKS FAQSSGLVRHWRIH
O15535_HUMAN	945	HKCDECGKS FTQSSGLIRHQRIH
O60893_HUMAN	946	HYCHECGKS FAQSSGLTKHRRH
ZN24_HUMAN	947	HICDECGKH FSQGSALILHQRIH
Z191_HUMAN	948	HICDECGKH FSQGSALILHQRIH
Z140_HUMAN	949	YACKECGKT FSQISNLVKHQMIH
Q14585_HUMAN	950	YECKECGK D F S FVSVLVRHQRIH
O75123_HUMAN	951	FECKECGKG FSQSSLLIRHQRIH
UKLF_HUMAN	952	FKCNHCDRC FSRSDHLALHMKRH

O95600_HUMAN	953	FRCTDCNRSFSRSDHLSLHRRRH
SP2_HUMAN	954	YACAQCQKRFMRS DHLT KHYKTH
SP4_HUMAN	955	YACPECSKRFMRS DHLT SKHV KTH
O60402_HUMAN	956	YACPECSKRFMRS DHLT SKHV KTH
O75411_HUMAN	957	YACPMCDRRFMRS DHLT KHARRH
Q13118_HUMAN	958	YACPMCDRRFMRS DHLT KHARRH
O14901_HUMAN	959	YACPVCDDRRFMRS DHLT KHARRH
BTE1_HUMAN	960	YACPLCEKRFMRS DHLT KHARRH
SP2_HUMAN	961	FVCNWF FCGKR FTR SDEL QRHARTH
SP4_HUMAN	962	FICNWM FCGKR FTR SDEL QRHRRTH
O60402_HUMAN	963	FICNWM FCGKR FTR SDEL QRHRRTH
EZF_HUMAN	964	YHCDWDGCGWK FAR SDEL TRHYRKH
O95600_HUMAN	965	YKCTWDGCSWK FAR SDEL TRHFRKH
UKLF_HUMAN	966	YKCSWEGCEWR FAR SDEL TRHYRKH
EKLF_HUMAN	967	YACTWEGCGWR FAR SDEL TRHYRKH
BTE2_HUMAN	968	YKCTWEGCDWR FAR SDEL TRHYRKH
O14901_HUMAN	969	FNCSDWDGCDKK FAR SDEL SRHRRTH
Q13118_HUMAN	970	FSCSWKGCERR FAR SDEL SRHRRTH
O75411_HUMAN	971	FSCSWKGCERR FAR SDEL SRHRRTH
BTE1_HUMAN	972	FPCTWPDCLKKFSR SDEL TRHYRTH
EGR4_HUMAN	973	FACPVE SCVRS FAR SDEL NRHLRIH
EGR2_HUMAN	974	YPCPAEGCDRRFSR SDEL TRHIRIH
EGR1_HUMAN	975	YACPVE SCDRRFSR SDEL TRHIRIH
EGR3_HUMAN	976	HACPAEGCDRRFSR SDEL TRHLRIH
Q16256_HUMAN	977	YQCDFKDCERR FFSR SDQLKR HQRRH
WT1_HUMAN	978	YQCDFKDCERR FFSR SDQLKR HQRRH
Q15881_HUMAN	979	YQCDFKDCERR FFSR SDQLKR HQRRH
Q15881_HUMAN	980	FQCKACQRKFSR SDHLKTHTRTH
Q16256_HUMAN	981	FQCKTCQRKFSR SDHLKTHTRTH
WT1_HUMAN	982	FQCKTCQRKFSR SDHLKTHTRTH
EGR4_HUMAN	983	FQCRICLRNFSR SDHLTSHV RTH
EGR3_HUMAN	984	FQCRICMR SFSR SDHLTTHIRTH
EGR2_HUMAN	985	FQCRICMRNFSR SDHLTTHIRTH
EGR1_HUMAN	986	FQCRICMRNFSR SDHLTTHIRTH
EVI1_HUMAN	987	YTCRYCGKIFPR SANLTRHLRTH
O95878_HUMAN	988	YRCTVCGKHFSR SSNLIRHQKTH
Z140_HUMAN	989	YVCKVCNK SFSW SSNLAKHQ RTH
O60893_HUMAN	990	YECEE CGKVFSH SSNLIKHQ RTH
Z135_HUMAN	991	YECSE CGKSFSFR SSFSQHERTH
O95878_HUMAN	992	YICCEE CGKSFSN SSSFGVHH RTH
ZN80_HUMAN	993	CKCSE CGKTFTYRSVFFRHS MTH
ZN80_HUMAN	994	YECSE CGKTFSYHSVFIQHRVTH
Z135_HUMAN	995	YGCNE CGKSFSH SSSLSQHERTH
Z135_HUMAN	996	YGCNE CGKTFSH SSSLSQHERTH
Z263_HUMAN	997	YKCPE CGKSFSR SSSLVIHERTH
Z263_HUMAN	998	YKCSE CGESFSR SRLMSHQ RTH
Z202_HUMAN	999	CRCNE CGKSFSR RDHLVRHQ RTH
ZN74_HUMAN	1000	FKCSDCEKAFNSR SRLTLHQ RTH

ZN42_HUMAN	1001	FACPECGQRFSQRLKLTRHQORTH
Z205_HUMAN	1002	YPCPECGKCFSQRSNLIHNRTH
ZN75_HUMAN	1003	FKCDECGKRFIQNSHLIKHQORTH
ZN07_HUMAN	1004	FKCDECGKGFVQGSHLIQHQRIH
O15090_HUMAN	1005	YPCPLCGKRFRFNSILSLHMIRTH
O94892_HUMAN	1006	YRCSECGKGFIVNSGLMLHQORTH
O95270_HUMAN	1007	HKCQVCGKAFSQSSNLITHSRKH
GFI1_HUMAN	1008	HKCQVCGKAFSQSSNLITHSRKH
Z135_HUMAN	1009	YKCQECGKAFSHSSALIEHHRTH
O60765_HUMAN	1010	FKCKECSKAFSQSSALIQHQITH
O60765_HUMAN	1011	CKCKVCGKAFRQSSALIQHQRMH
O60792_HUMAN	1012	CKCNECGKAFSYCSALIRHQORTH
Z151_HUMAN	1013	YVCERCGRFVQSSQLANHIRHH
EVI1_HUMAN	1014	YECENCAKVFTDPSNLQRHIRSQH
Z205_HUMAN	1015	YVCDRCAKRFTRRSDLVTHQGTH
Z205_HUMAN	1016	HKCPICAKCFTQSSALVTHQORTH
Z124_HUMAN	1017	YGCTICEKVFNIPSSFQIHQRNH
Z200_HUMAN	1018	YTCPLCGKQFNESSYLIHQORTH
O15361_HUMAN	1019	YTCPLCGKQFNESSYLIHQORTH
ZN07_HUMAN	1020	YKCNKCTKAFGCSSRLIRHQORTH
Z263_HUMAN	1021	YQCNICGKCFSCNSNLHRHQORTH
Q13134_HUMAN	1022	YKCELCPYSSSQKTHLTRHMIRTH
Q13127_HUMAN	1023	YKCELCPYSSSQKTHLTRHMIRTH
CTCF_HUMAN	1024	FQCSLCSYASRDYKLRHMIRTH
Q99592_HUMAN	1025	YTCSLCGKTFSCMYTLKRHERTH
Q13397_HUMAN	1026	YTCSLCGKTFSCMYTLKRHERTH
Q60765_HUMAN	1027	YKCSLCEKTFINTSSLRKHEKNH
ZN74_HUMAN	1028	YKCSACEKAFFSCSSLLSMHLRVH
ZN75_HUMAN	1029	YKCQQCDRRFRWSSDLNKHFMTH
Z189_HUMAN	1030	YQCNQCKQSFSQRRSLVKHQRIH
O75802_HUMAN	1031	YQCNQCKQSFSQRRSLVKHQRIH
Z186_HUMAN	1032	YACNCCEKLFSYKSSLTIHQRIH
Z186_HUMAN	1033	YACDHCEKAFFSHKSKLTVHQORTH
ZN84_HUMAN	1034	YECRDCEKAFFSQKSQLNTHQRIH
O60792_HUMAN	1035	YQCNKCEKTFSSQSSHLTQHQRIH
O75066_HUMAN	1036	YACQYCDVFAQSIELSRHVIRTH
O95878_HUMAN	1037	YRCDICGKSFSQSATLAVHHIRTH
P91805_SARPE	1038	YQCKVCQKRFPQLSTLHNHERTH
Z133_HUMAN	1039	YACKECGRCFRQRTTLVNHQORTH
Z133_HUMAN	1040	YVCGVCGHSFSQNSTLISHRRTH
O43336_HUMAN	1041	YVCIIECGKSLSSKYSLVHQRTH
O75467_HUMAN	1042	YACAQCGRRFCRNSHLIQHERTH
Z124_HUMAN	1043	YECKQCGKAFFSRSSHLRDHERTH
Z177_HUMAN	1044	YECNQCGKSFSSTGSYLIVHKRTH
Z177_HUMAN	1045	YECDHCGKSFSQSSHLNVHKRTH
ZN84_HUMAN	1046	YACGNCGKTFPQKSQFITHHRTH
Z135_HUMAN	1047	YECHECGKAFTQITPLIQHQORTH
Z135_HUMAN	1048	YECNQCGRAFSQLAPLIQHQRH

Z135_HUMAN	1049	YKCTQCGRTFNQIAPLIQHQRTH
O60893_HUMAN	1050	YQCDTCGKGFTRTSYLVQHQRSH
O43337_HUMAN	1051	YKCKQCGKGFNWKWYLVRHQRVH
Z205_HUMAN	1052	YRCEQCGKGFSWHSVLVTHRRTH
Z202_HUMAN	1053	YRCDDCGKHFRWTSDLVRHQORTH
ZN45_HUMAN	1054	YRCDVCGKRFRQRSYLOAHQORVH
ZN45_HUMAN	1055	YQCDACGKGFSRSSDFNIHFRVH
Z239_HUMAN	1056	YQCYECGKGFSQSSDLRIHLRVH
Z239_HUMAN	1057	YKCDKCGKGFSQSSKLHIHQRVH
Z239_HUMAN	1058	YHCGKCGKGFSQSSKLLIHQRVH
Z239_HUMAN	1059	YKCGECGKGFSQSSNLHIHRCIH
O15322_HUMAN	1060	YKCDMCGKEFSQSSCLQTHERVH
Z239_HUMAN	1061	YACQYCGKNFSQSSELLELHQRDH
ZN07_HUMAN	1062	YPCKECGKAFSQSSTLAQHQRMH
Z133_HUMAN	1063	YVCKTCGRGFSCLKSHLSRHRKTH
Z133_HUMAN	1064	YVCGVCGRGFSCLKSHLNRHQNIH
Z133_HUMAN	1065	YVCGVCEKGFSCLKKSLARHQKAH
EVI1_HUMAN	1066	YRCKYCDRSFSSISSNLQRHVRNIH
RRE1_HUMAN	1067	YKCQTCERTFTLKHSLVRHQRIH
O75850_HUMAN	1068	YACAQCGRRFSRKSHLGRHQAVH
O75850_HUMAN	1069	HACAVCARSFSSKTNLVRHQAIH
O75850_HUMAN	1070	YQCAQCARSFTHKQHLVRHQRVH
ZN42_HUMAN	1071	FVCSECGRSFSRSSHLLRHQLTH
Z132_HUMAN	1072	FECSECGRDFSQSSHLLRHQKVH
ZN35_HUMAN	1073	YECEKCGAAFI SNSHLMRHHRTH
Z132_HUMAN	1074	YECSECGRAFSSNSHLVRHQRVH
Z202_HUMAN	1075	YKCMCECGKSYTRSSHLLARHQKVH
Z134_HUMAN	1076	YECSECGKAYSLSSHLLNRHQKVH
Z239_HUMAN	1077	YECCKCGKGFSQSSNLHSHQRVH
Z165_HUMAN	1078	YECSECGRAFSQSSNLSQHQRTH
Z132_HUMAN	1079	YECSECGRAFNNNSNLAQHQKVH
Z239_HUMAN	1080	YECSECGMSFSQRSNLHIHQRDH
O00153_HUMAN	1081	HQCQVCGKTFSQSGSRNVHMRKH
Q13398_HUMAN	1082	YVCGECGKSFSHSSNLKNHQRVH
O15322_HUMAN	1083	YKCEICGKSFCRLRSSNLNRHYMVH
O75123_HUMAN	1084	FKCAQCGKAFCHSSDLIRHQRVH
O14913_HUMAN	1085	YKCEECDKAFLYHSFLRRHKAVH
O14913_HUMAN	1086	YKCEECDKAFLHHSYLRKHQAVH
ZN83_HUMAN	1087	FKCNECGKLFRDNSYLVRHQRFH
O15322_HUMAN	1088	HTCNECGKSFCYISALRIHQRVH
O60792_HUMAN	1089	FGCNDGKSFYRSALNKHQRLH
Z137_HUMAN	1090	YKCNKCGKIFRHRSYLAVYQORTH
O75123_HUMAN	1091	YVCNVCGKDFIHYSGLIEHQRVH
Z134_HUMAN	1092	YKNECGKYFSHHSNLIVHQRVH
O43361_HUMAN	1093	FECSCGKFFSHRSTLNMHQRVH
Z134_HUMAN	1094	FECIECGKFFSRSSDYIAHQRVH
Z134_HUMAN	1095	FVCSKCGKDFIRTSHLVRHQRVH
O14913_HUMAN	1096	YKCQECGKSFCYRSYLREHYRMH

Z174_HUMAN	1097	YKCDDCGKSFTWNSELKRHKRVH
O60765_HUMAN	1098	YRCKECGKSFSRRSGLFIHQKIH
O43167_HUMAN	1099	YSCGICGKSFSDDSSAKRRHCILH
O43829_HUMAN	1100	FVCEMCTKGFTTQAHLEHLKIH
O00403_HUMAN	1101	FVCEMCTKGFTTQAHLEHLKIH
O75626_HUMAN	1102	FKCQTCNKGFTQLAHLQKHVLVH
O15322_HUMAN	1103	FKCEQCGKGFRCAILQVHCKLH
BCL6_HUMAN	1104	YKCETCGARFVQVAHLRAHVLIH
Z195_HUMAN	1105	YKCEKCGKAFTQFSHLTVHESI
ZN85_HUMAN	1106	YKCKKCGKAFNQSAHLTTHEVIH
Z239_HUMAN	1107	YKCEKCGKGFTRSSSLIHHAVH
Z239_HUMAN	1108	YKCEQCGKGFTRSSSLIHQAVH
O15322_HUMAN	1109	YKCEECGKGFTDSLHLKHQIIH
O15322_HUMAN	1110	YICEKCGRAFIHDLKLQKHQIIH
O14913_HUMAN	1111	YKCEKCGKGFFRSSDLQHHQKIH
O14913_HUMAN	1112	YKCEECGKCFSSFTSLKRHQIIH
O14913_HUMAN	1113	YPYKCEECGKGFSRSSKLQEHQTIH
ZN45_HUMAN	1114	YKGEHCVKSFSWSSHLQINQRAH
ZN45_HUMAN	1115	YKCEECGKGFSWSSSLIHHQRVH
ZN45_HUMAN	1116	YKCEECGKVFSWSSYLQAHQRVH
ZN45_HUMAN	1117	YKCEKCDNAFRRFSSLQAHQRVH
ZN45_HUMAN	1118	YKCERCCKAFSQQFSSLQVHQRVH
ZN45_HUMAN	1119	YKCEECGVGFSQRSYLQVHLKVH
ZN45_HUMAN	1120	YKCEECGKSFSWRSRLQAHERI
ZN45_HUMAN	1121	YKCEECGKGFSVGSHLQAHQISH
ZN45_HUMAN	1122	YQCAECGKGFSVGSQQLQAHQORCH
ZN45_HUMAN	1123	YQCEECGKGFCRASNFLAHRGVH
ZN45_HUMAN	1124	YKCEECGKGFCRASNLLDHQORGH
ZN45_HUMAN	1125	YKCEECGKGFSQASNLLAHQORGH
O75467_HUMAN	1126	FVCALCGAAFSQGSLSLFKHQRVH
ZN42_HUMAN	1127	YHCGECGLGFTQVSRLLTEHQRIH
O60765_HUMAN	1128	YRCNECGKGFTSISRLNRHRIIH
TYY1_HUMAN	1129	YVCPFDGCNKKFAQSTNLKSHILTH
O15391_HUMAN	1130	FVCPFDVCNRKFAQSTNLKTHILTH
TYY1_HUMAN	1131	FQCTFEGCGKRFSLDNLRTHVRIH
O15391_HUMAN	1132	FQCTFEGCGKRFSLDNLRTHLRIH
Q14872_HUMAN	1133	YQCTFEGCPRTYSTAGNLRTHQKTH
GLI1_HUMAN	1134	HKCTFEGCRKSYSRLENLKTHLRSH
GLI3_HUMAN	1135	HKCTFEGCTKAYSRLLENLKTHLRSH
O60255_HUMAN	1136	HKCTFEGCSKAYSRLLENLKTHLRSH
O60254_HUMAN	1137	HKCTFEGCSKAYSRLLENLKTHLRSH
O60253_HUMAN	1138	HKCTFEGCSKAYSRLLENLKTHLRSH
O60252_HUMAN	1139	HKCTFEGCSKAYSRLLENLKTHLRSH
GLI2_HUMAN	1140	HKCTFEGCSKAYSRLLENLKTHLRSH
O95409_HUMAN	1141	FQCEFEFGCDRRFANSSDRKKHMHVH
Q15915_HUMAN	1142	FKCEFEFGCDRRFANSSDRKKHMHVH
ZIC3_HUMAN	1143	FKCEFEFGCDRRFANSSDRKKHMHVH
GLI1_HUMAN	1144	YMCEHEGCSKAFSNASDRAKHQNRTH

O60255_HUMAN	1145	YVCEHEGCGNKAFSNASDRAKHQNRTH
O60254_HUMAN	1146	YVCEHEGCGNKAFSNASDRAKHQNRTH
O60253_HUMAN	1147	YVCEHEGCGNKAFSNASDRAKHQNRTH
O60252_HUMAN	1148	YVCEHEGCGNKAFSNASDRAKHQNRTH
GLI3_HUMAN	1149	YVCEHEGCGNKAFSNASDRAKHQNRTH
GLI2_HUMAN	1150	YVCEHEGCGNKAFSNASDRAKHQNRTH
Z143_HUMAN	1151	YVCTVPGCDKRFTTEYSSLYKHHVVH
TF3A_HUMAN	1152	FKCTQEGCGKHFASPSKLKRHAKAH
TF3A_HUMAN	1153	FVCDYEGCGKAFIRDYHLSRHILTH
Q14872_HUMAN	1154	FECDEVQGEKAFNTLYRLKAHQRLH
Q14872_HUMAN	1155	FVCNQEGCGKAFLTSHSLRIHVRVH
ZN76_HUMAN	1156	YRCDFPSCGKAFAATGYGLKSHVRTH
Z143_HUMAN	1157	YQCEHAGCGKAFATGYGLKSHVRTH
Q14872_HUMAN	1158	FRCDHDGCGKAFAASHHLKTHVRTH
O00153_HUMAN	1159	FICPAEGCGKSFYVLQRLKVHMRTH
ZN76_HUMAN	1160	FQCPFEGCGRSFTTTSNIRKVHVRTH
Z143_HUMAN	1161	FKCPFEGCGRSFTTTSNIRKVHVRTH
Q15915_HUMAN	1162	FPCPFPGCGKVFARSENLIKHKRTH
O95409_HUMAN	1163	FPCPFPGCGKVFARSENLIKHKRTH
ZIC3_HUMAN	1164	FPCPFPGCGKIFARSENLIKHKRTH
ZN76_HUMAN	1165	YTCPEPHCGRGFTSATNYKNHVRIH
Z143_HUMAN	1166	YYCTEPGCGRAFASATNYKNHVRIH
O00153_HUMAN	1167	FMCHESGCGKQFTTAGNLKNHRRIH
ZN76_HUMAN	1168	YKCPEELCSKAFKTSGLDLQKHVRTH
Z143_HUMAN	1169	YRCSEDNCTKSFKTSGLDLQKHIRTH
Q14872_HUMAN	1170	FNCSESEGCSKYFTTSLDLRKHIRTH
ZN76_HUMAN	1171	FRCGYKGCGRLYTTAHHLKVHERAH
Z143_HUMAN	1172	FRCEYDGCGLYTTAHHLKVHERSH
BTE1_HUMAN	1173	HKCPYSGCGKVYGKSSHLKAHYRVH
BTE2_HUMAN	1174	HYCDYPGCTKVYTKSSHLKAHLRTH
O43839_HUMAN	1175	HRCHFNGCRKVYTKSSHLKAHQORTH
UKLF_HUMAN	1176	HRCQFNGCRKVYTKSSHLKAHQORTH
O95600_HUMAN	1177	HQCDFAGCSKVYTKSSHLKAHRRIH
Q13118_HUMAN	1178	HICSHPGCGKTYFKSSHLKAHTRTH
O75411_HUMAN	1179	HICSHPGCGKTYFKSSHLKAHTRTH
EZF_HUMAN	1180	HTCDYAGCGKTYTKSSHLKAHLRTH
O14901_HUMAN	1181	YVCSFPGCRKTYFKSSHLKAHLRTH
SP4_HUMAN	1182	HICHIEGCGKVYGKTSHLRAHLRWH
O60402_HUMAN	1183	HICHIEGCGKVYGKTSHLRAHLRWH
EKLF_HUMAN	1184	HTCAHPGCGKSYTKSSHLKAHLRTH
WT1_HUMAN	1185	FMCAYPGCNKRYFKLSHLQMHSRKH
Q16256_HUMAN	1186	FMCAYPGCNKRYFKLSHLQMHSRKH
Q15881_HUMAN	1187	FMCAYPGCNKRYFKLSHLQMHSRKH
SP2_HUMAN	1188	HVCHIPDCGKTFRKTSLLRAHVRLH
O43167_HUMAN	1189	YACKDCHRKFMVDVSQLKKHLRTH
O75467_HUMAN	1190	YACRACSKVFKSSDLLKHLRTH
ZEP1_HUMAN	1191	YICEYCNRAKAPSVLLKKHIRSH
Q02646_HUMAN	1192	YICPYCSRACAKPSVLKKHIRSH

O75362_HUMAN	1193	YACSYCGKFFRSNYYLNIHLRTH
Q92981_HUMAN	1194	YKCVQPCGKAFVSRYKLMRHMATH
O76019_HUMAN	1195	YKCVQPCGKAFVSRYKLMRHMATH
RRE1_HUMAN	1196	YACSVCNKRFWSLQDLTRHMRSH
O75626_HUMAN	1197	HECQVCHKRFSSTSNLKTHLRLH
Z202_HUMAN	1198	HDCSVCGKSFTCNHLVRHLRTH
O75123_HUMAN	1199	YACDICGKTFTFNSDLVRHRISH
Z151_HUMAN	1200	HKCSVCSKAFVNVGDLKHHIIH
SNAI_HUMAN	1201	YACVCGTCGKAFSRPWLLQGHVRTH
O43623_HUMAN	1202	YACVCKICGKAFSRPWLLQGHIRTH
O95409_HUMAN	1203	HVCFWEECPREGKPFKAKYKLVNHIRVH
ZIC3_HUMAN	1204	HVCYWEECPREGKSFKAKYKLVNHIRVH
O00146_HUMAN	1205	HECKLCGASFRTKGSLLRHRRH
O00146_HUMAN	1206	HVCQFCSRGFREKGSLLRVHRH
IKAR_HUMAN	1207	FQCNQCGASFRTQKGNLLRHHKLH
CTCF_HUMAN	1208	HKCHLCGRAFRTVTLLRNHLNTH
HKR3_HUMAN	1209	HVCEFCSHAFTQKANLNMHLRTH
Q15552_HUMAN	1210	HVCEHCNAAFRTNYHLQRHVFIH
O43591_HUMAN	1211	HVCEHCNAAFRTNYHLQRHVFIH
PLZF_HUMAN	1212	YICSECNRTFPSHTALKRHLRSH
Z151_HUMAN	1213	YVCIHCQRQFADPGALQRHVRIH
MAZ_HUMAN	1214	YICALCAKEFKNGYNLRRHEAIIH
O14753_HUMAN	1215	HLCTGCGKGFNDTFDLKRHVTRTH
O95365_HUMAN	1216	YECNICKVRFTTRQDKLKVHMRKH
O15156_HUMAN	1217	YACEVCGVRFTTRNDKLKIHMRRKH
O75066_HUMAN	1218	YSCEECGAKFAANSTLKNHLRLH
O95365_HUMAN	1219	YLCQQCGAAFAHNYDLKNHMRVH
O15156_HUMAN	1220	YSCPHCPARFLHSYDLKNHMLH
Z151_HUMAN	1221	HKCEDCGKEFTHTGNFKRHIRIH
Z151_HUMAN	1222	YRCEDCGKLFTTSGNLKRHQLVH
Z151_HUMAN	1223	YKCRECGKQFTTSGNLKRHLRIH
O15090_HUMAN	1224	YDCPYCGKTFRTSHHLKVHLRIH

Example 3: Non-human zinc finger databases.

For providing novel combinations of non-antigenic, optimised zinc fingers, for use in species other than humans, separate species-specific zinc finger databases are required,
 5 such as mouse, chicken, pig, cow, *etc.*

The fingers listed below are in a format that can be linked with classical wild-type canonical "TGEKP" linkers (i.e. ...TGEKP – zinc finger peptide sequence – TGEKP – zinc finger peptide sequence – TGEKP - etc...). For each peptide sequence, an
 10 oligonucleotide is designed to encode the peptide sequence; the oligonucleotide can then be linked into a library selection system, as described in the Examples *infra*.

Mouse Zinc Finger Database.

15 544 zinc finger units

Name	SEQ ID NO	Peptide sequence
O35745_MOUSE	1225	HQCTHCEKTFNRKDHLKNHLQTH
ZFX2_MOUSE	1226	HRCEYCKKGFRRPSEKNQHIMRH
ZFX1_MOUSE	1227	HRCEYCKKGFRRPSEKNQHIMRH
ZFY2_MOUSE	1228	HKCDMCSKGFHRPSELKKHVATH
ZFY1_MOUSE	1229	HKCDMCSKGFHRPSELKKHVATH
ZFX2_MOUSE	1230	HKCDMCDKGFHRPSELKKHVAAH
ZFX1_MOUSE	1231	HKCDMCDKGFHRPSELKKHVAAH
ZFA_MOUSE	1232	HKCDMCDKGFHRPSELKKHVAAH
Q9Z162_MOUSE	1233	YTCSVCGKGF SRPDHLSCHVKHVVH
MAZ_MOUSE	1234	YNCSHCGKSFSRPDHLNSHVRQVH
Q08376_MOUSE	1235	YSCEVCGKSFI RAPDLKKHERVH
Z151_MOUSE	1236	HKCPHCDKKFNQVGNLKAHLKIH
ZFX2_MOUSE	1237	FRCKRCRKGFRQQSELKKHMKTH
ZFX1_MOUSE	1238	FRCKRCRKGFRQQSELKKHMKTH
Q62518_MOUSE	1239	YVCTMCGKGYTLNSNLQVHLRVH
Q60636_MOUSE	1240	YECNVCAKTFGQLSNLKVHLRVH
Q9Z117_MOUSE	1241	CSCPECCKVLHQLSHLRSHYRLH
Q61898_MOUSE	1242	CSCPECGREFHQLSHLRKHRYRLH
O88631_MOUSE	1243	YSCQYCGKVFHQLSHFKSHFTLH
Q61164_MOUSE	1244	HKCPDCDMAFVTSGELVRHRRYKH
O35483_MOUSE	1245	FRCADCGRGFAQRSNLAKHRRGH
O35483_MOUSE	1246	FVCGVCGAGFSRRAHLTAHGRAH
O70162_MOUSE	1247	FVCRDCGQGFVRSARLEEHRRVH
Q9Z1D8_MOUSE	1248	HRCGDCGKFFLQASNFIQHRRRIH
O35483_MOUSE	1249	HRCPCDCGKGFGHSSDFKRHRRTH
O35483_MOUSE	1250	---ADCGKS FVYGSHLARHRRTH

O35483_MOUSE	1251	FPCPDCGKRFVYKSHLVTHRRIH
O88282_MOUSE	1252	YKCQLCRSAFRYKGNLASHRTVH
Q61065_MOUSE	1253	YKCDRCQASFRYKGNLASHKTVH
BCL6_MOUSE	1254	YKCDRCQASFRYKGNLASHKTVH
O70162_MOUSE	1255	FACQDCGRRFNQSTKLIQHQRVH
O70162_MOUSE	1256	--CVECGERFGRRSVLLQHRRVH
Q9Z0G7_MOUSE	1257	-DCPVCNKKFKMKHHLTEHMKTH
Q08376_MOUSE	1258	---HMCDKAFKHKSHLKDHERRH
Q64318_MOUSE	1259	HECGICRKAFKHKHHLIEHMLRH
Q64318_MOUSE	1260	FKCTECGKAFKYKHHLKEHLRIH
Q9Z1D8_MOUSE	1261	FKCNECGKGFGRSHLAGHLRLH
Q9Z1D8_MOUSE	1262	YGCNECGKSFGRHSHLIEHLKRH
Q9Z2X6_MOUSE	1263	----YVCKQCGKAFTLSSSLRRH
KID1_MOUSE	1264	YVCKECGKAFTLSTSLYKHLRTH
Q9Z1D7_MOUSE	1265	HGCDECGKSFTQHSRLIEHKRVH
ZF90_MOUSE	1266	YRCNLCGRSFRHSTSLTQHEVTH
Q9Z2X6_MOUSE	1267	YVCKECGKAFASTSLHIHEGTH
Q9Z2X6_MOUSE	1268	YVCKHCGKAYTTYNTLRAHERSH
Q9Z2X6_MOUSE	1269	YVCKHCGKAYTTYNTLRAHERSH
Q9Z2X6_MOUSE	1270	YVCKHCGKAYTSYSTLRAHERSH
Q9Z2X6_MOUSE	1271	YVCKHCGKAYTSYSTLRAHERSH
Q9Z2X6_MOUSE	1272	YVCKHCGKAYTSYSTLRAHERSH
Q9Z2X6_MOUSE	1273	YVCKHCGKAFTQSSYLRIHKRTH
ZF37_MOUSE	1274	YECEQCGKAHGHKHALTDHLRIH
Q62514_MOUSE	1275	YECEQCGKAHGHKHALTDHLRIH
Q61491_MOUSE	1276	YECNQCCKAFTQFFPLKRHEITH
ZF37_MOUSE	1277	YKCDCEGKAFGHSSSLTYHMRTH
Q62514_MOUSE	1278	YKCDCEGKAFGHSSSLTYHMRTH
Q61491_MOUSE	1279	YQCNQCAKAFYPYHRTLQIHERTH
Q61491_MOUSE	1280	CEYNQCWKAFAFYHKTLOIHERTH
Q61491_MOUSE	1281	YECNQCCKAFACYQSFQIHKRTH
Q61491_MOUSE	1282	YECNQCCKAFACNRYLQIHKRTH
Q61491_MOUSE	1283	YECNQCCKAFACPRYLQIHKRTH
Q61491_MOUSE	1284	YECNQCCKAFACLRLNLQNHKTTH
Q61491_MOUSE	1285	FECNQCCKAFAAHSTLQRHKRTH
Q61491_MOUSE	1286	YECNQCCKAFTRHSTLQIHKRTH
Q61491_MOUSE	1287	YECNQCCKAFTCRSNLQIHKRTH
Q9Z2X6_MOUSE	1288	YVCKQCGKAFTRSSHLQIHKITH
Q9Z2X6_MOUSE	1289	YICKQCGKAFARSSHLQIHKRSH
Q61491_MOUSE	1290	YKCKQCGKDFTHHSTLHIHKRIH
Q9Z2X6_MOUSE	1291	YSCKLCGKAFTHSNYLQIHKRIH
Q61491_MOUSE	1292	YECNQCCKAFARNSNLLDHKRIH
Q64247_MOUSE	1293	YICKQCGKTFRYLSCFQKHERIH
Q9Z2X6_MOUSE	1294	YACKQCDKAFKYLSSLQNHKRIH
Q9Z2X6_MOUSE	1295	HACKQCGKSFKRQSNVQAHERNH
Q64247_MOUSE	1296	YTCKHCTKTFTTSSTRNSHEKTH
Q64247_MOUSE	1297	YACKHCGKAFTTSSARNSHERI
Q64247_MOUSE	1298	YACKHCGKAFTSSSDRNSHERIH

Q64247_MOUSE	1299	YPCKYCGKAFATSSDRNSHERIH
Q64247_MOUSE	1300	YSCTHCGKAFSSPSDYNSCERIH
O88412_MOUSE	1301	YVCNECGKAFTCSSYLLIHQRIH
ZF35_MOUSE	1302	YMCNHICYKHFSQSSDLIKHQRIH
Q9Z2X6_MOUSE	1303	YVCKQCGKAFAQSSYLHIHQRS
ZF38_MOUSE	1304	YQCKDCGKAFFSGKGLIRHYRIH
OZF_MOUSE	1305	YECNKCAGKAFSRITSLIVHVRIH
Q9Z0Q5_MOUSE	1306	YECNECGKAFSQRTSLIVHVRIH
ZF90_MOUSE	1307	YQCNVCGKAFFKRSTSFIEHHRIH
OZF_MOUSE	1308	YECKICGKAFCQSSSLTVHMRSH
Q9Z0Q5_MOUSE	1309	YECNVCGKAFSQSSSLTVHVRSH
ZF90_MOUSE	1310	YECIDCGKAFSQSSSLIQHERTH
Z151_MOUSE	1311	CQCVICGKAFTQASSLIAHVQRH
OZF_MOUSE	1312	YECKGCGKAFFIQKSSLIRHQRS
Q9Z0Q5_MOUSE	1313	FECKDCGKAFFIQKSNLIRHQRT
Q9Z162_MOUSE	1314	---TYCSKAFRDSYHLRRHQSCH
Q9Z162_MOUSE	1315	HACEMCGKAFRDVYHLNRHKL
MAZ_MOUSE	1316	HACEMCGKAFRDVYHLNRHKL
Q61898_MOUSE	1317	FRCTECDKSFIRSSHLREHQKI
Q60585_MOUSE	1318	FDCKECGKTFSRGYHLTLHQRI
O35483_MOUSE	1319	YACAECGRRFGQSAALTRHQWA
Q60585_MOUSE	1320	YACTECGKSFRQVAHLTRHQRL
Q9Z1D9_MOUSE	1321	YACPECGECFRQSSHLSRHQRT
Q9Z1D9_MOUSE	1322	YKCFQCGERFRQSTHLVRHQRI
O88631_MOUSE	1323	YKCTKCDKLFTQYSHLRRHQRI
Q60585_MOUSE	1324	YKCTECKKAFRQHSHLTYHQRI
MLZ4_MOUSE	1325	HKCTECAKASAASPHLIQHQR
Q9Z116_MOUSE	1326	YECTECSKAFCQKSHLTQHQRV
O70237_MOUSE	1327	YPCQFCGKRFBHQKSDMKKHTY
GFI1_MOUSE	1328	YPCQYCGKRFBHQKSDMKKHTF
Q61624_MOUSE	1329	FRCDECGMRFIQKYHMERHKR
P97475_MOUSE	1330	FRCDECGMRFIQKYHMERHKR
Q61624_MOUSE	1331	FQCSQCDMRFIQKYLLQRHEKI
P97475_MOUSE	1332	FQCSQCDMRFIQKYLLQRHEKI
ZFP1_MOUSE	1333	FVCNYCDKTF SFKSLLVSHKRI
Q9Z116_MOUSE	1334	YICFECKRKAFFYRKSELTDHQ
Q9Z116_MOUSE	1335	YECKECCGKAFCQKPQLTLHQRI
ZFP1_MOUSE	1336	YGCSECGKTFAQKFELTTHQRI
Q06054_MOUSE	1337	YKCSDCGKCFIQKANLRTHQKI
Q06054_MOUSE	1338	YKCSDCGKCFIQKANLRTHERI
Q06054_MOUSE	1339	YKCSDCDKCFIQKAKLKKHQRI
Q06054_MOUSE	1340	YKCSDCDKCFIQKDHLRTHQRL
Q06054_MOUSE	1341	YKCSDCDKCFIRKANLRRHHR
Q06054_MOUSE	1342	YKCSCHCKCFIRKAHLRRHQRI
Q06054_MOUSE	1343	YKCSCHCKCFIQQAHLRRHQKI
Q06054_MOUSE	1344	YICAECNKCFFIQKSQLKTHQRI
MLZ4_MOUSE	1345	HICSQCGKAFSQISDLNRHQKTH
ZF37_MOUSE	1346	YECNECGIAFSQKSHLVVHQRT

Q62514_MOUSE	1347	YECNECGIAFSQKSHLVLHQORTH
ZF37_MOUSE	1348	YECVECGKAFSQKSHLIVHQRP
Q62514_MOUSE	1349	YECVECGKAFSQKSHLIVHQORTH
ZF37_MOUSE	1350	FECNECGKTFSKKSHLVIHQORTH
Q62514_MOUSE	1351	FECNECGKTFSKKSHLVIHQORTH
MFG3_MOUSE	1352	FECKECGKAFHFSSQLNNHKTS
Q62514_MOUSE	1353	FECYECGKAFAKSQLVIHQORSH
ZF37_MOUSE	1354	FECYECGKAFAKSQLVIHQORSH
Q9Z116_MOUSE	1355	YECKICGKCFYWKTSEFNRHQSTH
O88412_MOUSE	1356	YSCNECGKAFRQKSSLTVHQORTH
Q9Z116_MOUSE	1357	YECACGKAFASTKSYLTVHQORTH
P70405_MOUSE	1358	YECCKGKTFRGKYSLDQHQORVH
ZF90_MOUSE	1359	HECADCGKTFLWRTQLTEHQORIH
KR2_MOUSE	1360	YECMICGKHFTGRSSLTVHQVH
KR2_MOUSE	1361	YECDCGKAFAKNSSLIVHQORIH
Q9Z1D7_MOUSE	1362	YKCSVCGKAFAKISLIEHEQIH
Q61116_MOUSE	1363	YKCDTCGKAFAFSQKSSLQVHQORIH
O70237_MOUSE	1364	--CRMCGKAFAKRSSTLSTHLLIH
GFI1_MOUSE	1365	-DCKICGKSFAKRSSTLSTHLLIH
Q9Z150_MOUSE	1366	HSCGICGKCFTQKSTLHDHLNLH
Q9Z1D7_MOUSE	1367	YKCEVCGKTFRWRTVLIRHKVVH
ZF35_MOUSE	1368	-YKCMCGKAFAFSQCSAFTLHQORIH
ZF38_MOUSE	1369	YKCKECGKAFAHSSNFNKHHRH
OZF_MOUSE	1370	YGCNECGKAFAFSQFSTLALHMRH
Q9Z0Q5_MOUSE	1371	YGCNECGKAFAFSQFSTLALHLRH
ZFP1_MOUSE	1372	YECTECGKTFSQRSTLRLHLRH
MLZ4_MOUSE	1373	YKCDECGKNFSQNSDLVRHRRH
Q62514_MOUSE	1374	YECNECGKAFAKYGSSLTKHMRH
ZF37_MOUSE	1375	YECNECGKAFAKYGSSLTKHMRH
KR2_MOUSE	1376	YKCHDCGKAFAKNSSLTQHRRH
P70405_MOUSE	1377	CRDCGKFFSQTSHLNDHRRH
Q61117_MOUSE	1378	YKCSTCGKGFSRSSLNVHCRH
ZF92_MOUSE	1379	YLCQQCGKSFSRSEFNLIKHRH
ZF29_MOUSE	1380	YACKECGESFSYNSNLIRHQORH
O88282_MOUSE	1381	YRCSICGARFNRPANLKTHSRH
Q61065_MOUSE	1382	YRCNICGAQFNRPANLKTHTRH
BCL6_MOUSE	1383	YRCNICGAQFNRPANLKTHTRH
ZF29_MOUSE	1384	YKCRDCGKSFSRSANLITHQORH
Q9Z1D7_MOUSE	1385	YQCLQCNKSFNRRSTLSQHQGVH
ZF35_MOUSE	1386	YPCNSCSKSFSRGSDLIKHQORH
ZF35_MOUSE	1387	YPCSWCIKSFSRSSLIKHQORH
ZF35_MOUSE	1388	YPCNQCTKSFSRSLDLINHQORH
ZFP1_MOUSE	1389	YECDCVCQKTFSHKANLIKHQORH
ZF35_MOUSE	1390	YECCKGKTFSQSSNLILHQORH
O88412_MOUSE	1391	YECNECGKTFTRSSNLIVHQORH
MLZ4_MOUSE	1392	YDCNECGKSFGRSSHLIQHQTIH
MLZ4_MOUSE	1393	YECTACGKSFSRSSHLITHQKIH
KR2_MOUSE	1394	YECTECGKAFAFSQSAYLIEHRRH

ZF90_MOUSE	1395	YACKECGRNFSRSSALTKHHRVH
MLZ4_MOUSE	1396	YECTECDKSFSRSSALIKHKRVH
P70405_MOUSE	1397	YKCSECGKSFSQSSILIQHRRIH
P70405_MOUSE	1398	YKCSECGNSFSQSAILNQHRRIH
Q9Z1D8_MOUSE	1399	HQCNECGKSFIQSAHLIQHRRIH
KID1_MOUSE	1400	YRCQECGMSFGQSSALIQHRRIH
P70405_MOUSE	1401	YECSQCGKSFSQKSGLIQHVVH
P70405_MOUSE	1402	YECRECGKSFSQKATLIKHQRVH
P70405_MOUSE	1403	YECSQCGKSFSQKATLVKHKRVH
Q9Z1D8_MOUSE	1404	HQCNECGRGFSLKSHLSQHQRH
OZF_MOUSE	1405	YQCSECGKAFSQKSHHIRHQRIH
Q9Z0Q5_MOUSE	1406	YQCSECGKAFSQKSHHIRHQKIH
O88412_MOUSE	1407	YDCSECGKAFSQLSCLIVHQRIH
ZF35_MOUSE	1408	YKCSECGKAFNQSSVLILHQRIH
ZF35_MOUSE	1409	YKCDVCGKAFSQSSDRILHQRIH
KID1_MOUSE	1410	FKCNTCGKTFRQSSSRIAHQRIH
OZF_MOUSE	1411	YKCNECGTIFRQKQYLIKHHNIH
Q9Z0Q5_MOUSE	1412	FKCNECGTAFGQKKYLIKQHNIH
OZF_MOUSE	1413	FECSCGGRAFSQKQYLIKQHNIH
Q9Z0Q5_MOUSE	1414	FECNECGKAFSQKQYVIKHQSTH
OZF_MOUSE	1415	FKCNECGKAFSQKENLI IHQRIH
Q9Z0Q5_MOUSE	1416	FECSDCGKAFSQKENLLTHQKIH
KID1_MOUSE	1417	FKCSECGRAFSQSASLIQHERIH
O88412_MOUSE	1418	FECHECGKAFIQSANLVVHQRIH
O88412_MOUSE	1419	FTCSECGKGFSQSANLVVHQRIH
O88412_MOUSE	1420	FACSDCGKAFTQSANLIVHQRS
KR2_MOUSE	1421	YKCHECGKAFSQSMNLTVHQRT
ZF38_MOUSE	1422	YQCNECGKSFSQHAGLSSHQRLH
KID1_MOUSE	1423	YNCNECGKALSSHSTLI IHERIH
O35700_MOUSE	1424	YKCDQCPKAFNWKSNLIRHQMSH
EVI1_MOUSE	1425	YKCDQCPKAFNWKSNLIRHQMSH
Q62518_MOUSE	1426	YKCDVCGKSFGWRSNLI IHHRH
Q9Z1D8_MOUSE	1427	YACHLCGKAFRVRSHLVQHQS
Q9Z1D8_MOUSE	1428	YKCQVCGKAFRVSSHLLVQHHS
Q9Z1D7_MOUSE	1429	YECNDCGKAFFVYNSSLATHQ
MFG3_MOUSE	1430	YKCNACGRAFNRRSNLMQHEK
MFG3_MOUSE	1431	YKCNVCGKAFNRRSNLLQHQK
O88412_MOUSE	1432	YVCGKCGKAFTQSSNLTVHQK
Q9Z116_MOUSE	1433	YECKECCRKAFFYDKSNLKRHQ
Q60585_MOUSE	1434	YECKECCRKFFRRYSELISHQ
Q60585_MOUSE	1435	YECKECGKAFRQCAHLRHRHQ
ZF37_MOUSE	1436	YECIECGKAFKQNASLTKHMK
Q62514_MOUSE	1437	YECIECGKAFKQNASLTKHMK
Q61849_MOUSE	1438	YECNECGKAFKRHRSFVRHQK
MFG3_MOUSE	1439	FECKDCGKVFRNLNIHLIRHQ
Q61849_MOUSE	1440	YECKECGKAFLPQQQLTRHQK
Q06054_MOUSE	1441	HRCNECGKSLSSSSGLQRHQ
O35700_MOUSE	1442	HACPECGKTFATSSGLKQHKH

EVI1_MOUSE	1443	HACPECGKTFATSSGLKQHKHIIH
ZF92_MOUSE	1444	YECGECGKTFTRSSNLVKHQVH
O88412_MOUSE	1445	FKCSECEKAFSYSSQLARHQKVH
ZF90_MOUSE	1446	FECNVCCKAFRHSSSLGQHENA
KID1_MOUSE	1447	YECNTCGKLFNHRSSLTNHYKIIH
ZF29_MOUSE	1448	YKCDECGKSFSDGSNFSRHQTTH
OZF_MOUSE	1449	YKCGECGKAFSQRGNFSLHQKQH
O70162_MOUSE	1450	CDVCGKVFSQRSNLLRHQKIHTG
ZFP1_MOUSE	1451	YECNECAKTFFKKSNLIIHQKIIH
O88412_MOUSE	1452	YKCKDCEKAFSCFSLHIVHQRIH
Q9Z1D7_MOUSE	1453	YKCNECGRAFGQWSALNQHQRLH
ZF90_MOUSE	1454	YQCSLCGKAFQRSSSLVQHQRH
Q64247_MOUSE	1455	-----CGKVFILSGDLIKHERIH
MFG3_MOUSE	1456	YECEQCGSAFRLPYQLTQHQRH
Q61849_MOUSE	1457	FECCLCGSAFRCRSQNLNHLRIH
MFG3_MOUSE	1458	FKCKLCESAFRRKYQLSEHQRIH
Q61849_MOUSE	1459	FKCQECGKAFVVLAYLIEHQSIH
Q64247_MOUSE	1460	FVCKQCGEAFVNSSHLISHERIH
MFG3_MOUSE	1461	FQCKECCGRAFVRSTGLRIHERIH
Q64247_MOUSE	1462	FVCKTCGKAFSRSDYLINHKRIH
Q64247_MOUSE	1463	FVCKKCGKAFKRLGHFMNHERIH
ZF90_MOUSE	1464	FQCKECCGKAFSRCSSLVQHHERTH
MFG3_MOUSE	1465	FECKDCGKAFTVLAQLTRHQTIIH
MFG3_MOUSE	1466	FHCKVCGKAFTVLAQLTRHENIIH
MFG3_MOUSE	1467	FECCKECCGKSFKRVSSLVEHRIIH
ZFP1_MOUSE	1468	FECPECCGKAFTHQSNLIVHQRAH
ZF92_MOUSE	1469	FECTECCGKAFSRSSNLIEHQRIH
O54978_MOUSE	1470	FECQECGEAFARRSELIEHQKIIH
O70162_MOUSE	1471	FRCTECCGQSFRQRSNLLQHQRH
O70162_MOUSE	1472	FACAECGQSFRQRSNLTQHQRH
O70162_MOUSE	1473	FACPECCGQSFRQHANLTQHRRH
O70162_MOUSE	1474	YACAECGKAFRQRPTLTQHHLRTH
O70162_MOUSE	1475	AECGKTFRQRATLTQHLCVHTGE
Q9Z1D8_MOUSE	1476	FRCEECCGKSYNQRVHLIQHHRVH
Q9Z1D8_MOUSE	1477	FKCGECCGKSYNQRVHLTQHQRVH
ZF37_MOUSE	1478	FECNQCCGKAFKQIEGLTQHQRVH
Q62514_MOUSE	1479	FECNQCCGKAFKQIEGLTQHQRVH
O88282_MOUSE	1480	YPCPTCCGTRFRHLQTLKSHVRIH
Q61065_MOUSE	1481	YPCEICGTRFRHLQTLKSHLRIH
BCL6_MOUSE	1482	YPCEICGTRFRHLQTLKSHLRIH
Q60585_MOUSE	1483	YDCKECCGKAFRVRQQLTLHERIH
Q60585_MOUSE	1484	YDCKECCGKAFRVRGQLMLHQRIH
Q60585_MOUSE	1485	YECGECGKAFKVRQQLTFHQRIH
OZF_MOUSE	1486	YACKECCGKAFFNGKSYLKEHEKIIH
OZF_MOUSE	1487	YTCKECCGKAFFSGKSNLTEHEKIIH
Q9Z0Q5_MOUSE	1488	FICKECCGKTFSGKSNLTEHEKIIH
MFG3_MOUSE	1489	YKCKDCGKCFGCKSNLHQHESI
Q61849_MOUSE	1490	YQCKECCGKCFRQRSKLTEHESI

Q61849_MOUSE	1491	YECKECCGKCFGCRSTLTQHQS VH
Q61849_MOUSE	1492	FECCECGKKFRTARHLVKHQRIH
ZF92_MOUSE	1493	FVCRMCGKVFRRSFALLEHTRI H
ZF92_MOUSE	1494	YECSECGKQFQRSLALLEHQRIH
ZF35_MOUSE	1495	YECCECGKA FRMSSALVLHQRIH
P70405_MOUSE	1496	YECSECGKLFRQNSSLVDHQKTH
REX1_MOUSE	1497	HVCAECGKAFTESSKLKRHFLVH
TTY1_MOUSE	1498	HVCAECGKA FVESSKLKRHQLVH
ZFX2_MOUSE	1499	HICVECGKGFRHPSELKKHMR IH
ZFX1_MOUSE	1500	HICVECGKGFRHPSELKKHMR IH
ZFA_MOUSE	1501	HICVECGKG FCHPSELKKHMR IH
ZFY2_MOUSE	1502	HICGECGKGFRHPSALKKKHIRV H
ZFY1_MOUSE	1503	FICGECGKGFRHPSALKKKHIRV H
Q61116_MOUSE	1504	--CHECGKGFRQSSALQTHQRVH
Q06054_MOUSE	1505	YQCRKCGKCFRTYSSLYRHRTH
Q9Z117_MOUSE	1506	HQCEKCRKCFSTASSLTVHKRIH
Q61898_MOUSE	1507	HQCGKCGKCFNTSSSLTVHHR IH
Q60585_MOUSE	1508	YDCKECCGKA FRLFSQLTQHQS IH
Q60585_MOUSE	1509	YKCMCEKTFRLLSQLTQHQS IH
Q60585_MOUSE	1510	YDCKECCGKA FRLHSSLIQHQR IH
KR2_MOUSE	1511	YQCKECCGKA FRKNSSLIQH ER IH
KID1_MOUSE	1512	YLCNECGNTFKSSSSSLRYHQRI H
KR2_MOUSE	1513	YGCDECGKTFRQSSSL LKHQR IH
ZF37_MOUSE	1514	YKCNECGKTFRHSSNLMQH LRS H
Q62514_MOUSE	1515	YKCNECGKTFRHSSNLMQH LRS H
KID1_MOUSE	1516	YKCNECGKTFRCNSSLSNHQR TH
ZF37_MOUSE	1517	YECKECCGKSFRYNSSLTEHV RTH
Q62514_MOUSE	1518	YECKECCGKSFRYNSSLTEHV RTH
Q9Z117_MOUSE	1519	YKCKECCGKS FLELSHLKRHYR IH
O88631_MOUSE	1520	HKCKECCGKS FFILSHLKT HYR IH
Q61898_MOUSE	1521	YECKECCGKS FIELSHLKRHYR IH
Q9Z1D7_MOUSE	1522	HGCDECGKSFTQHSRL IEHKRVH
O35738_MOUSE	1523	FKCADCDRRFSRSDHLALHRR RH
O89090_MOUSE	1524	--CPEC PKRFMRSDHLSKH I KTH
Q64167_MOUSE	1525	--CPEC PKRFMRSDHLSKH I KTH
O89087_MOUSE	1526	--CPEC PKRFMRSDHLSKH I KTH
Q62445_MOUSE	1527	--CPEC SKRFMRSDHLSKH V KTH
O89091_MOUSE	1528	--CPMCDRRFMRSDHLTKHARR H
Q61596_MOUSE	1529	--CPMCDRRFMRSDHLTKHARR H
BTE1_MOUSE	1530	--CPLCEKRFMRSDHLTKHARR H
Q62445_MOUSE	1531	FICNWMFCGKRFTRSDELQRHRR TH
Q64167_MOUSE	1532	FMCNWSYCGKRFTRSDELQRH KR TH
O89090_MOUSE	1533	FMCNWSYCGKRFTRSDELQRH KR TH
O89087_MOUSE	1534	FMCNWSYCGKRFTRSDELQRH KR TH
Q60843_MOUSE	1535	YHCNWE GCGWK FARSDELTRHYR KH
EZF_MOUSE	1536	YHCDWDGCGWK FARSDELTRHYR KH
Q60980_MOUSE	1537	YKCTWEGCTWK FARSDELTRHFR KH
O35738_MOUSE	1538	YKCTWEGCTWK FGRSDELTRHYR KH

Q9Z0Z7_MOUSE	1539	YKCTWEGCDWRFARSDELTRHYRKH
O70261_MOUSE	1540	YACSWDGCDFWRFARSDELTRHYRKH
EKLF_MOUSE	1541	YACSWDGCDFWRFARSDELTRHYRKH
Q61596_MOUSE	1542	FSCSWKGCERRFARSDELSTRHRRTH
O89091_MOUSE	1543	FSCSWKGCERRFARSDELSTRHRRTH
BTE1_MOUSE	1544	FPCTWPDCLKKFSRSDDELTRHYRTH
EGR2_MOUSE	1545	YPCPAEGCDRRFSRSDDELTRHIRIH
WT1_MOUSE	1546	YQCDFKDCERRFSRSDQLKRHQRRH
WT1_MOUSE	1547	FQCKTCQRKFSRSDHLKTHTRTH
EGR1_MOUSE	1548	FQCRICMRNFSRSDHLTTHIRTH
KR2_MOUSE	1549	YQCNECGKPFSTRSTNLTRHQORTH
O35700_MOUSE	1550	YTCRYCGKIFPRSANLTRHLRTH
EVI1_MOUSE	1551	YTCRYCGKIFPRSANLTRHLRTH
ZF29_MOUSE	1552	FQCAECGKSFSRSPNLIAHQORTH
ZF38_MOUSE	1553	YVCTKCGKAFSHSSNLTLHYRTH
Q9Z1D8_MOUSE	1554	YQCDSGKAFSYSSDLIQHYRTH
ZF29_MOUSE	1555	YQCGECGKNFSRSSNLATHRRTH
ZF29_MOUSE	1556	YRCPECCKGFSNSSNFITHQORTH
ZF38_MOUSE	1557	YICAECKGAFSNSSNLTKHRRTH
ZF29_MOUSE	1558	YECLTCGESFSWSSNLIKHQORTH
ZF90_MOUSE	1559	YECNECGEAFSRLSSLTQHERTH
MLZ4_MOUSE	1560	YHCNECGENFSRISHLVQHQORTH
ZF29_MOUSE	1561	YKCLMCKGKSFSRGSILVMHQRAH
MLZ4_MOUSE	1562	YECNECGKSFSSRSSHQAQHQORTH
MLZ4_MOUSE	1563	YKCYECCKGFSRSSHLLIQHQORTH
O70162_MOUSE	1564	FACPECCKQRFSSQRLKLTRHQORTH
O35483_MOUSE	1565	FPCPECCKRFSQRSVLVTHQORTH
O35483_MOUSE	1566	--CDECKGKFVYRSHLAHQORTH
ZFP1_MOUSE	1567	YECSECKGSFIQNSQLIHRRTTH
GFI1_MOUSE	1568	HKCQVCKGKAFSSQSSNLITHSRKH
O70237_MOUSE	1569	HKCQVCKGKAFSSQSSNLITHSRKH
ZF29_MOUSE	1570	YKCTECCKQKFSQSSALITHRRTH
KID1_MOUSE	1571	FKCKECKSKAFSSQSSALIQHQITH
KID1_MOUSE	1572	CKCKVCKGKAFRQSSALIQHQRMH
Z151_MOUSE	1573	YVCERCCKRQFVQSSQLANHIRHH
O35700_MOUSE	1574	YECENCAKVFTDPSNLQRHIRSQH
EVI1_MOUSE	1575	YECENCAKVFTDPSNLQRHIRSQH
Q60585_MOUSE	1576	YECKKCAKIFTCSSDLRGHQQRSH
Q9Z116_MOUSE	1577	YECTVCRKSFICKSSFSHHWRTH
KR2_MOUSE	1578	YTCNVCDKHFIERSSSLTVHQORTH
Q61164_MOUSE	1579	FQCSLCSYASRDYKLRHMRTH
P97365_MOUSE	1580	FQCWLCSAKFKISSDLKRHMVRH
KID1_MOUSE	1581	YKCSMCCKTFINTSSLRKHEKNH
ZF35_MOUSE	1582	YTCNLCSKSFSSQSSDLTKHQVRH
ZF35_MOUSE	1583	YHCSSCNKAFRQSSDLILHHRVRH
ZF38_MOUSE	1584	YWCSHCKGKTFCKSKSNLSKHQVRH
Q9Z1D9_MOUSE	1585	YKCGDCEKSFQRSDLFKHQORTH
Q9Z1D9_MOUSE	1586	YKCDSCCKGFRQRSDLFKHQRIH

ZF35_MOUSE	1587	YPCSQCSKMF'SRRSDLVKHYRIH
ZF35_MOUSE	1588	YQCSHCSKSFSQHS GMVKHLRIH
ZF35_MOUSE	1589	YACTQCPRSFSQKSDLIKHQRIH
ZF35_MOUSE	1590	YPCAQC NKSF'SQNSDLIKHRRIH
ZF35_MOUSE	1591	YMCNH CYKHFSQSSDLIKHQRIH
ZF35_MOUSE	1592	YNCDECDQSFAWSTGLIRHQORTH
Q9Z1D9_MOUSE	1593	YQCQECGKRFSQSAALVKHQORTH
Q9Z1D9_MOUSE	1594	YACVVCGRRFSQSATLIKHQORTH
Q9Z116_MOUSE	1595	YECKQCMKTFYRKSGLTRHQORTH
Q06054_MOUSE	1596	YECKQCSKSFYTSSHLENHYRTH
Q9Z116_MOUSE	1597	YECQLCQKAFYCTSHLIVHQORTH
ZF29_MOUSE	1598	YECPCGKTF'SRKSHLITHERTH
MLZ4_MOUSE	1599	YECVQCGKGFTQSSNLITHQRVH
ZF37_MOUSE	1600	YECNHCGKVL SHKQGLLDHQORTH
Q62514_MOUSE	1601	YECNHCGKVL SHKQGLLDHQORTH
ZF90_MOUSE	1602	YECNECGRAFRKKTNLHDHQORTH
Q61491_MOUSE	1603	YECNQCGRAFRQYVYLQCHERIH
ZF35_MOUSE	1604	YPCAQCGKSFSQSRSDLVNHQRVH
Q64247_MOUSE	1605	YVCEQCGKGFIQLKYLLMHQORSH
Q61116_MOUSE	1606	YTCQQCGKGFSQASYFHHMHQRVH
O35483_MOUSE	1607	YRCVFCGAGFGRRSYCVTHQORTH
ZF29_MOUSE	1608	YRCGDCGKGFSQRSQ LVVHQORTH
Q61117_MOUSE	1609	YRCDICGKRFRQRSYLHDHHR IH
Q9Z2U2_MOUSE	1610	FKCVVP SCKTFTRNSNLRAHCQLVH
Q61116_MOUSE	1611	YRCDSCGKGFSRSSDLNIHRRVH
Q61117_MOUSE	1612	YQCHACWKS FCHSSEFNNHIRVH
Z239_MOUSE	1613	YQCYECGKGFSQSSDLRIHLRVH
Z239_MOUSE	1614	FKCDRCGKGFSQSSKLHIHKRVH
Z239_MOUSE	1615	YHCGKCGQGFSQSSKLLIHQRVH
Z239_MOUSE	1616	YKCGECGKGFSQSSNLHIHRCTH
ZF35_MOUSE	1617	YKCDECGKAFSQSSDLMIHQRIH
ZF38_MOUSE	1618	YDCKCGKA FGQSSDLLKHQORMH
O35700_MOUSE	1619	YRCKYCDRSFSISSNLQRHV RN IH
EV11_MOUSE	1620	YRCKYCDRSFSISSNLQRHV RN IH
O35483_MOUSE	1621	YRCVFCGRSFSQSSALARHQAVH
O35483_MOUSE	1622	YLC SNCGRRFSQSSHLLTHMKTH
O70162_MOUSE	1623	FVCGECGRSFSRSSHLLRHQLTH
O88412_MOUSE	1624	YECAKCGAAFISNSHLMRHHRTH
O88631_MOUSE	1625	YKCMECDRSYIQYSHLKRHQKVH
O88631_MOUSE	1626	YKCKECGKS YAYRTGLKRHQKIH
Z239_MOUSE	1627	YEC SKCGKGFSQSSNLHIHQRVH
Z239_MOUSE	1628	YACEECGMSFSQRSNLHIHQRVH
MLZ4_MOUSE	1629	YECNECWRSFGERSDLIKHQORTH
MLZ4_MOUSE	1630	YECHECGRGFSERSDLIKHYRVH
Q61116_MOUSE	1631	YECNECGKRFSLSGNLDIHQRVH
Q61116_MOUSE	1632	YKCGDCGKRFS CSSNLHTHQRVH
Q62518_MOUSE	1633	YKCGECGKS FICSSNLYIHQRVH
Q9Z150_MOUSE	1634	CPRCGKQFNHSSNLNRHMNVHRG

Q61116_MOUSE	1635	FHCSVCGKNFSRSSHFLDHQRIH
Q61116_MOUSE	1636	KCNVCQKQFSKTSNLQAHQRVH
Q62518_MOUSE	1637	YSCDVCGKGFSRSSQLQSHQRVH
Q62518_MOUSE	1638	FKCDACGKSFSRSSHLRSHQRVH
Q61898_MOUSE	1639	YKCRECDKSFTQRAYLRNHHNRVH
Q61898_MOUSE	1640	YKCMECDKSFTHNSNFRTHQRVH
Q9Z117_MOUSE	1641	YKCMECNKSFTQDSHLRTHQRVH
Q61898_MOUSE	1642	YKCIECDKSFTQVSHLRTHQRVH
O88631_MOUSE	1643	YKCSECDKSFTQASQLRTHQRVH
Q61898_MOUSE	1644	YKCNECDRSFTHYASLRWHQKTH
Q9Z117_MOUSE	1645	YKCKECDKSFAHCSSFRRHQKTH
Q61898_MOUSE	1646	YKCKECDKSFAHYPNFRTHQKIH
O88631_MOUSE	1647	YKCKDCDIFFNHYSSLRRHQKVH
Q9Z117_MOUSE	1648	YKCKDCDISFIQISNLRRHQRVH
Q61898_MOUSE	1649	YKCRDCDISFSQISNLRRHQKLH
Q9Z117_MOUSE	1650	FKCRECDKSFTKCSHLRRHQSVH
Q61898_MOUSE	1651	YKCRECDKSFIHSSHLRRHQNVH
Q9Z117_MOUSE	1652	YKCRECDKSFIQRSNLI IHQRVH
Q06054_MOUSE	1653	YKCSECEKSFTCGSVLRKHQKIH
Q06054_MOUSE	1654	YKCSECEKSFTVGSDLRMHQKIH
Q06054_MOUSE	1655	YKCSECEKCFTVVSDLRTHQKIH
Q06054_MOUSE	1656	YKCSECEKSFTVGSSLRIHQRIH
Q06054_MOUSE	1657	YKCECGKSFTVGSDLRKHQKCH
Q61898_MOUSE	1658	YKCIECGKSFTNNSYLRTHQKVH
Q61898_MOUSE	1659	YRCKECDKSFHESATLREHEKSH
Q61898_MOUSE	1660	YRCAECDKSFTTRCSYLRAHQKIH
Q9Z117_MOUSE	1661	YRCKECDKSFTTECSTLRAHQKIH
Q61898_MOUSE	1662	YRCKECDKSFTSCSTLKAHQSIH
Q9Z117_MOUSE	1663	YICKECGKSFTTRCSYLRAHQKIH
O88631_MOUSE	1664	YVCKECGKSLTTCAILRAHQKIH
Q61898_MOUSE	1665	YECKECGKSFTTCSTLRIHQTIH
Q9Z117_MOUSE	1666	YICKECGKSFTKCSLQIHQKIH
O88631_MOUSE	1667	YTCKQCGKSFTTRGSTLRVHQRIH
O88631_MOUSE	1668	YKCNICDKSFTTECSSLKEHRKTH
Q9Z117_MOUSE	1669	YKCEVCDKSFTVNSTLKTHLKIH
Q61898_MOUSE	1670	YKCEICDKSFTTTTTLKTTHQKIH
Q9Z117_MOUSE	1671	YKCSVCGKSFTQCTNLKTHQRLH
Q61898_MOUSE	1672	YKCSVCDKSFTQCTHLKIHQRRH
KID1_MOUSE	1673	YRCKECGKSFGRRSGLFIHQKVH
ZF29_MOUSE	1674	YSCPECGKSFGNRSSLNTHQGIH
Q9Z117_MOUSE	1675	YKCKECGKSFPQLSALKSHQKIH
Q61898_MOUSE	1676	YKCKECEKSFFVQLSALKSHQKLH
O88631_MOUSE	1677	YKCNDCGKSFSYLSALQSHHKRH
Q08376_MOUSE	1678	FVCEMCTKGFTTQAHLEHLKIH
Q60636_MOUSE	1679	FKCQTCNKGFTQLAHLQKHVLVH
Q61116_MOUSE	1680	YKCEVCGKGFTQWAHLQAHHERIH
O88282_MOUSE	1681	YKCETCGSRFVQVAHLRAHVLIH
Q61065_MOUSE	1682	YKCETCGARFVQVAHLRAHVLIH

BCL6_MOUSE	1683	YKCETCGARFVQVAHLRAHVLIH
O88631_MOUSE	1684	YRCEVCDKWFTLSSSLSRHQKIH
Q61116_MOUSE	1685	YRCEVCGKRFPWSLSLHSHQSVH
Z239_MOUSE	1686	YKCDKCGKGFTRSSSLLVHHS LH
ZF29_MOUSE	1687	YKCGLCGKSFSQSSSLIAHQGTH
Q62518_MOUSE	1688	YKCVDCGKEFSRPSSLQAHQGIH
Q61117_MOUSE	1689	YRCEECEGKGFSWSSSLLIHQRAH
Q61117_MOUSE	1690	YKCEECEGKVFSWSSYLKAHQRVH
Q61116_MOUSE	1691	FKCEECEGKEFRWSVGLSSHQRVH
Q61117_MOUSE	1692	YKCETCGKAFSRVSILQVHQRVH
Q61116_MOUSE	1693	YKCEECEGKGFSASSFQSHQRVH
Q61116_MOUSE	1694	YKCGECGKGFSHASSLQAHHSVH
Q61117_MOUSE	1695	YQCAECGRGFTVESHLQAHQORSH
Q61117_MOUSE	1696	YQCEECEGRGFCRASNFLAHRGVH
Q61117_MOUSE	1697	YKCEECEGKGFTRASTLLDHQORGH
Q61117_MOUSE	1698	YVCEECEGKGFSQASHLLAHQORGH
Q62518_MOUSE	1699	YNCETCGSAFSQASHLQDHQORLH
ZF29_MOUSE	1700	YRCPECEGKGFSWNSVLI IHQRIH
O70162_MOUSE	1701	YCCGECDLGFTQVSRLTEHQRIH
KID1_MOUSE	1702	YRCSECEGKGFTSISR LNRHRI IH
TYY1_MOUSE	1703	YVCPFDGCNKKFAQSTNLKSHILTH
REX1_MOUSE	1704	YQCTFEGCGKRFSLDFNLRTHIRIH
TYY1_MOUSE	1705	FQCTFEGCGKRFSLDFNLRTHVRIH
MTF1_MOUSE	1706	YQCTFEGCPRTYSTAGNLRTHQKTH
GLI_MOUSE	1707	HKCTFEGCRKSYSRLENLKTHLRSH
GLI3_MOUSE	1708	HKCTFEGCTKAYSRLLENLKTHLRSH
ZIC2_MOUSE	1709	FQCEFEFGCDRRFANS SDRKKHMHVH
ZIC1_MOUSE	1710	FKCEFEFGCDRRFANS SDRKKHMHVH
ZIC3_MOUSE	1711	FKCEFEFGCDRRFANS SDRKKHMHVH
ZIC4_MOUSE	1712	FRCEFEFGCERRFANS SDRKKHSHVH
GLI_MOUSE	1713	YMCEQEGCSKAFSNASDRAKHQNRTH
GLI3_MOUSE	1714	YVCEHEGCNKAFSNASDRAKHQNRTH
O70230_MOUSE	1715	YVCTVPGCDKRFTEYSSLYKHHVVH
MTF1_MOUSE	1716	FECDVQGCEKAFNTLYRLKAHQORLH
MTF1_MOUSE	1717	FVCNQEGCGKAFLTSYSLRIHV RVH
O70230_MOUSE	1718	YQCEHSGCGKAFATGYGLKSHFRTH
MTF1_MOUSE	1719	FRCDHDGCGKAFAASHHLKTHV RTH
O70230_MOUSE	1720	FKCPIEGCGRSFTTSNIRKVHIRTH
ZIC4_MOUSE	1721	FPCPFPGCGKVFARSEN LKIHKRTH
ZIC2_MOUSE	1722	FPCPFPGCGKVFARSEN LKIHKRTH
ZIC1_MOUSE	1723	FPCPFPGCGKVFARSEN LKIHKRTH
ZIC3_MOUSE	1724	FPCPFPGCGKIFARSEN LKIHKRTH
O70230_MOUSE	1725	YYCTEPGCGRAFASATNYKNHVRIH
O70230_MOUSE	1726	YRCSEDNCTKSFKTSGDLQKHIRTH
MTF1_MOUSE	1727	FNCESQGCSKYFTTLSDLRKHIRTH
O70230_MOUSE	1728	FRCKYDGCCKLYTTAHLKVHERSH
BTE1_MOUSE	1729	HKCPYSGCGKVYGKSSHLKAHYRVH
Q9Z0Z7_MOUSE	1730	--CDYNGCTKVYTKSSHLKAHLRTH

Q60980_MOUSE	1731	HRCDDYDGCNKVYTKSSHLKAHRRTH
O35738_MOUSE	1732	HRCDFEGCNKVYTKSSHLKAHRRTH
Q61596_MOUSE	1733	HICSHPGVGKTYFKSSHLKAHVRRTH
O89091_MOUSE	1734	HICSHPGCGKTYFKSSHLKAHVRRTH
Q60843_MOUSE	1735	HTCSYTNCGKTYTKSSHLKAHLRTH
EZF_MOUSE	1736	HTCDYAGCGKTYTKSSHLKAHLRTH
Q64167_MOUSE	1737	HICHIQGCCKVYGKTSHLRAHLRWH
O89090_MOUSE	1738	HICHIQGCCKVYGKTSHLRAHLRWH
O89087_MOUSE	1739	HICHIQGCCKVYGKTSHLRAHLRWH
Q62445_MOUSE	1740	HVCHIEGCGKVYGKTSHLRAHLRWH
O70261_MOUSE	1741	HTCGHEGCGKSYTKSSHLKAHLRTH
EKLF_MOUSE	1742	HTCGHEGCGKSYTKSSHLKAHLRTH
WT1_MOUSE	1743	FMCAYPGCNKRYFKLSHLQMHRSKH
ZEP1_MOUSE	1744	YICEYCNRAKPSVLLKHIRSH
Q61479_MOUSE	1745	YICQYCSRPCAKPSVLQKHIRSH
O55140_MOUSE	1746	YICPYCSRACAKPSVLKKHIRSH
Q60636_MOUSE	1747	HECQVCHKRFSSTSNLKTHLRLH
SNAI_MOUSE	1748	CVCTTCGKAFSRPWLLQGHVRRTH
P97469_MOUSE	1749	CVCKICGKAFSRPWLLQGHIRTH
ZIC2_MOUSE	1750	HVCFWEECPREGKPFKAKYKLVNHIRVH
ZIC3_MOUSE	1751	HVCYWEECPREGKSFKAKYKLVNHIRVH
Q62065_MOUSE	1752	HECKLCGASFRTKGSLIRHRRH
Q62065_MOUSE	1753	HVCQFCSRGFREKGS�VRHVRRH
IKAR_MOUSE	1754	FQCNQCGASFRTQKGNLLRHIKLH
Q9Z2Z2_MOUSE	1755	FHCNQCGASFRTQKGNLLRHIKLH
HELI_MOUSE	1756	FHCNQCGASFRTQKGNLLRHIKLH
Q61164_MOUSE	1757	HKCHLCGRAFRVTLLRNHLNTH
Q61624_MOUSE	1758	HVCEHCNAAFRTNYHLQRHVFIH
P97475_MOUSE	1759	HVCEHCNAAFRTNYHLQRHVFIH
Z151_MOUSE	1760	YVCTHCQRQFADPGGLQRHVRIH
Q62511_MOUSE	1761	YICEYCARAFKSSHNLAHVHRMIH
MAZ_MOUSE	1762	YICALCAKEFKNGYNLRRHEAIIH
O88939_MOUSE	1763	YECNICKVRFTRQDKLKVHMRKH
Q64321_MOUSE	1764	--CEVCGVRFTRNDKLLKIHMRRKH
P97365_MOUSE	1765	PHKCEVCGKCFSRKDKLKTHMRCH
O88939_MOUSE	1766	YLCQQCGAAFAHNYDLKNHMRVH
Q64321_MOUSE	1767	YSCPHCPARFLHSYDLKNHMHLLH
Z151_MOUSE	1768	HKCEDCGKEFTHTGNFKRHIRIH
Z151_MOUSE	1769	YRCGDCGKLFTTSGNLKRHQLVH
Z151_MOUSE	1770	-KCRECGKQFTTSGNLKRHLRIH

Chicken database.

5	35 finger units	SEQ ID NO	
	Q92010_CHICK	1771	YSCEVCGKSFI RAPDLKKHERVH

Q90851_CHICK	1772	YPCTICGKKFTQRGTMTRHMRS
Q90850_CHICK	1773	YPCTICGKKFTQRGTMTRHMRS
Q90851_CHICK	1774	--CDACGMRFTROYLTEHMR
Q90850_CHICK	1775	--CDACGMRFTROYLTEHMR
CTCF_CHICK	1776	HKCPDCDMAFVTSGELVRHRRYKH
ZKR1_CHICK	1777	-TCGDCGKGFAWASHLQRHRRVH
ZKR1_CHICK	1778	HRCGDCGKGFAWASHLQRHRRVH
ZKR1_CHICK	1779	HRCGDCGKGFWASHLERHRRVH
ZKR1_CHICK	1780	--CPDCGKSFPWASHLERHRRVH
Q92010_CHICK	1781	--CHMCDKAFKHKSHLKDHERRH
O42408_CHICK	1782	HECGICKKAFKHKHHLIEHMR
DEFI_CHICK	1783	HECGICKKAFKHKHHLIEHMR
O42408_CHICK	1784	FKCTECGKAFKYKHHLKEHLRIH
DEFI_CHICK	1785	FKCTECGKAFKYKHHLKEHLRIH
O42409_CHICK	1786	YPCQYCGKRFHQKSDMKKHTYIH
O42409_CHICK	1787	FECKMCGKTFKRSSTLSTHLLIH
ZKR1_CHICK	1788	YECPECGEAFSQGSHLTKHRRSH
ZKR1_CHICK	1789	YECPECGEAFSQGSHLTKHRRSH
ZKR1_CHICK	1790	YSCPECGESYSQSSHLVQHRRTH
O42409_CHICK	1791	HKCQVCGKAFSQSSNLITHSRKH
O57415_CHICK	1792	YQCNICDYIAADKAALIRHLRTH
CTCF_CHICK	1793	FQCSLCSYASRDYKLKRHMRT
O57415_CHICK	1794	YKCQTCERTFTLKHSLVRHQRIH
Q92010_CHICK	1795	FVCEMCTKGFTTQAHLEHLKIH
O57415_CHICK	1796	-TCPYCPRVFSWASSLQRHMLTH
O57415_CHICK	1797	HSCSICGKSLSSASSLDRHMLVH
O57415_CHICK	1798	--CTVCNKRFWSLQDLTRHMRS
Q91051_CHICK	1799	CVCKICGKAFSRPWLLQGHIRTH
O12939_CHICK	1800	CVCKMCGKAFSRPWLLQGHIRTH
O57415_CHICK	1801	YKCSVCGQSFTTNGNMHRHMKIH
IKAR_CHICK	1802	FQCNQCGASFTQKGNLLRHIKLH
CTCF_CHICK	1803	HKCHLCGRAFRVTLLRNHLNTH
O93567_CHICK	1804	YECNICNVRFTRQDKLKVHMRKH
O93567_CHICK	1805	YLCQQCGAAFAHNYDLKNHMRVH

Plant Database.

52 finger units

SEQ ID NO

O22089_PETHY	1806	HECSICGEQFLLGQALGGHMRKH
O22088_PETHY	1807	HECSFCGEDFPTGQALGGHMRKH
O22087_PETHY	1808	-ECSFCGEDFPTGQALGGHMRKH
Q39092_ARATH	1809	HKCKLCWKSFANGRALGGHMRSH
Q39217_ARATH	1810	HKCSICSQSFGTGQALGGHMRRH
P93713_PETHY	1811	HECSICGLEFAIGQALGGHMRRH
O22086_PETHY	1812	HECSICGLEFPIGQALGGHMRRH
O22085_PETHY	1813	HECSICGMEFSLGQALGGHMRRH
O22084_PETHY	1814	HECSICGMEFSLGQALGGHMRRH
Q42453_ARATH	1815	HPCPICGVKFPMGQALGGHMRRH
Q42410_ARATH	1816	HPCPICGVVEFPMGQALGGHMRRH
O65150_TOBAC	1817	HVCSICHKAFPTGQALGGHKRRH
Q40897_PETHY	1818	HVCSICHKAFPTGQALGGHKRRH
Q40896_PETHY	1819	HVCSICHKAFPSGQALGGHKRRH
Q42430_WHEAT	1820	HRCISICQKEFPTGQALGGHKRKH
Q40899_PETHY	1821	HECSICHKCFPTGQALGGHKRCH
P93166_SOYBN	1822	HECSICHKSFPTGQALGGHKRCH
Q96289_ARATH	1823	HVCTICNKSFPSPGQALGGHKRCH
Q42423_ARATH	1824	HVCTICNKSFPSPGQALGGHKRCH
O22533_ARATH	1825	HVCSICHKSFATGQALGGHKRCH
Q40898_PETHY	1826	HECSICHKCFSSGQALGGHKRRH
Q38895_ARATH	1827	YTCSFCKREFRSAQALGGHMNVH
O23621_ARATH	1828	YTCNFCRREFRSAQALGGHMNVH
O80942_ARATH	1829	YTCSFCRREFKSAQALGGHMNVH
P93714_PETHY	1830	HECSYCGAEFTSGQALGGHMRRH
Q43614_PETHY	1831	HECAICGAEFTSGQALGGHMRRH
O22083_PETHY	1832	HECSICGAEFTSGQALGGHMRRH
Q41070_PEA	1833	HECSICGAEFTSGQALGGHMRRH
Q42375_ARATH	1834	HECSICGSEFTSGQALGGHMRRH
O65499_ARATH	1835	HKCNICFRVFSSGQALGGHMRCH
O22090_PETHY	1836	HECPVCFRVFSSGQALGGHKRTH
O22082_PETHY	1837	HECPVCYRVFSSGQALGGHKRSH
P93717_PETHY	1838	HECSICHRVFSTGQALGGHKRCH
O04177_BRARA	1839	HTCSICFKSFSSGQALGGHKRCH
O04176_BRARA	1840	HTCSICFKSFSSGQALGGHKRCH
P93715_PETHY	1841	HQCSICHRVFSSGQALGGHKRCH
Q39092_ARATH	1842	HECPICAKVFTSGQALGGHKRSH
P93719_PETHY	1843	HECPYCDRVFKSGQALGGHKRSH
P93718_PETHY	1844	HACPFCDPRMFKSGQALGGHKRSH
O22091_PETHY	1845	YECPLCFKIFQSGQALGGHKRSH
Q42430_WHEAT	1846	-KCSVCGKSFSSYQALGGHKTSH
O04177_BRARA	1847	YKCTVCGKSFSSYQALGGHKTSH
O04176_BRARA	1848	YKCTVCGKSFSSYQALGGHKTSH
Q96289_ARATH	1849	YKCSVCDKTFSSYQALGGHKASH
Q42423_ARATH	1850	YKCSVCDKTFSSYQALGGHKASH
Q40897_PETHY	1851	YKCSVCDKSFSSYQALGGHKASH
Q40896_PETHY	1852	YKCSVCDKSFSSYQALGGHKASH
Q40898_PETHY	1853	YKCNVCNKSFHSYQALGGHKASH
O65150_TOBAC	1854	YKCSVCDKAFSSYQALGGHKASH
P93166_SOYBN	1855	YKCSVCDKSFPSYQALGGHKASH
Q40899_PETHY	1856	YKCSVCGKGFSGSYQALGGHKASH
O22533_ARATH	1857	YKCSVCDKAFSSYQALGGHKASH

Arabidopsis Database

SEQ ID NO

Q9ZU64/169-191	1858	YTCPKCNSIFDTSQKFAAHMSSH
O23621/40-62	1859	YTCNFCRREFRSAQALGGHMNVH
O23504/5-27	1860	HKCKLCSKSFCONGRALGGHMKSH
Q9SYC5/250-275	1861	WYCSCGSDFKHKRSLKDHSVKAFGNGH
Q9SYC5/224-246	1862	FACRMCGKAFAVKGDWRTHEKNC
O22533/89-111	1863	YKCSVCDKAFSSYQALGGHKASH
O22533/148-170	1864	HVCSI CHKS FATGQALGGHKRCH
Q9SN24/149-171	1865	HNCSICFKSFPSGQALGGHKRCH
Q9SN24/94-116	1866	YKCSVCGKSFPSYQALGGHKTS
Q9STI7/117-140	1867	YFCGVCDRRFYTNEKLINHFQI
Q9STM3/1296-1320	1868	LKCPWKGCKMTFKWAWSRTEHIRVH
Q9STM3/1243-1268	1869	YQCNMEGCTMSFSSEKQLMLHKRNIC
Q9STM3/1271-1290	1870	KGCGKNFFSHKYL VQHQRVH
Q9STM3/1326-1352	1871	YVCAEPDCGQTFRFVSDFSRHKRKTGH
Q9STM3/1296-1320	1872	LKCPWKGCKMTFKWAWSRTEHIRVH
O81801/142-164	1873	PMCNVCGKGFASWKAVFGHLRQH
O65601/61-83	1874	QKCEKCSREFCSPVNFRRHNRMH
Q9SFY6/118-140	1875	YKCSVCDKTFSSYQALGGHKASH
Q9SFY6/174-196	1876	HVCTICNKSFPSPGQALGGHKRCH
O65245/147-171	1877	FYCELCSKQYRTVMEFEGHLSSYDH
Q39261/52-74	1878	FSCNYCQRKFYSSQALGGHQNAH
Q9SSW0/118-140	1879	HVCSVCGKSFATGQALGGHKRCH
Q9SSW0/75-97	1880	YKCGVCYKTFSSYQALGGHKASH
Q39262/61-83	1881	FSCNYCQRTFYSSQALGGHQNAH
Q9SSW1/164-186	1882	HTCSICFKSFASGQALGGHKRCH
Q9SSW1/97-119	1883	YKCTVCGKSFSSYQALGGHKTS
Q9ZPT0/145-167	1884	WVCERCSKGYAVQSDYKAHLKTC
Q9ZPT0/67-89	1885	YICEICNQGFQORDQNLQMHRRRH
Q9ZPT0/172-193	1886	HSCDCGRVFSRVESFIEHQDNC
Q39263/85-107	1887	FSCNYCQRKFYSSQALGGHQNAH
Q9SGD1/291-316	1888	WYCTCGSDFKHKRSLKDHIRSFGSGH
Q9SGD1/265-287	1889	FSCGKCGKALAVKGDWRTHEKNC
Q9SGD1/180-202	1890	FACSI CSKTFNRYNNMQMHMWGH
Q9SSW2/106-128	1891	YKCNVCEKAFFPSYQALGGHKASH
Q9SSW2/165-187	1892	HECSI CHKV FPTGQALGGHKRCH
Q39264/60-82	1893	HECQYCGKEFANSQALGGHQNAH
P93815/7-30	1894	QECVACKRVFLSSHQLISHYNAAH
Q39265/41-63	1895	YECQYCCREFANSQALGGHQNAH
Q39266/59-81	1896	FSCNYCRRKFYSSQALGGHQNAH
Q39267/93-115	1897	FECHYCFRNFPSTQALGGHQNAH
Q9SVY1/301-323	1898	FMCRCGKAFAVRGDWRTHEKNC
Q9SVY1/217-239	1899	FSCPVCFKTFNRYNNMQMHMWGH
Q9SGH2/1804-1827	1900	IHCLICHKTFASDDEFEDHTESKC
Q38895/47-69	1901	YTCSFCKREFRSAQALGGHMNVH
Q9SLB8/49-71	1902	YTCSFCRREFRSAQALGGHMNVH
Q9SL35/188-210	1903	HECSI CGSEFTSGQALGGHMRRH
Q9SL35/113-135	1904	YECKTCNRTFSSFQALGGHRASH

O81013/49-71	1905	HFCVICEKQFSSGKAYGGHVRIH
O81013/119-141	1906	IRCCLCGKEFQTMHSLFGHMRRH
O23395/664-686	1907	LHCEKCGKALQPTMEKHLKVFH
Q9SI97/34-56	1908	FACKTCNKEFSPSFQALGGHRASH
Q9SI97/78-100	1909	HECPICGAFAVGQALGGHMRKH
Q9SR34/35-57	1910	YVCSFCIRGFSNAQALGGHMNIH
Q42485/68-90	1911	FSCNYCQRKFYSSQALGGHQNAH
O82389/126-149	1912	FPCNSCGEIFPKINLLENHIAIKH
Q9SQX8/182-204	1913	YQCKTCDRTFSPSFQALGGHRASH
Q9SQX8/261-283	1914	HECGICGAFTSGQALGGHMRRH
O65499/222-244	1915	HKCNICFRVFSSGQALGGHMRCH
O65499/77-99	1916	RPCTECGRKFWSWKALFGHMRCH
O65499/162-184	1917	FECGGCKKVFGSHQALGGHRASH
Q9SCM4/220-243	1918	DVCPKCSRGFRDPVDLLKHIDKDH
Q96289/80-102	1919	YKCSVCDKTFSSYQALGGHKASH
Q96289/136-158	1920	HVCTICNKSFPSPGQALGGHKRCH
Q9SCQ6/139-161	1921	WKCDKCSKKYAVQSDWKAHSKIC
Q9SCQ6/166-187	1922	YKCDCGTLFSRRDSFITHRAFC
Q9SCQ6/63-85	1923	FVCEICNKGFFORDQNLQLHRRGH
Q9SFS1/70-92	1924	YVCEICNQGFORDQNLQMHRRRH
Q9SFS1/148-170	1925	WICERCSKGYAVQSDYKAHLKTC
Q9SFS1/175-196	1926	HSCDCGRVFSRVESFIEHQDTC
Q9SSA6/575-598	1927	IHCLICHKTFASDDEFEDHTESKC
Q42410/39-61	1928	FTCKTCLKQFHSFQALGGHRASH
Q42410/82-104	1929	HPCPICGVEFPMGQALGGHMRRH
Q9XFP6/12-35	1930	VWCYYCDREFDDEKILVQHQAQKH
Q9XFP6/36-59	1931	FKCHVCHKKLSTASGMVIHVLQVH
O22238/218-241	1932	VSCGSCKKTFNSGNALESHNKAKH
Q42453/40-62	1933	FRCKTCLKEFSSFQALGGHRASH
Q42453/86-108	1934	HPCPICGVKFPMGQALGGHMRRH
Q42375/113-135	1935	YECKTCNRTFSSFQALGGHRASH
Q42375/188-210	1936	HECSICGSEFTSGQALGGHMRRH
O22759/159-181	1937	WKCEKCSKFYAVQSDWKAHTKIC
O22759/186-207	1938	YRCDCGTLFSRKDTFITHRAFC
O22759/82-104	1939	FVCEICNKGFFORDQNLQLHRRGH
Q9ZUL3/81-103	1940	FICEVCNKGFFOREQNLQLHRRGH
Q9ZUL3/157-179	1941	WKCDKCSKRYAVQSDWKAHSKTC
Q9ZUL3/184-205	1942	YRCDCGTLFSRRDSFITHRAFC
P93751/95-117	1943	FECHYCFRNFTSQALGGHQNAH
O81827/196-219	1944	VSCHKCGEKFSKLEAAEAHHLTKH
Q9ZUL4/82-104	1945	WKCEKCSKRYAVQSDWKAHSKTC
Q9ZUL4/109-130	1946	YRCDCGTIFSRRDSYITHRAFC
Q9ZUL4/6-28	1947	FICDVCNKGFFOREQNLQLHRRGH
Q9SHD0/194-216	1948	FKCETCGKVFKSYQALGGHRASH
Q9SHD0/243-265	1949	HECPICFRVFTSGQALGGHKRSH
Q9SHD0/4-26	1950	YKCRFCFKSFINGRALGGHMRSH
O64936/131-153	1951	YQCNVCGRELPSYQALGGHKASH

O64936/179-201	1952	HKCSICHREFSTGHSLGGHKRLH
Q9SIJ0/65-87	1953	RPCTECGKQFGSLKALFGHMRCH
Q9SIJ0/148-170	1954	FEC DGCKKVFGSHQALGGHRATH
Q9SIJ0/211-233	1955	HRCNICSRVFSSGQALGGHMRCH
Q9SLD4/47-69	1956	FECKTCNKRFSSFQALGGHRASH
Q9SLD4/94-116	1957	HKCSICSQSFGTGQALGGHMRRH
Q9ZU93/121-143	1958	FECPICKNPFTSEEEVSVHVESC
Q9SFT3/177-200	1959	CACPQCGEVFPKLESLEHHQAVRH
Q9ZQE0/244-266	1960	YTCPKCNGVFNTSQKFAAHMSSH
Q42423/80-102	1961	YKCSVCDKTFSSYQALGGHKASH
Q42423/136-158	1962	HVCTICNKSFPSPGQALGGHKRCH
Q9ZWA6/146-168	1963	WKCEKCAKRYAVQSDWKAHSTC
Q9ZWA6/173-194	1964	YRCDCGTIFSRRDSFITHRAFC
Q9ZWA6/70-92	1965	FLCEICGKGFQORDQNLQLHRRGH
O80942/39-61	1966	YTCSFCRREFKSAQALGGHMNVH
Q39217/90-112	1967	HKCSICSQSFGTGQALGGHMRRH
Q39217/43-65	1968	FECKTCNKRFSSFQALGGHRASH
Q39092/160-182	1969	FECETCEKVFKSYQALGGHRASH
Q39092/209-231	1970	HECPICAKVFTSGQALGGHKRSH
Q39092/5-27	1971	HKCKLCWKSFANGRALGGHMRSH
O81793/138-160	1972	PVCHICGRGFGSWKAVFGHMRAH
O64828/530-553	1973	LQCI PCGSHFGDKEQLLVHVQAVH
O64828/599-622	1974	FVCKFCGLKFNLLPDLGRHHQAEH
O64828/496-519	1975	FACAI CLDSFVRRKLLEIHVEERH
O49591/251-278	1976	FMCLYCNELCRPFSSLEAVRKHMEAKSH
O49591/26-50	1977	LTCNACNMEFKDEEERNLHYKSDWH
O49591/90-114	1978	YTCAICAKGYRSSKAHEQHLQSRSH

There follow several examples of how to construct and select DNA-binding sub-domains from libraries of natural zinc fingers.

5

Example 4: Human Zinc Finger Module ‘Mini-Library’.

As a preliminary test of the efficacy of using natural zinc finger modules for constructing novel DNA-binding domains, a ‘mini-library’ of natural, human zinc finger modules is generated. The mini-library comprises 8 zinc finger modules, which have the following nomenclature assigned to them in the human genome database: Zif268 finger 1, Zif268 finger 2, Sp1 finger 3, WT1 finger 1, O15391, O75626, ZN45 and Z165. Since there is more than one zinc finger module belonging to the zinc fingers proteins ZN45 and Z165, we have called the selected modules ZN45-(AAA) and Z165-(GCC) respectively,

10

according to their predicted binding site. We have also predicted the binding sites for the zinc fingers O15391 and O75626. The preferred binding sites for Zif268 finger 1, Zif268 finger 2, Sp1 finger 3 and WT1 finger 1 are already known. The amino acid sequences of each of the stated modules, and their predicted or previously determined binding
 5 sequences are shown in Table 3.

Two 3-zinc finger peptide libraries are prepared, containing the 8 zinc finger modules stated. All novel 3-finger peptides contain a leader sequence, MAEERP (SEQ ID NO:16), at the start of the peptide and are tagged by the sequence
 10 LRQKDGGGSYPYDVPDYA (SEQ ID NO:1989) at the C-terminus. This sequence provides: (in the absence of a further C-terminal finger) a suitable terminus to the final α -helix of the peptide –LRQKD- (SEQ ID NO:1987) as found in wild-type Zif268; a short, flexible linker sequence, GGGS (SEQ ID NO:2121); and an HA-tag (YPYDVPDYA (SEQ ID NO:2122)), which is recognised by the HA-antibody. Adjacent zinc finger
 15 modules are fused using the linker peptide sequence TGEKP (SEQ ID NO:3). The peptide sequences described above are also displayed in Table 3.

In the first library (library 1), the 8 zinc finger modules are recombined in random order to create 3-finger peptides with all possible combinations of the 8 zinc finger modules.
 20 Such a procedure results in a library diversity of 512 ($=8^3$), comprising peptides that are predicted to bind to any possible combination of the binding sites assigned in Table 3. Library 1 allows novel 3-finger domains to be selected as a unit, for specified 9 bp target sequences. Such 3-finger units may be used for the construction of poly-zinc finger peptides as described in Moore, M., Choo, Y. & Klug, A. (2001) *Proc. Natl. Acad. Sci.*
 25 *USA* 98: 1432-1436; and WO 01/53480.

In the second library (library 2), the 8 zinc finger modules are randomly recombined to create 2-finger peptides which are all joined to the C-terminus of Zif268 finger 1. The invariant finger 1 acts as an anchor for the selection, both by providing extra affinity to
 30 stabilise the selection, and by fixing the register of the protein DNA interaction (as discussed *supra*). Such a library has a diversity of 64 ($=8^2$), and allows novel 2-finger units to be selected for a given 6 bp target sequence. The resulting 2 finger units can be

recovered by PCR and used in the construction of poly-zinc finger peptides (based on strings of 2-finger units), as described in WO 01/53480.

These two libraries (encoding 3-finger peptides) are screened, as described below, for the ability of their encoded proteins to bind three different 9 bp binding sequences: 5'-GCG-TGG-GCG-3'; 5'-GGA-TAA-GCG-3'; and 5'-GCC-GAG-TGG-3'.

As positive controls, the genes encoding the 3-finger peptides predicted to bind the above target sequences are specifically constructed and tested in a similar manner.

x	FINGER/UNIT	SEQ ID NO:	PEPTIDE SEQUENCE	SITE
1	ZIF268 F1	1979	YACPVESCDRRFSRSDDELTRHIRIH	GCG
2	ZIF268 F2	1980	FQCRICMRNFSRSDHLSTHIRTH	TGG
3	Sp1 F3	1981	FSCPICEKRFMRS DHLT KHARRH	GGG
4	WT1 F1	1982	FMCAYPGCNKRYFKLSHLQMHSRKH	GAG
5	O15391	1983	FVCPFDVCNRKFAQSTNLKTHILTH	TAA ¹
6	O75626	1984	FKCQTCNKGFTQLAHLQKHVYLVH	GGA ¹
7	ZN45-AAA	1985	YKCEECGKGFSQASNLLAHQRGH	AAA ¹
8	Z165-GCC	1986	YECNECGKSFAESSDLTRHRRIH	GCC ¹
9	leader	16	MAEERP	-
10	linker	3	TGEKP	-
11	G ₃ S-HA-tag	1989	LRQKDGGGSYPYDVPDYA*	-

¹Predicted binding site. *indicates a translation stop codon.

Table 3. Nomenclature, amino acid sequences and known or predicted binding sequences ("SITE") of zinc finger modules and other peptide units used in library construction.

a. Human Zinc Finger Mini-Library Construction.

Two libraries are prepared, according to the scheme shown in Figure 2. The N-terminal finger of the 3-finger construct is referred to as 'cassette A'. The central finger is encoded by cassette B, and the third (C-terminal) finger module is called cassette C.

Zinc Finger Cassettes

Polynucleotide sequences encoding the amino acid sequences of the 8 zinc finger modules shown in Table 3 are determined, taking into account *E. coli* codon preferences,

and the corresponding nucleotide sequences are synthesised as single stranded oligonucleotides, examples of which are shown in Table 4. Also shown are the sequences of exemplary linkers and an exemplary 3'-tag required for the assembly of 3-finger domains. Double stranded cassettes encoding the zinc finger modules and relevant leader, linker, and terminator sequences are generated by PCR according to the procedure described below, using the appropriate oligonucleotide templates of Table 4, and primers of Table 5.

x	CODE	FINGER	SEQ ID NO	NUCLEOTIDE SEQUENCE
1	AS144	ZIF268 F1	1990	TATGCGTGCCCCGGTGGAAAGCTGCGATCGTCGTTTTCAG CCGTAGCGATGAACTGACCCGTCATATTCGTATTCAT
2	AS145	ZIF268 F2	1991	TTTCAGTGCCGTATTTGCGATGCGTAACTTTAGCCGTAG CGATCATCTGAGCACCCATATTCGTACCCAT
3	AS148	Sp1 F3	1992	TTTAGCTGCCCCGATTTGCGAAAAACGTTTTATGCGTAG CGATCATCTGACCAAACATGCGCGTCGTCAT
4	AS149	WT1 F1	1993	TTTATGTGCGCGTATCCGGGCTGCAACAAACGTTATTT TAAACTGAGCCATCTGCAGatgCATAGCCGTAAACAT
5	AS150	O15391	1994	TTTGTGTGCCCCGTTTGATGTGTGCAACCGTAAATTTGC GCAGAGCACCAACCTGAAAACCCATATTCTGACCCAT
6	AS151	O75626	1995	TTTAAATGCCAGACCTGCAACAAAGGCTTTACCCAGCT GGCGCATCTGCAGAAACATTATCTGGTGCAT
7	AS152	ZN45- AAA	1996	TATAAATGCGAAGAATGCGGCAAAGGCTTTAGCCAGGC GAGCAACCTGCTGGCGCATCAGCGTGGCCAT
8	AS153	Z165-GCC	1997	TATGAATGCAACGAATGCGGCAAAAGCTTTGCGGAAAG CAGCGATCTGACCCGTCATCGTCGTATTCAT
9		MAEERP leader	1998	ATGGCGGAAGAACGTCCG
10		TGEKP linker	1999	ACCGGCGAAAAACCG
11		G ₃ S-HA- tag (tag)	2000	CATCTGCGCCAGAAGGACGGCGGGCGGCAGCTATCCGTA TGATGTGCCGGATTATGCGTAA

10 **Table 4.** Nucleotide sequences encoding zinc finger modules and other peptide sequences used in the construction of 3-finger proteins.

x	CODE	NAME	SEQ ID NO	SEQUENCE
1	AS5	pETFwd1	2001	CGCTGACTTCCGCGTTTCC
2	AS86	SDRev	2002	ATGTATATCTCCTTCTTAAAGTT
3	AS93	ZnF1Fwd	2003	AACTTTAAGAAGGAGATATACATATGGCGGAAGAA CGTCCGTATGCGTGCCCGGTGGAAAG
4	AS94	ZnF2Fwd	2004	AACTTTAAGAAGGAGATATACATATGGCGGAAGAA CGTCCGTTTCAGTGCCGTATTTGCATG

5	AS95	ZnF3Fwd	2005	AAC TTTAAGAAGGAGATATACATATGGCGGAAGAA CGTCCGTTTAGCTGCCCCGATTTGCG
6	AS96	ZnF4Fwd	2006	AAC TTTAAGAAGGAGATATACATATGGCGGAAGAA CGTCCGTTTATGTGCGCGTATCCGGG
7	AS97	ZnF5Fwd	2007	AAC TTTAAGAAGGAGATATACATATGGCGGAAGAA CGTCCGTTTATGTGCGCGTATCCGGG
8	AS98	ZnF6Fwd	2008	AAC TTTAAGAAGGAGATATACATATGGCGGAAGAA CGTCCGTTTAAATGCCAGACCTGCAAC
9	AS99	ZnF7Fwd	2009	AAC TTTAAGAAGGAGATATACATATGGCGGAAGAA CGTCCGTATAAATGCGAAGAATGCGGC
10	AS100	ZnF8Fwd	2010	AAC TTTAAGAAGGAGATATACATATGGCGGAAGAA CGTCCGTATGAATGCAACGAATGCGGC
11	AS101	1Link1Rev	2011	CGGTTTTTCGCCGGTATGAATACGAATATGACGGG
12	AS102	1Link2Rev	2012	CGGTTTTTCGCCGGTATGGGTACGAATATGGGTGC
13	AS103	1Link3Rev	2013	CGGTTTTTCGCCGGTATGACGACGCGCATGTTTGG
14	AS104	1Link4Rev	2014	CGGTTTTTCGCCGGTATGTTTACGGCTATGCATCT G
15	AS105	1Link5Rev	2015	CGGTTTTTCGCCGGTATGGGTCAGAATATGGGTTT TC
16	AS106	1Link6Rev	2016	CGGTTTTTCGCCGGTATGCACCAGATAATGTTTCT GC
17	AS107	1Link7Rev	2017	CGGTTTTTCGCCGGTATGGCCACGCTGATGCGC
18	AS108	1Link8Rev	2018	CGGTTTTTCGCCGGTATGAATACGACGATGACGGG
19	AS109	1Link1Fwd	2019	CATACCGGCGAAAAACCGTATGCGTGCCCGGTGGA AAG
10	AS110	1Link2Fwd	2020	CATACCGGCGAAAAACCGTTTCAGTGCCGTATTTG CATG
11	AS111	1Link3Fwd	2021	CATACCGGCGAAAAACCGTTTAGCTGCCCCGATTTG CG
12	AS112	1Link4Fwd	2022	CATACCGGCGAAAAACCGTTTATGTGCGCGTATCC GGG
13	AS113	1Link5Fwd	2023	CATACCGGCGAAAAACCGTTTGTGTGCCCGTTTGA TGTG
14	AS114	1Link6Fwd	2024	CATACCGGCGAAAAACCGTTTAAATGCCAGACCTG CAAC
15	AS115	1Link7Fwd	2025	CATACCGGCGAAAAACCGTATAAATGCGAAGAATG CGGC
16	AS116	1Link8Fwd	2026	CATACCGGCGAAAAACCGTATGAATGCAACGAATG CGGC
17	AS117	2Link1Rev	2027	TGGCTTCTCACC CGTGTGATGAATACGAATATGAC GGGTC
18	AS118	2Link2Rev	2028	TGGCTTCTCACC CGTGTGATGGGTACGAATATGGG TGC
19	AS119	2Link3Rev	2029	TGGCTTCTCACC CGTGTGATGACGACGCGCATGTT TGG
20	AS120	2Link4Rev	2030	TGGCTTCTCACC CGTGTGATGTTTACGGCTATGCA TCTG

21	AS121	2Link5Rev	2031	TGGCTTCTCACCCGTGTGATGGGTCAGAATATGGG TTTTC
22	AS122	2Link6Rev	2032	TGGCTTCTCACCCGTGTGATGCACCAGATAATGTT TCTGC
23	AS123	2Link7Rev	2033	TGGCTTCTCACCCGTGTGATGGCCACGCTGATGCG C
24	AS124	2Link8Rev	2034	TGGCTTCTCACCCGTGTGATGAATACGACGATGAC GGG
25	AS125	2Link1Fwd	2035	CACGGGTGAGAAGCCATATGCGTGCCCGGTGGAAA G
26	AS126	2Link2Fwd	2036	CACGGGTGAGAAGCCATTTTCAGTGCCGTATTTGCA TG
27	AS127	2Link3Fwd	2037	CACGGGTGAGAAGCCATTTAGCTGCCCGATTTGCG
28	AS128	2Link4Fwd	2038	CACGGGTGAGAAGCCATTTATGTGCGCGTATCCGG G
29	AS129	2Link5Fwd	2039	CACGGGTGAGAAGCCATTTGTGTGCCCGTTTGATG TG
30	AS130	2Link6Fwd	2040	CACGGGTGAGAAGCCATTTAAATGCCAGACCTGCA AC
31	AS131	2Link7Fwd	2041	CACGGGTGAGAAGCCATATAAATGCGAAGAATGCG GC
32	AS132	2Link8Fwd	2042	CACGGGTGAGAAGCCATATGAATGCAACGAATGCG GC
33	AS133	3HA1Rev	2043	CTAGGAATTCTTACGCATAATCCGGCACATCATA GGATAGCTGCCGCCGCCGTCCTTCTGGCGCAGATG AATACGAATATGACGGGTC
34	AS134	3HA2Rev	2044	CTAGGAATTCTTACGCATAATCCGGCACATCATA GGATAGCTGCCGCCGCCGTCCTTCTGGCGCAGATG GGTACGAATATGGGTGC
35	AS135	3HA3Rev	2045	CTAGGAATTCTTACGCATAATCCGGCACATCATA GGATAGCTGCCGCCGCCGTCCTTCTGGCGCAGATG ACGACGCGCATGTTTGG
36	AS136	3HA4Rev	2046	CTAGGAATTCTTACGCATAATCCGGCACATCATA GGATAGCTGCCGCCGCCGTCCTTCTGGCGCAGATG TTTACGGCTATGCATCTG
37	AS137	3HA5Rev	2047	CTAGGAATTCTTACGCATAATCCGGCACATCATA GGATAGCTGCCGCCGCCGTCCTTCTGGCGCAGATG GGTCAGAATATGGGTTTTC
38	AS138	3HA6Rev	2048	CTAGGAATTCTTACGCATAATCCGGCACATCATA GGATAGCTGCCGCCGCCGTCCTTCTGGCGCAGATG CACCAGATAATGTTTCTGC
39	AS139	3HA7Rev	2049	CTAGGAATTCTTACGCATAATCCGGCACATCATA GGATAGCTGCCGCCGCCGTCCTTCTGGCGCAGATG GCCACGCTGATGCGC
40	AS140	3HA8Rev	2050	CTAGGAATTCTTACGCATAATCCGGCACATCATA GGATAGCTGCCGCCGCCGTCCTTCTGGCGCAGATG AATACGACGATGACGGG

41	AS141	Rev3	2051	CTAGGAATTCTTACGCATAATC
42	AS142	1LinkRev	2052	CGGTTTTTCGCCGGTATG
43	AS143	2LinkRev	2053	TGGCTTCTCACCCGTGTG

Table 5. Modifying oligonucleotides used for mini-library construction.

5 **1. Library 1.**

Once made into double stranded DNA cassettes, the finger units are attached to T7 upstream expression sequences by PCR overlap extension, using the following protocol.

10 (a) Upstream sequences are first extracted from pET23a by PCR using primers pETFwd1 and SDRev, generating the fragment pET5'.

 (b) The fingers for cassette A are amplified with forward primers ZnFxFwd (AS93-100) and reverse primers 1LinkxRev (AS101-AS108), where x is the number of a
15 particular finger from Tables 3 and 4, as indicated.

 (c) The fingers for cassette B are amplified with forward primers 1LinkxFwd (AS109-116) and reverse primers 2LinkxRev (AS117-AS124), where x refers to the
20 finger module number.

 (d) The fingers for cassette C are amplified with forward primers 2LinkxFwd (AS125-132) and reverse primers 3HAxRev (AS133-AS140), where x refers to the
25 appropriate zinc finger module.

 The steps to create cassettes A, B and C are performed separately, however, mixed populations of template oligonucleotides can be added to each PCR of steps (a), (b), and (c) to produce a library of each cassette.

 The final 3-finger library is assembled by overlap extension as outlined in Figure 2. In
30 the first step the mixed pool of cassette A is appended to the upstream sequences, pET5'.

Equimolar amounts are mixed and PCR-cycled in the absence of primers. The reaction product is either purified immediately or reamplified before purification using primers pETFwd1 and 1LinkRev.

5 In the second step cassette B (mixed pool) is appended to the product of the above step. Again, equimolar amounts are mixed and PCR-cycled in the absence of primers. The reaction product is either purified immediately or reamplified before purification using primers pETFwd1 and 2LinkRev.

10 In the final step cassette C (mixed pool) is appended to the above product. Equimolar amounts are mixed and PCR-cycled in the absence of primers. As before, the reaction product may be purified immediately or reamplified before purification using primers pETFwd1 and Rev3. (see, also Figure 2).

15 **2. Library 2.**

Library 2 is assembled in a similar manner to Library 1 except that cassette A is represented by Zif268 finger 1 only.

20 The final PCR products containing T7 promoter sequences and encoding 3-finger peptides attached to an HA-antibody tag are purified and used for the production of protein.

25 **b. Zinc Finger Library Screening.**

Two exemplary methods for screening zinc finger libraries, such as those produced above, are described in Protocol A and Protocol B, below.

Protocol A:

The peptides of library 1 and library 2 are screened to select 3-zinc finger domains which bind the sequences: 5'-GCG-TGG-GCG-3'; 5'-GGA-TAA-GCG-3'; and 5'-GCC-GAG-TGG-3'. Since library 2 contains Zif268 finger 1 in the N-terminal position, in theory, these peptides should only bind the sequences, 5'-GCG-TGG-GCG-3', and 5'-GGA-TAA-GCG-3'. Hence, library 2 is effectively used to select 2-finger units which bind strongest to the 6 bp sequences, 5'-GCG-TGG-3', and 5'-GGA-TAA-3'. Double stranded binding sites for use in the selection protocol are generated by annealing the complimentary oligonucleotides: Zif.b site and Zif site RC (AS154 and AS155); #1#5#6.b and #1#5#6 RC (AS156 and AS157); and #2#4#8.b and #2#4#8 RC (AS158 and AS159). The top strand of each binding site is biotinylated, allowing capture of binding site/zinc finger/HA-antibody ternary complexes to the streptavidin-coated plate in an ELISA screening assay. The oligonucleotides are displayed in Table 6, below.

15

x	Code	Name	SEQ ID NO	Sequence
1	AS154	Zif.b site	2054	TTTTTTTTTTTGC GTGGGCGTTTTTTTTTTT
2	AS155	Zif site RC	2055	AAAAAAAAAAACGCCACGCAAAAAAAAAAA
3	AS156	#1#5#6.b	2056	TTTTTTTTTTTGGATAAGCGTTTTTTTTTTT
4	AS157	#1#5#6 RC	2057	AAAAAAAAAAACGCTTATCCAAAAAAAAAAA
5	AS158	#2#4#8.b	2058	TTTTTTTTTTGCCTGTTGGTTTTTTTTTTT
6	AS159	#2#4#8 RC	2059	AAAAAAAAAAACCAACAGGCAAAAAAAAAAA

Table 6. Oligonucleotide sequences used to generate double stranded binding sites used in the selection procedure.

20

The PCR-amplified 3-finger constructs are gel-purified from a 1% TAE-agarose gel using the Gel Extraction Kit (Qiagen) and quantified based on absorbance at 260 nM. Dilutions (in 0.25 mg/ml λ DNA) of DNA template encoding for either library 1 or 2 are prepared at the final total template concentration of 4.2 fM and 1 fM, respectively. At these concentrations 1 μl of template contains approximately 2500 and 600 molecules of library 1 or library 2, respectively. At such low concentrations, such samples must be PCR amplified to generate enough template for protein expression. Hence, these 1 μl aliquots

25

are taken and added to 1 ml PCR pre-mix, containing primers Rev3 (AS141) and pETFwd2 (primer sequences shown below, see Table 7). The PCR pre-mixes are then aliquoted into 96 (or 384) well plates at 10 μ l per well, which is the equivalent of approximately 25 or 6 molecules of library 1 or library 2 template, respectively.

5 Templates are amplified using 30 cycles of PCR. After this first round of PCR, 0.5 μ l aliquots of PCR product are added to new 10 μ l PCR pre-mixes (in 96 or 384 well format), containing nested primers, pETFwd3 and Rev3, and amplified for another 30 cycles. The resultant product is concentrated enough to perform *in vitro* transcription / translation.

10

In vitro translation experiments using TNT PCR coupled transcription-translation mix (Promega) are assembled according to the manufacturer's instructions. Typically 5 μ l final volume contains 1 μ l of each PCR product and 4 μ l rabbit reticulocyte pre-mix (containing 20 μ M methionine, 12.5 μ g/ml λ Hind III digest (Roche), 500 μ M ZnCl₂ (Sigma), 0.7 μ l H₂O, 40 nM PCR-amplified DNA template). Reactions are incubated at 15 30°C for 90 minutes. 50 μ l PBS binding buffer containing 0.1 % BSA (Sigma), 0.5% Tween 20 (Sigma), 50 μ M ZnCl₂, 10 nM of the appropriate biotinylated binding site, 25 μ U/ml rat 3F10 anti-HA HRP conjugate (Roche) is added to the translation mix and incubated for 45 minutes at room temperature. The binding mix is thereafter transferred 20 to pre-blocked black streptavidin-coated 8-well strips or 96 / 384 well plates (Roche), and the ternary complexes containing 3-finger peptide, biotinylated binding site and anti-HA HRP antibody are captured while shaking at 200 rpm for 45 minutes at room temperature. The wells are then washed five times with 100 μ l PBS binding buffer containing 0.1 % BSA (Sigma), 0.5% Tween 20 (Sigma), 50 μ M ZnCl₂ to remove unbound components. 25 Finally, the retained HRP activity is measured by adding 50 μ l QuantaBlu fluorogenic HRP substrate (Pierce). Figure 3 demonstrates the capture and detection of target site-binding zinc finger peptides using the assay described. Fluorescence is measured on a SpectraMax Gemini XS (Molecular Devices) fluorescence microplate reader at 320 nm excitation, 433 nm emission and 420 nm cut-off values.

30

The wells that give the highest levels of fluorescence are those which contain the highest number of, or tightest binding 3-finger peptides. PCR products from the second PCR

amplification stage, corresponding to such samples, are purified from TAE-agarose gels and quantified, as above. Pure PCR products are diluted to approximately 50 molecules per μl (which is equivalent to approximately 100 aM concentration) in 0.25 mg/ml λ DNA. As above, 1 μl samples of template are added to 1 ml PCR pre-mix containing
 5 primers, pETFwd4 and Rev3. 10 μl aliquots are placed in each well of a 96 well plate. At this stage, there is (on average) 0.5 template molecules per aliquot. Therefore, generally speaking, half of the samples will contain no template and half will contain a single template molecule. Samples are then PCR amplified using 30 cycles. Again, 0.5 μl PCR samples are taken from each well and amplified again by 30 cycles of PCR using
 10 the nested primers, pETFwd5 and Rev3. 1 μl of each of these PCR products is used for protein expression, as described above. At this stage, the highest levels of fluorescence correspond to the samples containing the tightest binding 3-finger peptides. The PCR product encoding such peptides is purified, as before, and can be sequenced to determine the protein sequence of the optimal 3-zinc finger domain for the appropriate binding site.

15

If further rounds of selection are required, PCR amplification can be conducted with the nested primers pETFwd6, pETFwd9 and pETFwd7, also shown below (Table 7).

NAME	SEQ ID NO	SEQUENCE
pETFwd1	2060	CGCTGACTTCCGCGTTTCC
pETFwd2	2061	TCCAGACTTTACGAAACACGG
pETFwd3	2062	CGAAGACCATTCATGTTGTTGC
pETFwd4	2063	GTCGCAGACGTTTTGCAGC
pETFwd5	2064	GCAGTCGCTTCACGTTTCGC
pETFwd6	2065	CGCTCGCGTATCGGTGATTC
pETFwd9	2066	CATTCTGCTAACCAGTAAGGC
pETFwd7	2067	GCCTAGCCGGGTCCTCAAC

20 **Table 7:** Primers used for PCR amplification of 3-finger cassettes (as constructed by the procedure of Figure 2) to provide template used in screening zinc finger libraries.

Protocol B:

The peptides of library 2 were screened to select 3-zinc finger domains which bind the sequences: 5'-GCG-TGG-GCG-3', and 5'-GGG-AGG-CCT-3'. Double stranded binding sites for use in the selection protocol were generated by annealing the complementary oligonucleotides: Zif.b site and Zif site RC (AS154 and AS155, shown above), which generated the 5'-GCG-TGG-GCG-3' binding site; and the oligonucleotides 5'-TTTTTTTTTTTGGGAGGCCTTTTTTTTTTTT-3' (SEQ ID NO:2123) and 5'-AAAAAAAAAAAGGCCTCCCAAAAAAAAAA-3' (SEQ ID NO:2124), which generated the 5'-GGG-AGG-CCT-3' binding site. The top strand of each binding site was biotinylated, allowing capture of binding site/zinc finger/HA-antibody ternary complexes onto streptavidin-coated plate in an ELISA screening assay.

15 The 3-finger library 2 constructs were cloned into the multiple cloning site of vector pET23a (Novagen), using appropriate restriction sites. This library was then transformed into *E.coli* and plated out to grow single colonies. 384 colonies (which should represent the vast majority of the 64 member library) were picked into 2xYT media with ampicillin and cultures grown at 37°C overnight. Library 2 expression cassettes were recovered from bacteria by PCR using primers pETFwdx (where x is 1-7, eg pETFwd1) and Rev3 as described in Protocol A above.

In vitro coupled transcription / translation of PCR products was conducted as described above, with the difference that each of the 384 zinc finger peptides was screened individually in a well of a 384 well plate. The library was screened against the 5'-GCG-TGG-GCG-3', and 5'-GGG-AGG-CCT-3' binding sites, as detailed in Protocol A. Wells that yielded the highest levels of fluorescence were those which contain the tightest binding 3-finger peptides. The ELISA results from the screen of the 384 samples against the 5'-GCG-TGG-GCG-3' site are shown in Figure 4. Six constructs displayed significant binding to the target site and these are termed C8, G16, I19, I23, J19 and K19 according to their coordinates on the 384-well plate. Similarly, one construct (B10)

showed strong binding to the 5'-GGG-AGG-CCT-3' target site. PCR products encoding the tightest binding peptides can be purified, as described *supra*, and sequenced.

5 Some of the selected constructs: C8, J19, K19, I23, G16 (which bind the 5'-GCG-TGG-GCG-3' site) and B10 (which binds the 5'-GGG-AGG-CCT-3' site), were selected and screened against a range of different binding sites to test their specificity. The sites used were: 5'-GCG-TGG-GCG-3'; 5'-CCA-CTC-GGC-3'; 5'-CCT-AGG-GGG-3'; 5'-GGA-TAA-GCG-3'; 5'-GGG-AGG-CCT-3'; 5'-GCG-TAA-GGA-3'; and 5'-GCG-GGG-
10 GGA-3'. The binding assay was conducted as described above. The results (Figure 5) show that the selected 3-zinc finger peptides bind preferentially to their target site, in comparison to the alternative binding sites tested.

**Example 5: Human Zinc Finger Module Libraries for Rapid Selection of 2-Finger
15 Units.**

The preferred subunits of a poly-zinc finger construction strategy are in the form of two-finger sub-domains. Assuming that there are 1,000 individual natural finger modules, a library of all combinations of such zinc finger modules, in 2-finger units, would contain
20 1,000,000 members. All of the 1,000 natural finger modules would have to be made from oligonucleotides, and the expense would be considerable. Furthermore, this figure is likely to be an underestimate of the number of natural fingers. Hence, due to the huge numbers of natural, human zinc finger modules available, it is advantageous to limit the size of the libraries screened, as discussed in the Description. One way in which library
25 size can be reduced is to limit the library members to zinc finger modules which are predicted to bind the desired sequence. For instance, based on the target sites in Example 1, if 2-finger domains are required to bind the sequence 5'-GCG-TGG-3', an individual library can be constructed from the zinc finger modules predicted to bind the sequences 5'-GCG-3' and 5'-TGG-3'. Equally, if the sequence 5'-GGA-TAA-3' is to be targeted,
30 zinc finger modules predicted to bind the sequences and 5'-GGA-3' and 5'-TAA-3' can be used. Table 8 shows the natural, human zinc finger modules from Example 1, which are predicted to bind the aforementioned 3 bp sequences.

5'-GCG-3'	5'-TGG-3'	5'-GGA-3'	5'-TAA-3'
Zif268 finger 1 (GCG)	Zif268 finger 2 (TGG)	BCL6 (NGA)	TYY1 (NAA)
Zif268 finger 3 (GCG)	MAZ finger 2 (TGG)	O75626 (GGA)	O15391 (YAA)
Sp1 finger 2 (GCG)	WT1 finger 3 (TGG)	ZN45 (N ^N /T _A)	O75626 (YAA)
WT1 finger 4 (GCG)	SP4 (NGG)	O15535 (GNA)	ZN45 (N ^N /T _A)
BTE1 (GCG)	BTE1 (NGG)	Q15776 (GNA)	Z136 (TNN)
O43296 (GNG)	Z136 (TNN)	O60893 (GNA)	Z239 (YAA)
Z174 (GCG, RNA)	Q15776 (NGG)	Z132 (a) (GGA)	Q15776 (a) (TNA)
Z202 (GCG, RNA)	ZN84 (YGG)	Z132 (b) (GGA)	Q15776 (b) (TNA)
		Z132 (GGN)	Z195 (YAA)
		ZN85 (GGA)	ZN84 (YAA)
			O75346 (TAA)
			ZN43 (TAA)

Table 8. The natural, human zinc finger modules predicted to bind the sequences 5'-GCG-3', 5'-TGG-3', 5'-GGA-3' and 5'-TAA-3'.

5

On the basis of the specificities shown in Table 5, a library of 2-finger units to target the 6 bp sequence 5'-GCG-TGG-3' has 64 (8x8) members, and a library to target the sequence 5'-GGA-TAA-3' has 120 (10x12) members. To screen sample sizes of this magnitude we can construct each 2-finger unit specifically (using for example, an 8x8 or 10x12 matrix arrangement), and assay the samples containing individual clones using the fluorescent-ELISA protocol of Example 4. Such a procedure can save time in comparison to constructing all possible 64 or 120 variants in a random fashion (as a library), as described in Example 4, because the number of constructs screened would have to be considerably higher.

15

a. Construction of 2-Finger Domains to Bind 5'-GCG-TGG-3'

A 64 member, 2-finger library is constructed from the natural, human zinc finger modules predicted to bind the sequences 5'-GCG-3' and 5'-TGG-3' (Table 8, columns 1 and 2).

20

The 2-finger library units are all attached to the C-terminus of Zif268 finger 1, which acts as an anchor finger. The construction protocol is different from that described in Example 4, as described below.

5 *Zinc Finger Cassettes*

Nucleotide sequences encoding the amino acid sequences of the 16 zinc finger modules (Table 8, columns 1 and 2) are determined, taking into account human codon preferences, and the corresponding nucleotide sequences are synthesised as single stranded oligonucleotides, shown in Table 9. Double stranded cassettes encoding the zinc finger modules and flanking linker sequences are generated by PCR using the appropriate primers, shown in Table 10.

X	FINGER	SEQ ID NO	NUCLEOTIDE SEQUENCE
1	Zif268 F1	2068	TACGCCTGCCCCGTGGAGAGCTGCGACCGCCGCTTCAG CCGCAGCGACGAGCTGACCCGCCACATCCGCATCCAC
2	Zif268 F3	2069	TTCGCCTGCGACATCTGCGGCCGCAAGTTCGCCCCGAG CGACGAGCGCAAGCGCCACACCAAGATCCAC
3	Sp1 F2	2070	TTCGCCTGCAGCTGGCAGGACTGCAACAAGAAGTTCGC CCGCAGCGACGAGCTGGCCCCGCCACTACCGCACCCAC
4	WT1 F4	2071	TTCAGCTGCCGCTGGCCCAGCTGCCAGAAGAAGTTCGC CCGCAGCGACGAGCTGGTGCGCCACCACAACATGCAC
5	BTE1	2072	TTCCCCTGCACCTGGCCCCGACTGCCTGAAGAAGTTCAG CCGCAGCGACGAGCTGACCCGCCACTACCGCACCCAC
6	O43296	2073	TACGAGTGCGTGGAGTGCGGCAAGGCCTTCACCCGCAT GAGCGGCCTGACCCGCCACAAGCGCATCCAC
7	Z174	2074	TACAAGTGCGACGACTGCGGCAAGAGCTTCACCTGGAA CAGCGAGCTGAAGCGCCACAAGCGCGTGCAC
8	Z202	2075	TACCGCTGCGACGACTGCGGCAAGCACTTCCGCTGGAC CAGCGACCTGGTGCGCCACCAGCGCACCCAC
9	Zif268 F2	2076	TTCCAGTGCCGCATCTGCATGCGCAACTTCAGCCGCAG CGACCACCTGAGCACCCACATCCGCACCCAC
10	MAZ F2	2077	TACAACTGCAGCCACTGCGGCAAGAGCTTCAGCCGCCC CGACCACCTGAACAGCCACGTGCGCCAGGTGCAC
11	WT1 F3	2078	TTCCAGTGCAAGACCTGCCAGCGCAAGTTCAGCCGCAG CGACCACCTGAAGACCCACACCCGCACCCAC
12	Sp4	2079	CACAAGTGCCCCCTACAGCGGCTGCGGCAAGGTGTACGG CAAGAGCAGCCACCTGAAGGCCCACTACCGCGTGCAC
13	BTE1	2080	CACAAGTGCCCCCTACAGCGGCTGCGGCAAGGTGTACGG CAAGAGCAGCCACCTGAAGGCCCACTACCGCGTGCAC

14	Z136	2081	TTCGAGTGCAAGCGCTGCGGCAAGGCCTTCCGCAGCAG CAGCAGCTTCCGCCTGCACGAGCGCACCCAC
15	Q15776	2082	TACGAGTGCGACGAGTGCGGCAAGACCTTCCGCCGCAG CAGCCACCTGATCGGCCACCAGCGCAGCCAC
16	ZN84	2083	TACGAGTGCGGCGAGTGCGGCAAGGCCTTCAGCCGCAA GAGCCACCTGATCAGCCACTGGCGCACCCAC

¹ RNA Binding.

Table 9. Nucleotide sequences of zinc finger modules and nucleotide sequences encoding other peptide sequences used in the construction of peptides to bind the sequence 5'-GCG-TGG-3'.

The primers used to amplify the N-terminal finger of the pair (the equivalent of cassette B, above) add TGEKP (SEQ ID NO:3) linker sequences, and the restriction site *Xma*I (5'-CCC-GGG-3') at the 5' end and an *Age*I site (5'-ACC-GGT-3') at the 3' end. *Age*I and *Xma*I create compatible ends, but have unique restriction sites. These primers are called CasBxFwd and CasBxRev, respectively, where x refers to the number of the zinc finger module in Table 9. The primers used to amplify the C-terminal finger of the pair (the equivalent of cassette C, above) add TGEKP (SEQ ID NO:3) linker sequences, and the restriction site *Xma*I at the 5' end and a sequence encoding LRQKDGGGS (SEQ ID NO:2125), containing a restriction site for *Bam*HI at the 3' end. These primers are referred to as CasCxFwd and CasCxRev, respectively. The 16 individual zinc finger cassettes are then purified using the QIAquick PCR purification kit (Qiagen).

Name	SEQ ID NO	Sequence
CasB9Fwd	2084	GATCCCCGGGGAGAGAAGCCCTTCCAGTGCCGCATCTGCAT
CasB10Fwd	2085	GATCCCCGGGGAGAGAAGCCCTACAACTGCAGCCACTGCGG
CasB11Fwd	2086	GATCCCCGGGGAGAGAAGCCCTTCCAGTGCAAGACCTGCCA
CasB12Fwd	2087	GATCCCCGGGGAGAGAAGCCCCACAAGTGCCCCCTACAGCG
CasB13Fwd	2088	GATCCCCGGGGAGAGAAGCCCCACAAGTGCCCCCTACAGCG
CasB14Fwd	2089	GATCCCCGGGGAGAGAAGCCCTTCGAGTGCAAGCGCTGCG
CasB15Fwd	2090	GATCCCCGGGGAGAGAAGCCCTACGAGTGCGACGAGTGCG
CasB16Fwd	2091	GATCCCCGGGGAGAGAAGCCCTACGAGTGCGGCGAGTGCG
CasC1Fwd	2092	GATCCCCGGGGAGAGAAGCCCTACGCCTGCCCCGTGGAG

CasC2Fwd	2093	GATCCCCGGGGAGAGAAGCCCTTCGCCTGCGACATCTGCG
CasC3Fwd	2094	GATCCCCGGGGAGAGAAGCCCTTCGCCTGCGAGCTGGCAGG
CasC4Fwd	2095	GATCCCCGGGGAGAGAAGCCCTTCAGCTGCCGCTGGCCC
CasC5Fwd	2096	GATCCCCGGGGAGAGAAGCCCTTCCCCTGCACCTGGCCC
CasC6Fwd	2097	GATCCCCGGGGAGAGAAGCCCTACGAGTGCGTGAGTGCG
CasC7Fwd	2098	GATCCCCGGGGAGAGAAGCCCTACAAGTGCGACGACTGCGG
CasC8Fwd	2099	GATCCCCGGGGAGAGAAGCCCTACCGCTGCGACGACTGCG
CasB9Rev	2100	CTTCTCACC GGT GTGGGTGCGGATGTGGGTG
CasB10Rev	2101	CTTCTCACC GGT GTGCACCTGGCGCACGTG
CasB11Rev	2102	CTTCTCACC GGT GTGGGTGCGGGTGTGGGT
CasB12Rev	2103	CTTCTCACC GGT GTGCACGCGGTAGTGGGC
CasB13Rev	2104	CTTCTCACC GGT GTGCACGCGGTAGTGGGC
CasB14Rev	2105	CTTCTCACC GGT GTGGGTGCGCTCGTGCA
CasB15Rev	2106	CTTCTCACC GGT GTGGCTGCGCTGGTGGCC
CasB16Rev	2107	CTTCTCACC GGT GTGGGTGCGCCAGTGGCT
CasC1Rev	2108	GATCGGATCCGCCGCCGTCCTTCTGGCGCAGGTGGATGC GGATGTGGCGG
CasC2Rev	2109	GATCGGATCCGCCGCCGTCCTTCTGGCGCAGGTGGATCT TGGTGTGGCGC
CasC3Rev	2110	GATCGGATCCGCCGCCGTCCTTCTGGCGCAGGTGGGTGC GGTAGTGGCG
CasC4Rev	2111	GATCGGATCCGCCGCCGTCCTTCTGGCGCAGGTGCATGT TGTGGTGGCGC
CasC5Rev	2112	GATCGGATCCGCCGCCGTCCTTCTGGCGCAGGTGGGTGC GGTAGTGGCG
CasC6Rev	2113	GATCGGATCCGCCGCCGTCCTTCTGGCGCAGGTGGATGC GCTTGTGGCGG
CasC7Rev	2114	GATCGGATCCGCCGCCGTCCTTCTGGCGCAGGTGCACGC GCTTGTGGCG
CasC8Rev	2115	GATCGGATCCGCCGCCGTCCTTCTGGCGCAGGTGGGTGC GCTGGTGGCG

ScaRev	2116	GTCATGCCATCCGTAAGATGC
GSFwd	2117	<u>GGCGGATCCTATCCGTATGATGTG</u>
Zif1Fwd	2118	AGAGAGAGAGAGATCTATGGCGGAAGAACGTCCGTATGC GTGCCCCGGTGGAAAG
Zif1Rev	2119	AGCCGGATCCCAAACACCGGTATGAATACGAATATGACG GG
pETRev1	2120	AGTGTAGCGGTCACGCTGC

Table 10. Oligonucleotides used for PCR construction of rapid zinc finger library. Annealing sequences are shown in bold, restriction sites are underlined.

5 ***3-Finger Library Peptides***

The 2 natural zinc finger modules for each construct are appended to the C-terminus of Zif268 finger 1 (as in Example 4, library 2). Hence, a plasmid construct containing Zif268 finger 1 and appropriate restriction sites for cloning of the two natural finger
10 modules is also prepared. The construction and cloning procedure for the 3-finger libraries follows (see also Figure 6).

(a) The plasmid pET23a/TZF-HA was assembled by PCR amplification of plasmid pTFZ-KOX (described in co-owned WO 01/53480) with primers AS1 and AS2.
15 The sequences of these primers are as follows:

AS1: CGATGGATCCATGGGAGAGAAGGCGCTGC (SEQ ID NO:2126)
AS2: GCGTAAAGCTTACGCATAATCCGGCACATCATACGGATAAGAG
 CCGCCGCCGTCCTTCTGTCTTAAATGGATTT (SEQ ID NO:2127)

The PCR product was gel purified and digested with BamHI and HindIII, then
20 repurified and cloned into BamH I/Hind III-digested pET23a vector (Novagen), yielding pET23a/TFZ-HA. A number of clones were picked and sequenced to verify the correctness of the inserts.

(b) A fragment of approximately 1.2 kb is amplified from the vector
25 pET23a/TFZ-HA, using the primers ScaRev and GSFwd (Table 10). This fragment

contains the HA-epitope tag sequence (YPYDVPDYA* (SEQ ID NO: 2122)) and part of the GGGG (SEQ ID NO:1988) linker sequence at the 5' end. Additionally, the GSFwd primer adds a BamHI site at the extreme 5' end. The ScaRev primer does not contain a restriction site, but a *ScaI* site from the vector is present approximately 40 bp downstream of the primer binding site. This fragment is cut with *Bam*HI and *Sca*I and inserted into similarly cut pET23a.

(c) Zif268 finger 1 is then amplified using the PCR primers Zif1Fwd and Zif1Rev (Table 10), which add a *Bgl*II site at the 5' end and both *Age*I and *Bam*HI sites at the 3' end. This construct is then cut with *Bgl*II and *Bam*HI and inserted into the vector construct made in step (b), which has been linearised with *Bam*HI. At this stage the new construct, termed pET23aZif1HA is sequenced to find correctly oriented zinc finger inserts.

(d) Oligonucleotides encoding zinc finger modules for the C-terminus of the 3-finger constructs (cassette C) are amplified using the primers CasCxFor and CasCxRev (where x is 1 to 8, see Table 10). These cassettes are then digested with the restriction enzyme *Bam*HI, and inserted into *Bam*HI cut, dephosphorylated pET23aZif1HA. At this stage the new vector construct is not recircularised.

(e) Oligonucleotides encoding zinc finger modules for cassette B are amplified using primers CasBxFor and CasBxRev (where x is 9 to 16, see Table 10). These fragments are cut with the enzymes *Xma*I and *Age*I, at 37 °C for 1-2 hours. The linear vector produced in stage (d) above, is also cut with *Age*I and *Xma*I (as described), and dephosphorylated. Digested cassette B fragments are ligated into *Age*I, *Xma*I cut vector, in the presence of the restriction enzymes *Age*I and *Xma*I at room temperature for 16 hours. During this incubation incorrectly ligated fragments are re-digested and re-ligated repeatedly, until the majority (or all) of the inserts are in the desired orientation. Correct 3-finger constructs have the assembly depicted in Figure 6.

(f) Finally, 3-finger constructs are amplified from the ligated vector (produced in step (e)) using the primers pETFwd1 (Table 5) and pETRev1 (Table 10). 1 µl of each

ligation mixture is amplified in a 10 µl (total volume) PCR reaction for 30 cycles. Alternatively, the ligated vector can be transformed into bacteria to produce samples containing single zinc finger clones.

- 5 The above procedure results in the majority of PCR products being the correct 3-finger constructs, so that any incorrect fragments will not significantly affect the selection protocol, and the PCR products can be used for screening without further processing. Alternatively, 3-finger PCR products may be purified from an agarose gel before use.

10 **b. Screening of the Library Against 5'-GCG-TGG-GCG-3'**

Members of the zinc finger library can be screened against the desired target site from a mixed population of clones, or from individual clones as described in Example 4, Protocol A or Protocol B (above), respectively. The target site for the screen is produced
15 by annealing the oligonucleotides Zif.b site (AS154) and Zif site RC (AS155), as before. Template for protein expression is in each case made by PCR using primers pETFwd1 (Table 5) and pETRev1 (Table 10). 1 µl of each PCR reaction is used to express protein and screen for binding to the Zif site in the manner described in Example 4. The DNA corresponding to the samples giving the highest fluorescence signals is collected, purified
20 from a 1% TAE-agarose gel, and sequenced to determine the sequence of the optimal binding 3-finger peptide.

Example 6: Reduced Human Zinc Finger Module Library for Universal DNA Recognition.

25

A library system similar to that described in Example 5 can be constructed using zinc finger modules from databases such as those in Examples 1, 2 and 3 to select 2-finger units which bind any 2-finger (6 bp) recognition sequence. There are only 4096 ($=4^6$) unique 6 bp sequences, therefore, a 2-finger library of natural zinc fingers (from specific
30 animals, plants or fungi) can easily be constructed with enough variability to provide a specific 2-finger combination for optimal binding to any 6 bp target site. Again, to reduce the number of natural zinc finger modules that have to be constructed, a small

selection of natural zinc finger modules (*e.g.*, 3) are chosen for each 3 bp binding sequence (according to their predicted or determined recognition sequence). There are 64 ($=4^3$) possible 3 bp binding sequences so in the first instance less than 200 (*i.e.* 192) natural zinc finger modules are constructed. These 200 zinc finger modules can be in either of 2 possible positions in the 2-finger construct, which gives approximately 40,000 ($=200^2$) combinations of fingers to bind the 4096 possible 6 bp target sites. As in Example 5, these 2-finger units are attached to Zif268 finger 1 which acts as an anchor for DNA recognition.

10 a. Library Construction

The selected zinc finger modules are reverse translated from their amino acid sequences and synthesised as oligonucleotides. Double stranded zinc finger cassettes for both N-terminal and C-terminal fingers are created by PCR using primers specific for the relevant zinc finger module. Each zinc finger module is amplified in 2 separate reactions, as described in Example 5. The first PCR reaction uses primers which add TGEKP (SEQ ID NO:3) linker peptides and *Age*I and *Xma*I restriction sites, to the 3' and 5' ends, respectively, to generate cassette B fragments. The second PCR reaction generates cassette C fragments by adding a TGEKP (SEQ ID NO:3) linker and an *Xma*I site at the 5' end (this primer is the same as that used in cassette B production), and a sequence encoding the sequence LRQKDGGGS (SEQ ID NO:2125) and a *Bam*HI restriction site at the 3' end. The final constructs are similar to that represented in Figure 6.

25 b. Library Selection

The collection of 3-finger zinc finger peptides produced above can be used to obtain specific domains for binding desired target sequences. Two exemplary approaches are described below.

30 i). Non-Cloning Selections.

A library constructed as described herein can be used to select optimal zinc finger domains for binding to any specified binding site. For instance, to select a peptide which binds the sequence 5'-GGA-TAA-3', the binding site formed by annealing the oligonucleotides #1#5#6.b and #1#5#6 RC (Table 6, above), can be used as a target site
5 (5'-GGA-TAA-GCG-3'). Selection of a zinc finger domain to bind such a target can be conducted, for example, in the manner described in Example 4. Briefly, the zinc finger library is diluted into 100 or more sub-libraries, which are screened as described above. The most active sub-libraries collected are further diluted to create much smaller sub-libraries, which are screened again, and so on. Following such a protocol, a library of
10 40,000 members can be fully screened and a high-affinity binder selected in just 3 rounds.

This selection procedure provides an extremely rapid method to select zinc finger peptides to bind any desired target site. The procedure also has the advantages of eliminating the need for cloning (as is required for methods such as phage display, see
15 below), and is not limited by library size.

ii). Phage Library Selections

Zinc finger polypeptide phage display libraries are made and used to select clones
20 encoding peptides that bind the desired nucleotide sequence, as described in co-owned WO 98/53057. An exemplary phage display library contains peptides which bind target sites with the sequence 5'-XXX-XXX-GCG-3', where X can be any nucleotide. Hence, libraries of phage can be selected using the same target sites as described above. The selection protocol for zinc fingers displayed on phage is briefly described below.

25

Protocol

The selection protocol is adapted from that described in co-owned international patent application WO98/53057.

30

The 3-finger constructs of the present Example are PCR amplified using universal forward and reverse primers which contain sites for *NotI* and *SfiI* respectively (called NatPhageF and NatPhageR, respectively).

5 NatPhageF: GCAACTGCGGCCCGAGCCGGCCATGGCAGAGGAACGCCCCGTATG (SEQ ID NO:2128)

NatPhageR: GAGTCATTCTGCGGCCGCGTCCTTCTGGCGCAGGTG (SEQ ID NO:2129)

Backward PCR primers in addition introduce Met-Ala-Glu as the first three amino acid
 10 residues of the zinc finger polypeptides, and these are followed by the residues of the wild type or library zinc finger polypeptides as required. Cloning overhangs are produced by digestion with *SfiI* and *NotI* where necessary. Nucleic acid encoding zinc finger polypeptide fragments is ligated into similarly prepared Fd-Tet-SN vector. This is a derivative of fd-tet-DOG1 (Hoogenboom *et al.* (1991) *Nucl. Acids Res.* 19:4133-4137),
 15 in which a section of the pelB leader and a restriction site for the enzyme *SfiI* (underlined) have been added by site-directed mutagenesis using the oligonucleotide:

5' CTCCTGCAGTTGGACCTGTGCCATGGCCGGCTGGGCCGCATA
 GAATGGAACAACCTAAAGC 3' (SEQ ID NO:2130)

20 that anneals in the region of the polylinker. Electrocompetent DH5 α cells are transformed with recombinant vector in 200 ng aliquots, grown for 1 hour in 2xTY medium with 1% glucose, and plated on TYE containing 15 μ g/ml tetracycline and 1% glucose.

25 To generate phage for selections, tetracycline resistant colonies are transferred from plates into 2xTY medium (16g/litre Bacto tryptone, 10g/litre Bacto yeast extract, 5g/litre NaCl) containing 50 μ M ZnCl₂ and 15 μ g/ml tetracycline, and cultured overnight at 30°C in a shaking incubator. Cleared culture supernatant containing phage particles is obtained by centrifuging at 300 xg for 5 minutes.

Double stranded binding sites for use in selections are generated by annealing complementary oligonucleotides, one of which is biotinylated.

Biotinylated DNA target sites (1 pmol) are bound to streptavidin-coated wells (Roche).

- 5 Phage supernatant solutions are diluted 1:10 in PBS selection buffer (PBS containing 50 μ M $ZnCl_2$, 2% Marvel, 1% Tween, 20 μ g/ml sonicated salmon sperm DNA, and 10-fold excess of competitor DNA), and 200 μ l is applied to each well for 1 hour at 20°C. After this time, the wells are emptied and washed 18 times with PBS containing 50 μ M $ZnCl_2$ and 1% Tween and 2 times in PBS containing 50 μ M $ZnCl_2$. Retained phage are eluted in
- 10 100 μ l 0.1M triethylamine and neutralised with an equal volume of 1M Tris (pH 7.4). Logarithmic-phase *E. coli* JM109 (100 μ l) are infected with eluted phage (100 μ l), and used to prepare phage supernatants for subsequent rounds of selection. After 4 rounds of selection, a 'pool' or 'mini-population' of phage is obtained, which bind the specified target sequence. These pools of phage can be stored at -70°C for later use. Additionally,
- 15 *E. coli* infected with these phage pools can be plated to obtain individual clones, which can be tested by ELISA for binding affinity and specificity to obtain the 'best' clone (see Example 9, Quality Control).

20 **Example 7: Complete Human Zinc Finger Module Library for Universal DNA Recognition.**

- An complete, or nearly complete, library containing all zinc finger sequences which bind a particular target site can be constructed using zinc finger modules to select 2-finger (or
- 25 3-finger) units which bind any 6 bp (or 9 bp) recognition sequence. Two exemplary methods for construction of such a library are described.

a. Oligonucleotide-Based Library Construction.

- 30 All zinc finger modules may be synthesised as a single stranded oligonucleotide, as described in Example 4. Zinc finger modules are made double stranded and TGEKP (SEQ ID NO:3) linkers added by PCR with 5' and 3' primers specific for each individual

zinc finger module, to make cassettes. These cassettes can then be recombined, as described in Example 5, to make random or deliberate combinations of zinc finger modules comprising 2, 3, or more linked fingers.

5 **b. PCR-Based Library Construction.**

Zinc fingers proteins (especially of the Cys₂His₂ family) form the second most abundant family of proteins in the human genome. Furthermore, in nature, zinc finger modules are often linked by the canonical linker peptide TGEKP (SEQ ID NO:3), which begins
10 immediately after the second zinc-coordinating histidine residue. Therefore, the peptide sequence HTGEKP (SEQ ID NO:2131) is commonly found between natural zinc finger modules. Because of this consensus sequence, it has been possible to clone natural zinc finger modules from the human genome (Becker, K.G., Nagel, J.W., Canning, R.D., Biddison, W.E., Ozato, K. & Drew, P.D. (1995) *Hum. Mol. Genet.* 4: 685-691; Bray, P.,
15 Lichter, P., Thiesen, H.-J., Ward, D.C. & Dawid, I.B. (1991) *Proc. Natl. Acad. Sci. USA* 88: 9563-9567), and the Arabidopsis genome (Meissner, R. & Michael, A.J. (1997) *Plant Mol Biol* 33: 615-624), using redundant primers for PCR. See also Pellegrino *et al.* (1991) *Proc. Natl. Acad. Sci. USA* 88:671-675. It is preferable to use genomic DNA or a genomic DNA (gDNA) library, rather than a cDNA library, because transcription factors,
20 such as zinc finger proteins, are strongly regulated during the cell cycle, development and in response to extracellular signals. Hence, a cDNA library will probably not contain the majority of zinc finger proteins, and will be biased towards highly expressed proteins.

A suitable protocol for the PCR-extraction of zinc finger modules from human genomic
25 DNA follows:

Genomic DNA is purified directly from human cells, or provided by a gDNA library. gDNA libraries are preferable as they are commercially available (for example from Clontech, ATCC, Stratagene etc) and can be easily manipulated. PCR to extract zinc
30 finger modules can be conducted directly on purified gDNA, or the gDNA library can be screened for zinc fingers containing the HTGEKP (SEQ ID NO:2131) motif before carrying out PCR. To screen the gDNA library, any method known to one of skill in the

art, *e.g.* colony hybridisation, can be used. Phage containing gDNA inserts are plated onto *Escherichia coli* XL-1 Blue bacterial lawns. At least 10^6 phage plaques are transferred to replica filters and screened with, for example, a 27-mer ^{32}P -radiolabelled degenerate oligonucleotide, which anneals to the conserved linker region of zinc finger proteins and adjacent sequences. The sequence of a suitable degenerate probe (SEQ ID NO:2132), and the amino acid sequence (SEQ ID NO:2133) to which it corresponds is shown below.

10 $\text{C}^{\text{G}}/\text{T}^{\text{C}}/\text{G}$ $\text{A}^{\text{T}}/\text{C}^{\text{C}}/\text{G}$ $\text{CA}^{\text{C}}/\text{T}$ $\text{AC}^{\text{C}}/\text{G}$ $\text{GG}^{\text{C}}/\text{G}$ $\text{GA}^{\text{G}}/\text{A}$ $\text{AA}^{\text{G}}/\text{A}$ $\text{CC}^{\text{C}}/\text{T}$ $\text{T}^{\text{A}}/\text{T}^{\text{C}}/\text{T}$
 R/L I/T/M H T G E K P Y/F

Hybridisation is performed, *e.g.*, for 16 hours at 42-50 °C, following which filters are washed 3-5 times, to remove non-specifically bound probe, in 0.2x standard saline citrate (SSC)/0.1% SDS. Filters are then subjected to autoradiography or phosphorimaging to determine positive plaques.

Positive plaques are picked into log-phase *E. coli* XL-1 Blue bacterial cultures and the phage are harvested for PCR. 1 µl phage supernatant is added to 49 µl PCR pre-mix, containing the oligonucleotide primers TGEKPfor (SEQ ID NO:2134) and TGEKPrev (SEQ ID NO:2135) (shown below, annealing sequence in bold), and zinc finger modules are amplified by 30 cycles of PCR. TGEKPfor (SEQ ID NO:2134) and TGEKPrev (SEQ ID NO:2135) also contain *Xba*I and *Eco*RI restriction sites (underlined), respectively. PCR products are separated on 1.5% TAE-agarose gels and fragments of approximately 120 bp (corresponding to 1 zinc finger module plus flanking sequences) are purified, as described in Example 4. Additionally, fragments of approximately 220 bp, corresponding to natural 2-finger units, can also be collected and used. Such products can be digested with *Xba*I and *Eco*RI and cloned into a vector that has been digested so as to generate compatible ends, such as, for example, pcDNA3.1(-) (Invitrogen) digested with *Eco*RI and *Xba*I. Such a vector pool can then be used as a source for natural 1- or 2-zinc finger modules, from which to construct 2- or 3-zinc finger peptides for selections as described above. Zinc finger modules for cassette B can be amplified from such vectors using the universal primers TGEKPXma (SEQ ID NO:2136) and TGEKPAge (SEQ ID NO:2137),

which anneal to the conserved TGEKP (SEQ ID NO:3) linker regions and add restriction sites for the enzymes *Xma*I at the 5' terminus and *Age*I at the 3' terminus, respectively (restriction sites underlined). Cassette C units can be amplified using the primer TGEKPXma (SEQ ID NO:2136) and TGEKPend (SEQ ID NO:2138), which adds a 3' TRQKDGGGS (SEQ ID NO:2139) sequence incorporating a *Bam*HI site (underlined, see below). Two- and 3-finger constructs can then be constructed and screened as described in the Examples above.

TGEKPfor: TTAGTCTAGA^C/GCA^C/TAC^C/GGG^C/GGA^G/AA^G/ACC (SEQ ID NO:2134)

TGEKPrev: TACTGAATTC^G/AGG^C/TTT^C/TTC^G/CCG^G/CGT^G/ATG (SEQ ID NO:2135)

TGEKPXma: TCTAGA^C/GCA^C/TCCCGGGGA^G/AA^G/ACC (SEQ ID NO:2136)

TGEKPage: GAATTC^G/AGG^C/TTT^C/TCAACCGGT^G/ATG (SEQ ID NO:2137)

TGEKPend: AGTGTGGTGGGAATTC^G/AGGGGATCCGCCCGCCGTC^C/TTT^C/TTG^G/CCG^G/CGT^G/ATG (SEQ ID NO:2138)

Example 8. Microarray Analysis.

Microarray analysis can also be used to determine the binding site specificity of 2- and 3-finger peptides. For example, a 3-zinc finger library, with finger 1 fixed as Zif268 finger one recognises the sequence 5'-XXX-XXX-GCG-3', where X is any specified nucleotide. Hence, there are 4096 (=4⁶) unique binding sites for such a library. All 4096 of these sites can be arrayed onto a single glass slide, allowing a specified 2-finger peptide to be screened against every possible binding site at once. A suitable protocol for such an experiment is described in Martha L. Bulyk, Xiaohua Huang, Yen Choo, & George M. Church (*Proc. Natl. Acad. Sci. USA*: Vol. 98, No. 13, 7158-7163, June 19, 2001) which is incorporated, by reference, in its entirety. See also co-owned WO 01/25417, the disclosure of which is hereby incorporated by reference in its entirety.

The amount of binding to each target sequence can be visualised and quantified using simple fluorescence measurements. For example, the zinc finger peptide can be expressed *in vitro*, or on the surface of phage. Isolated zinc finger peptides may contain an epitope tag for labelling purposes, whereas bound phage can be detected using a
5 primary antibody against a phage coat protein, such as gVIII. A secondary antibody, such as one conjugated to R-phycoerythrin may be used to provide a visible signal when a suitable substrate is applied.

10 **Example 9. Quality Control.**

Particular 2- or 3-finger peptides can be screened to determine their specificity or affinity, as desired.

a. Phage ELISA Assay

Phage supernatants from Round 4 of selection (Example 6, *supra*) are used to infect *E. coli* JM109 bacteria, and grown to prepare fresh supernatants for zinc finger phage ELISA, using standard procedures as described previously (Choo, Y. & Klug, A. (1994) *Proc. Natl. Acad. Sci. USA* 91, 11163-11167; Choo, Y. & Klug, A. (1994) *Proc. Natl. Acad. Sci. USA* 91, 11168-11172). Briefly, 5'-biotinylated, positionally randomised oligonucleotide libraries, containing Zif268 binding site variants, are synthesised by annealing complimentary oligonucleotides as described *supra*. DNA libraries are added to streptavidin-coated ELISA wells (Boehringer-Mannheim) in PBS containing 50µM ZnCl₂ (PBS/Zn). Phage solution (overnight bacterial culture supernatant diluted 1:10 in PBS/Zn containing 2% Marvel, 1% Tween and 20µg/ml sonicated salmon sperm DNA) is applied to each well (50µl/well). Binding is allowed to proceed for one hour at 20°C. Unbound phage are removed by washing 7 times with PBS/Zn containing 1% Tween, then 3 times with PBS/Zn. Bound phage are detected by ELISA using horseradish peroxidase-conjugated anti-M13 IgG (Pharmacia Biotech) and the colourimetric signal is quantitated using SOFTMAX 2.32 (Molecular Devices).

For rapid validation, the entire population of phage from Round 4 selection can be assayed in two ELISA wells: one containing the target DNA binding site, and one containing a control DNA binding site with between 1 and 5 base changes from the target sequence. A selection is deemed to be successful if the ELISA signal (representing DNA binding) is higher in the target well than in the control well.

The higher the signal measured above, the greater the *population* of specific binding clones. However, individual low values for such a procedure do not necessarily indicate a failure of the selection, as there may be individual high affinity / specificity clones within the round 4 phage population that may be masked by other non-specific clones. Nevertheless, this assay provides a quick profile of the overall quality of selection.

For a more detailed validation, individual phage clones are recovered from Round 4 by plating out infected bacterial colonies on agar. Fresh phage supernatants are prepared from these colonies and assayed by ELISA, as described above.

- 5 Finally, the coding sequence of individual zinc finger clones can be amplified by PCR using external primers complementary to phage sequence, and the PCR products are then sequenced to determine the amino acid sequence of the selected zinc fingers.

As an alternative, individual 3-finger peptides can be analysed by gel-shift assays or by
10 microarray screening, as described *infra*. See also WO 00/41566, WO 00/42219 and WO 01/25417.

b. Gel-Shift Assay

Peptides are assayed using ³²P end-labelled synthetic oligonucleotide duplexes containing
15 the appropriate binding site sequences.

DNA binding reactions contain the appropriate zinc-finger peptide, binding site and 1 µg competitor DNA (*e.g.*, poly dI-dC or salmon sperm DNA) in a total volume of 10 µl, which contains: 20 mM Bis-tris propane (pH 7.0), 100 mM NaCl, 5 mM MgCl₂, 50 µM ZnCl₂, 5 mM DTT, 0.1 mg/ml BSA, 0.1% Nonidet P40. Incubations are performed at
20 room temperature for 1 hour.

To determine the concentration of zinc finger peptide produced in the *in vitro* expression system, crude protein samples are used in gel-shift assays against a dilution series of the appropriate binding site. Binding site concentration is always well above the K_d of the peptide, but ranged from a higher concentration than the peptide (80 mM), at which all
25 available peptide binds DNA, to a lower concentration (3-5 mM), at which all DNA is bound. Controls are carried out to ensure that binding sites are not shifted (*i.e.*, bound) in the absence of zinc finger peptide. The reaction mixtures are then separated on a 7% native polyacrylamide gel. Radioactive signals are quantitated by PhosphorImager

analysis to determine the amount of shifted binding site, and hence, the concentration of active zinc finger peptide.

Dissociation constants (K_d) are determined in parallel to the calculation of active peptide concentration. For determination of K_d , serial 3, 4 or 5-fold dilutions of crude peptide are
5 made and incubated with radiolabelled binding site (10 pM – 10 nM depending on the peptide), as above. Samples are run on 7% native polyacrylamide gels and the radioactive signals quantitated by PhosphorImager analysis. The data is then analysed according to linear transformation of the binding equation and plotted in CA-Cricket Graph III (Computer Associates Inc. NY) to generate the apparent dissociation constants.
10 The K_d values reported are the average of at least two separate determinations.

c. Microarray Assay

Selected zinc finger domains can also be assayed for binding site specificity using the
15 microarray analysis outlined in Example 8.

All publications mentioned in the above specification are herein incorporated by reference. Various modifications and variations of the described methods and system of the invention will be apparent to those skilled in the art without departing from the scope
20 and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are apparent to those skilled in molecular biology or related fields are intended to be within the scope of
25 the following claims.

CLAIMS

1. A composite binding polypeptide comprising a first natural binding domain derived from a first natural binding polypeptide, and a second natural binding domain derived from a second natural binding polypeptide, wherein said first and second natural binding polypeptides may be the same or different; which polypeptide binds to a target, said target differing from the natural target of the both the first and the second binding polypeptides.
2. A composite polypeptide according to claim 1, wherein said first and second natural binding polypeptides are different polypeptides.
3. A composite polypeptide according to claim 1 or claim 2, comprising three or more natural binding domains.
4. A composite polypeptide according to any preceding claim, wherein the binding domains are nucleic acid binding domains.
5. A composite polypeptide according to claim 4, which is a nucleic acid binding polypeptide.
6. A composite polypeptide according to claim 4 or claim 5 which is a zinc finger polypeptide, and the natural binding domains are zinc finger domains.
7. A composite polypeptide according to claim 6, which comprises a Cys2-His2 zinc finger binding domain.
8. A composite polypeptide according to claim 6 or claim 7, which comprises a Cys3-His zinc finger binding domain.
9. A composite polypeptide according to any preceding claim, which comprises 6 or more natural binding domains.

10. A composite polypeptide according to claim 9, wherein 6 natural binding domains are arranged in a 3x2 conformation, separated by linker sequences.
11. A chimeric polypeptide comprising:
 - (a) a binding polypeptide according to any preceding claim, and
 - (b) a biological effector domain.
11. A library of natural binding domains.
12. A library according to claim 11, comprising a plurality of natural binding domains from which a polypeptide according to any one of claims 1 to 10 can be assembled.
13. A library of natural zinc finger nucleic acid binding domains, wherein said zinc finger domains comprise a linker attached thereto.
14. A library according to claim 13, wherein the linker comprises the sequence TGEKP.
15. A method for selecting a binding polypeptide capable of binding to a target site, comprising:
 - (a) providing a library of natural binding domains;
 - (b) assembling two or more of said domains to form a composite polypeptide;
 - (c) screening said composite polypeptide against the target site in order to determine its ability to bind the target site.
16. A method according to claim 15, wherein the natural binding domains are zinc finger binding domains.
17. A method according to claim 15 or claim 16, wherein two or more composite polypeptides comprising two or more domains which are selected for binding to two or

more target sites are combined to provide a composite polypeptide which binds to an aggregate binding site comprising the two or more target binding sites.

18. A method for designing a composite binding polypeptide, comprising:
 - (a) providing information defining a target site;
 - (b) selecting, from a database of natural binding domains, sequences of binding domains which are predicted to bind to the target site by the application of one or more rules which define target binding interactions for the binding domains; and
 - (c) displaying the sequences of the binding domains, separated by linker sequences, and optionally assembling the binding polypeptide from a library of said domains.
19. A method according to claim 18, wherein the binding domains are zinc finger domains.
20. A method according to claim 19, wherein the zinc fingers are considered to bind to a nucleic acid triplet and domains are selected according to one or more of the following rules:
 - (a) if the 5' base in the triplet is G, then position +6 in the α -helix is Arg; or position +6 is Ser or Thr and position ++2 is Asp;
 - (b) if the 5' base in the triplet is A, then position +6 in the α -helix is Gln and ++2 is not Asp;
 - (c) if the 5' base in the triplet is T, then position +6 in the α -helix is Ser or Thr and position ++2 is Asp;
 - (d) if the 5' base in the triplet is C, then position +6 in the α -helix may be any amino acid, provided that position ++2 in the α -helix is not Asp;
 - (e) if the central base in the triplet is G, then position +3 in the α -helix is His;
 - (f) if the central base in the triplet is A, then position +3 in the α -helix is Asn;
 - (g) if the central base in the triplet is T, then position +3 in the α -helix is Ala, Ser or Val; provided that if it is Ala, then one of the residues at -1 or +6 is a small residue;

(h) if the central base in the triplet is C, then position +3 in the α -helix is Ser, Asp, Glu, Leu, Thr or Val;

(i) if the 3' base in the triplet is G, then position -1 in the α -helix is Arg;

(j) if the 3' base in the triplet is A, then position -1 in the α -helix is Gln;

(k) if the 3' base in the triplet is T, then position -1 in the α -helix is Asn or Gln;

(l) if the 3' base in the triplet is C, then position -1 in the α -helix is Asp.

21. A method according to claim 19, wherein the zinc fingers are considered to bind to a nucleic acid quadruplet and domains are selected according to one or more of the following rules:

(a) if base 4 in the quadruplet is G, then position +6 in the α -helix is Arg or Lys;

(b) if base 4 in the quadruplet is A, then position +6 in the α -helix is Glu, Asn or Val;

(c) if base 4 in the quadruplet is T, then position +6 in the α -helix is Ser, Thr, Val or Lys;

(d) if base 4 in the quadruplet is C, then position +6 in the α -helix is Ser, Thr, Val, Ala, Glu or Asn;

(e) if base 3 in the quadruplet is G, then position +3 in the α -helix is His;

(f) if base 3 in the quadruplet is A, then position +3 in the α -helix is Asn;

(g) if base 3 in the quadruplet is T, then position +3 in the α -helix is Ala, Ser or Val; provided that if it is Ala, then one of the residues at -1 or +6 is a small residue;

(h) if base 3 in the quadruplet is C, then position +3 in the α -helix is Ser, Asp, Glu, Leu, Thr or Val;

(i) if base 2 in the quadruplet is G, then position -1 in the α -helix is Arg;

(j) if base 2 in the quadruplet is A, then position -1 in the α -helix is Gln;

(k) if base 2 in the quadruplet is T, then position -1 in the α -helix is His or Thr;

(l) if base 2 in the quadruplet is C, then position -1 in the α -helix is Asp or His;

(m) if base 1 in the quadruplet is G, then position +2 is Glu;

(n) if base 1 in the quadruplet is A, then position +2 Arg or Gln;

(o) if base 1 in the quadruplet is C, then position +2 is Asn, Gln, Arg, His or Lys;

(p) if base 1 in the quadruplet is T, then position +2 is Ser or Thr.

22. A method according to claim 19, wherein the zinc fingers are considered to bind to a nucleic acid quadruplet and domains are selected according to one or more of the following rules:

(a) if base 4 in the quadruplet is G, then position +6 in the α -helix is Arg; or position +6 is Ser or Thr and position ++2 is Asp;

(b) if base 4 in the quadruplet is A, then position +6 in the α -helix is Gln and ++2 is not Asp;

(c) if base 4 in the quadruplet is T, then position +6 in the α -helix is Ser or Thr and position ++2 is Asp;

(d) if base 4 in the quadruplet is C, then position +6 in the α -helix may be any amino acid, provided that position ++2 in the α -helix is not Asp;

(e) if base 3 in the quadruplet is G, then position +3 in the α -helix is His;

(f) if base 3 in the quadruplet is A, then position +3 in the α -helix is Asn;

(g) if base 3 in the quadruplet is T, then position +3 in the α -helix is Ala, Ser or Val; provided that if it is Ala, then one of the residues at -1 or +6 is a small residue;

(h) if base 3 in the quadruplet is C, then position +3 in the α -helix is Ser, Asp, Glu, Leu, Thr or Val;

(i) if base 2 in the quadruplet is G, then position -1 in the α -helix is Arg;

(j) if base 2 in the quadruplet is A, then position -1 in the α -helix is Gln;

(k) if base 2 in the quadruplet is T, then position -1 in the α -helix is Asn or Gln;

(l) if base 2 in the quadruplet is C, then position -1 in the α -helix is Asp;

(m) if base 1 in the quadruplet is G, then position +2 is Asp;

(n) if base 1 in the quadruplet is A, then position +2 is not Asp;

(o) if base 1 in the quadruplet is C, then position +2 is not Asp;

(p) if base 1 in the quadruplet is T, then position +2 is Ser or Thr.

23. The method of any of claims 18-22, further comprising the step of synthesizing a polynucleotide encoding the binding polypeptide.

24. A computer-implemented method for designing a zinc finger polypeptide, comprising the steps of:

(a) providing a system comprising at least storage means for storing data relating to a library of zinc fingers; storage means for storing a rule table; means for inputting target nucleic acid sequence data; processing means for generating a result; and means for outputting the result;

(b) inputting sequence data for a target nucleic acid molecule;

(c) defining a first target zinc finger binding site in said nucleic acid molecule;

(d) interrogating the zinc finger library and rule table storage means, comparing zinc fingers to the target zinc finger binding site according to the rule table and selecting zinc finger data identifying a zinc finger capable of binding to said target site;

(e) defining at least one further target zinc finger binding site and repeating step (d); and

(f) outputting the selected zinc finger data.

25. A method according to claim 24, further comprising sending instructions to an automated chemical synthesis system to assemble a zinc finger polypeptide as defined by the zinc finger data obtained in (f).

26. A method according to claim 25, wherein the zinc finger polypeptide is tested for binding to the target site, and data from said testing is used to select, from a plurality of candidates, a zinc finger polypeptide capable of binding to the target site.

27. A method according to any one of claims 24 to 26, wherein two or more zinc finger polypeptides are combined to form a zinc finger polypeptide capable of binding to an aggregate binding site comprising two or more target sites.

27. A method according to claim 24, wherein the rule table comprises rules as set forth in any one of claims 21 to 23.


FIGURE 1

FIGURE 2





FIGURE 3







FIGURE 4

FIGURE 5






FIGURE 6